

The Synthesis of Mycobactin Analogs and Heterocyclic Scaffolds From Acylnitroso Hetero-Diels-Alder Cycloadducts

Brian Scott Bodnar

**Publication Date** 

15-07-2008

## License

This work is made available under a All Rights Reserved license and should only be used in accordance with that license.

## Citation for this work (American Psychological Association 7th edition)

Bodnar, B. S. (2008). *The Synthesis of Mycobactin Analogs and Heterocyclic Scaffolds From Acylnitroso Hetero-Diels-Alder Cycloadducts* (Version 1). University of Notre Dame. https://doi.org/10.7274/2n49t15093d

This work was downloaded from CurateND, the University of Notre Dame's institutional repository.

For more information about this work, to report or an issue, or to preserve and share your original work, please contact the CurateND team for assistance at curate@nd.edu.

# THE SYNTHESIS OF MYCOBACTIN ANALOGS AND HETEROCYCLIC SCAFFOLDS FROM ACYLNITROSO HETERO-DIELS-ALDER CYCLOADDUCTS

VOLUME II

A Dissertation

Submitted to the Graduate School

of the University of Notre Dame

in Partial Fulfillment of the Requirements

for the Degree of

Doctor of Philosophy

by

Brian Scott Bodnar

Marvin J. Miller, Director

Graduate Program in Chemistry and Biochemistry

Notre Dame, Indiana

July 2008

# CONTENTS

# VOLUME II

8.1 General comments208.2 Experimental procedures for chapter 3208.3 Experimental procedures for chapter 4308.4 Experimental procedures for chapter 5348.5 Experimental procedures for chapter 6368.6 Experimental procedures for chapter 736	)8
8.2 Experimental procedures for chapter 3208.3 Experimental procedures for chapter 4308.4 Experimental procedures for chapter 5348.5 Experimental procedures for chapter 6368.6 Experimental procedures for chapter 738	)8
8.3 Experimental procedures for chapter 4308.4 Experimental procedures for chapter 5348.5 Experimental procedures for chapter 6368.6 Experimental procedures for chapter 738	)9
<ul> <li>8.4 Experimental procedures for chapter 5</li></ul>	)4
<ul> <li>8.5 Experimental procedures for chapter 6</li></ul>	43
8.6 Experimental procedures for chapter 7	57
	32
Appendix A	<del>)</del> 0
References	76

#### CHAPTER 8:

#### EXPERIMENTAL DATA

#### 8.1 General comments

All chemicals purchased were reagent grade and used without purification unless noted otherwise. Dry CH<sub>2</sub>Cl<sub>2</sub>, CH<sub>3</sub>CN, and Et<sub>3</sub>N were prepared by distillation from Dry THF was prepared by distillation from calcium hydride under Ar. sodium/benzophenone ketyl radical under Ar. Reactions were carried out under an inert atmosphere of argon only when specified in the experimental details, and were monitored by TLC as described in the experimental procedure using aluminum-backed 0.2 mm silica gel 60 F-254 plates. Visualization of TLC plates was performed under a UV lamp irradiating at 254 nm or by staining with CAM stain (Ceric Ammonium Molybdate stain, Hanessian's stain), ferric chloride stain, or ninhydrin stain. Column chromatography was conducted using silica gel 60 (230-400 mesh). All melting points were measured on a Thomas-Hoover Melting Apparatus and are uncorrected. All NMR spectra were recorded on a Varian 300 MHz or 500 MHz instrument under ambient temperatures unless otherwise noted. Chemical shift values for NMR spectra are reported as  $\delta$  in ppm relative to the solvent residual peak or to an internal tetramethylsilane standard. Infrared spectra were recorded using an FT-IR spectrometer and are reported in cm<sup>-1</sup>. Mass spectra were obtained as specified. Optical rotations were measured on a Rudolph Research Autopol III. Analytical LC/MS analyses were carried out on a Waters ZQ

instrument consisting of chromatography module Alliance HT, photodiode array detector 2996, and mass spectrometer Micromass ZQ, using a 3 x 50 mm Pro C18 YMC reverse phase column (Waters). Compounds were eluted using a gradient of 5-80% CH<sub>3</sub>CN in 10 mM ammonium acetate over 10 min at a flow rate of 0.7 mL/min. The MS electrospray source was operated at capillary voltage 3.5 kV and a desolvation temperature of 300  $^{\circ}$ C.

8.2 Experimental procedures for chapter 3



**2-Benzyloxy-benzoic acid (3.2).** Methyl salicylate (5.87 g, 5.0 mL, 38.6 mmol) was added to K<sub>2</sub>CO<sub>3</sub> (15.38 g, 111.3 mmol) and CH<sub>3</sub>CN (150 mL) at 20°C. The reaction was stirred at 60°C overnight. The reaction was filtered and concentrated to yield a clear oil. 10% aqueous KOH was added to the oil (65 mL) and the reaction was heated to 100°C for 2.5 h. The mixture was washed with CH<sub>2</sub>Cl<sub>2</sub> (2 x 100 mL), and acidified to an apparent pH of 2 using 1M HCl. The aqueous layer was extracted with EtOAc (3 x 100 mL) and the combined EtOAc layers were washed with H<sub>2</sub>O (2 x 100 mL) and brine (2 x 100 mL). The EtOAc was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to yield a white solid that was recrystallized from MeOH (8.54 g, 97%). mp 74-75°C; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  5.31 (s, 2H), 7.12-7.20 (m, 2H), 7.40-7.48 (m, 5H), 7.57 (ddd, *J* = 1.8, 7.5, 8.1 Hz, 1H), 8.22 (dd, *J* = 1.8, 7.8 Hz, 1H) ppm. HRMS (FAB) *m/z* [M+H]<sup>+</sup> calcd, 229.0865; obsd, 229.0841.



L-Serine benzyl ester hydrochloride (3.4). A 1-L round-bottomed flask was charged with Boc-L-serine (10.08 g, 49.11 mmol) and K<sub>2</sub>CO<sub>3</sub> (8.86 g, 64.1 mmol). DMF (400 mL) was added and the mixture was stirred under a stream of Ar for a few min. Benzyl bromide (6.20 mL, 52.2 mmol) was added to the reaction and the mixture was stirred vigourously at RT under Ar. After stirring overnight, the white solid was removed by vacuum filtration and washed with 200 mL of DMF. The filtrate was concentrated under vacuum and the residue was partitioned between  $H_2O$  (100 mL) and EtOAc (200 The layers were separated and the organic layer was washed with saturated mL). NaHCO<sub>3</sub> (3 x 100 mL) and brine (2 x 100 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated under vacuum to yield a yellow oil (14.159 g, 98% crude). The crude material was dissolved in 500 mL of anhydrous ether in a 1-L round-bottomed flask. The solution was cooled in an ice/ $H_2O$  bath to 4°C. Freshly prepared dry HCl gas was bubbled through the solution for 2h. (HCl gas was prepared by adding conc. HCl dropwise to anhydrous CaCl<sub>2</sub>, and the gas was then dried by bubbling through conc. H<sub>2</sub>SO<sub>4</sub>). The solution had turned opaque and the HCl gas flow was ceased. The opaque solution was stirred (with the attached septum) at RT overnight. The white solid was collected by vacuum filtration to yield only ~4 g (~35% yield). The filtrate was left to stand overnight to precipitate out more solid. A second crop of solid was collected to yield additional white solid. Compound 3.4 was isolated as a white solid (9.8439 g, 86.5% yield from Boc-serine).



(*S*)-Benzyl 2-(2-(benzyloxy)benzamido)-3-hydroxypropanoate (3.5) using EDC-mediated amide coupling. Dichloromethane (160 mL) was added to 3.4 (6.83 g, 29.5 mmol) and 3.2 (8.11 g, 35.5 mmol) under Ar. Et<sub>3</sub>N (4.80 mL, 34.5 mmol) was added followed by EDC•HCl (6.57 g, 34.3 mmol), and the mixture was stirred at rt overnight under Ar. The mixture was washed with  $H_2O$  (3 x 100 mL), saturated NaHCO<sub>3</sub> (3 x 100 mL), 5% aqueous citric acid (3 x 100 mL), and brine (3 x 100 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a white solid (11.2 g, 93.3% yield).

**Compound 3.5 from the acid chloride.** Compound **3.2** (13.21 g, 56.72 mmol) was dissolved in 150 mL of anhydrous  $CH_2Cl_2$  in a 500-mL round-bottomed flask. Oxalyl chloride (9.95 mL, 114 mmol) was added to the colorless solution slowly, followed by a catalytic amount of anhydrous DMF (0.10 mL, 1.3 mmol). The light yellow solution bubbled profusely and was stirred at RT under Ar. Bubbling ceased after ~2h, and the reaction became yellow in color. After 4h, the reaction was concentrated under vacuum, then dissolved in toluene and concentrated under vacuum (2x), then dissolved in CHCl<sub>3</sub> and concentrated under vacuum (2x) to yield the crude acid chloride.

The crude acid chloride (8.59 g, 34.1 mmol) was dissolved in 170 mL of anhydrous  $CH_2Cl_2$  in a 500-mL round-bottomed flask under Ar. **3.4** (8.39 g, 36.2 mmol) was added to the flask and the mixture cooled in an ice/H<sub>2</sub>O bath under Ar. Diisopropylethylamine (14.9 mL, 85.6 mmol) was added to the reaction slowly. Most of the solid material dissolved, and the light orange solution was warmed to RT and stirred

under Ar overnight. After 19h, TLC of the reaction (90% CH<sub>2</sub>Cl<sub>2</sub>/acetone) indicated the reaction was complete. The orange solution was concentrated under vacuum to yield an orange oil. The oil was partitioned between EtOAc (~300 mL) and H<sub>2</sub>O (~150 mL). The layers were separated and the organic layer was washed with H<sub>2</sub>O (1 x 100 mL), 5% citric acid (3 x 100 mL), H<sub>2</sub>O (1 x 100 mL), saturated NaHCO<sub>3</sub> (3 x 100 mL), H<sub>2</sub>O (1 x 100 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to yield **3.5** as a fluffy white solid (13.5 g, 97.8% yield). mp 119-120°C; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 1.76 (br, 1H, -OH), 3.92 (m, *J* = 3.9 Hz, 1H, -NH), 4.89 (dt, *J* = 3.9, 6.9 Hz, 1H, α-CH), 5.12-5.30 (m, 4H, OCH<sub>2</sub>-Ph), 7.05 (d, *J* = 8.4 Hz, 1H, Ar-H), 7.11 (d, *J* = 7.8 Hz, 1H, Ar-H), 7.30-7.48 (m, 10H, Ar-H), 8.20 (dd, *J* = 1.5, 7.8 Hz, 1H, Ar-H), 8.80 (d, *J* = 6.6 Hz, 1H, Ar-H) ppm; <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 55.62, 63.79, 67.39, 71.47, 113.0, 121.3, 121.7, 128.2, 128.3, 128.6, 128.8, 129.0, 132.5, 133.4, 135.5, 135.7, 157.2, 165.8, 170.4 ppm; HRMS-FAB (m/z) [M + H]<sup>+</sup> found 406.

**PEG-supported Burgess's Reagent (3.7).** Polyethylene glycol (PEG) (6.96 g, 9.29 mmol) was dried in vacuo for 4 h and added dropwise as a solution in 20 mL of benzene to a solution of chlorosulfonyl isocyanate (1.20 g, 0.74 mL, 8.47 mmol) in 20 mL of benzene at RT. The reaction was allowed to stir for an additional 1h before concentrating to yield a yellow-tan oil. The oil was stored under vacuum overnight. The oil was dissolved in 20 mL of benzene and added dropwise to a solution of Et<sub>3</sub>N (1.97 g,

2.74 mL, 19.5 mmol) in 15 mL of benzene at RT. The reaction was stirred for an additional 15 min, filtered, and concentrated to afford a tan solid (8.14 g). mp ~ 20-25 °C.



(S)-Benzyl 2-(2-(benzyloxy)phenyl)-4,5-dihydrooxazole-4-carboxylate (3.8) using PEG-Burgess reagent. Compound 3.5 (0.59 g, 1.46 mmol) and compound 3.7 (2.8 g, 2.8 mmol) were dissolved in 10 mL 1:1 THF/dioxane at RT. The reaction was heated to 95 °C for 3 h. The solvent was removed to yield a yellow oil. Chromatography through silica gel using a solvent system of 95%  $CH_2Cl_2/EtOAc$  provided 3.8 as a white solid (0.428 g ,76%).

General procedure for the preparation of oxazolines using DAST. Oxazoline 3.8 using DAST. Amide 3.5 (3.27 g, 8.07 mmol) was dissolved in 100 mL of  $CH_2Cl_2$  under Ar. The solution was cooled to -78 °C (dry ice/acetone bath) and DAST (1.20 mL, 9.16 mmol) was added dropwise over 4 min. The reaction was stirred at -78 °C for 3.5 h. The reaction was monitored by TLC (1:1 hexanes/EtOAc – UV lamp) for the disappearance of amide 3.5. K<sub>2</sub>CO<sub>3</sub> (3.03 g, 21.9 mmol) was added to the reaction in one portion and the mixture was allowed to reach RT over 45 min. The mixture was poured into 100 mL of saturated NaHCO<sub>3</sub> and the layers were separated. The aqueous layer was extracted with  $CH_2Cl_2$  (3 x 50 mL). The combined organic layers were washed with  $H_2O$  (2 x 100 mL) and brine (2 x 100 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a yellow oil that solidified in the freezer overnight. Recrystallized from MeOH to

yield oxazoline **3.8** as a white solid (2.87 g, 91.7%). mp = 69-70 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz)  $\delta$  4.55-5.70 (m, 2H), 5.03 (dd, *J* = 7.9, 10.6 Hz, 1H), 5.20-5.32 (m, 4H), 7.00 (m, 2H), 7.27-7.44 (m, 10H), 7.50 (dd, *J* = 0.9, 8.1 Hz, 2H), 7.82 (ddd, *J* = 0.6, 1.8, 8.1 Hz, 1H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  171.0, 165.7, 157.6, 136.8, 135.5, 132.6, 131.6, 128.5, 128.4, 128.3, 128.2, 127.5, 126.7, 120.7, 117.2, 113.8, 70.6, 69.2, 68.8, 67.1 ppm.  $[\alpha]_D^{23} = +105$  ° (c = 1, MeOH). HRMS (FAB) *m*/*z* [M+H]<sup>+</sup> calcd, 388.1549; obsd, 388.1549.



(*S*)-2-(2-Hydroxy-phenyl)-4,5-dihydro-oxazole-4-carboxylic acid (2.62). To an Ar-purged solution of **3.8** (2.00 g, 5.16 mmol) in MeOH (100 mL) was added 10 wt% Pd/C catalyst (3.75 mg, 18 wt %). The solution was stirred under H<sub>2</sub> (balloon) for 3.5h. The reaction was filtered through celite and the filtrate was concentrated and triturated from EtOAc/hexanes to yield **2.62** as a white solid (1.06 g, 99%). mp 149-150 °C (turns pink). <sup>1</sup>H NMR (500 MHz, d<sup>6</sup>-DMSO, 30 °C)  $\delta$  7.63 (dd, *J* = 8.0, 2.0 Hz, 1H), 7.46 (ddd, *J* = 9.0, 7.0, 1.5 Hz, 1H), 7.00 (dd, *J* = 8.5, 1.0 Hz, 1H), 6.94 (ddd, *J* = 8.0, 7.5, 1.0 Hz, 1H), 5.02 (dd, *J* = 10.0, 7.5 Hz, 1H), 4.63 (m, 2H) ppm. <sup>13</sup>C NMR (125 MHz, d<sup>6</sup>-DMSO, 30 °C)  $\delta$  171.6, 166.1, 159.1, 134.0, 127.9, 119.0, 116.5, 109.6, 69.2, 66.7 ppm. HRMS (FAB) *m*/*z* calcd for C<sub>10</sub>H<sub>9</sub>NO<sub>4</sub>, 208.0610; obsd, 208.0609.



(S)-Benzyl 2-benzamido-3-hydroxypropanoate (3.9). Compound 3.4 (8.018 g, 34.609 mmol) and benzoic acid (4.692 g, 38.42 mmol) were added to a flame-dried 500mL round-bottomed flask along with 150 mL of dry CH<sub>2</sub>Cl<sub>2</sub> under Ar. Et<sub>3</sub>N (5.20 mL, 37.0 mmol) was added to the suspension, followed by EDC•HCl (7.79 g, 40.6 mmol). After stirring for 2h under Ar at RT, the suspended material had fully dissolved and the resultant colorless solution was stirred at RT under Ar overnight. After ~26 h, the reaction was concentrated and the resultant orange oil was partitioned between EtOAc (300 mL) and  $_{\text{H2O}}$  (200 mL). The layers were separated and the aqueous layer was extracted with EtOAc (2 x 150 mL). The combined organic layers were washed with H<sub>2</sub>O (200 mL), 5% citric acid (3 x 200 mL), H<sub>2</sub>O (1 x 200 mL), saturated NaHCO<sub>3</sub> (3 x 200 mL), H<sub>2</sub>O (1 x 200 mL), and brine (2 x 200 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to yield **3.9** as a white solid (9.94 g, 95.9% yield). mp = 91-92 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  3.36 (t, J = 5.4 Hz, 1H), 3.90 – 4.05 (m, 2H), 4.82 (dt, J =7.5, 3.3 Hz, 1H), 5.15 (s, 2H), 7.29 – 7.34 (m, 7H), 7.42 (t, J = 7.5 Hz, 1H), 7.74 (d, J = 7.5 Hz, 2H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 55.48, 63.41, 127.19, 128.08, 128.44, 128.62, 131.76, 133.93, 135.39, 167.74, 170.44 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>17</sub>H<sub>17</sub>NO<sub>4</sub>, 300.1236; obsd, 300.1231.



(S)-Benzyl 2-phenyl-4,5-dihydrooxazole-4-carboxylate (3.10). Compound 3.9 (5.123 g, 17.12 mmol) was dissolved in 100 mL of CH<sub>2</sub>Cl<sub>2</sub> in a flame-dried 250-mL round-bottomed flask and cooled to -78°C in a dry ice/acetone bath under Ar. Diethylaminosulfur trifluoride (2.50 mL, 18.9 mmol) was added dropwise to the reaction over  $\sim 5$  min. The reaction was stirred under Ar at  $-78^{\circ}$ C. After 4 h, the reaction was complete by TLC (1:1 hexanes/EtOAc - UV lamp), and  $K_2CO_3$  (6.31 g, 45.7 mmol) was added to the reaction in one portion. The reaction was removed from the dry ice/acetone bath and allowed to warm to RT under Ar over ~30 min while stirring. The solution was poured into  $\sim 100$  mL of saturated NaHCO<sub>3</sub> and H<sub>2</sub>O was added until all of the solid material had dissolved. The layers were separated and the aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 x 60 mL). The combined CH<sub>2</sub>Cl<sub>2</sub> layers were washed with saturated NaHCO<sub>3</sub> (1 x 80 mL), H<sub>2</sub>O (2 x 80 mL), and brine (1 x 80 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to yield a yellow oil that solidified upon storage at -10°C overnight to a white solid (4.96 g). Recrystallization from EtOAc/hexanes afforded oxazoline **3.10** as a white solid (4.103 g, 85.2% yield). mp = 49-50 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.99 (m, 2H), 7.50 (m, 1H), 7.42-7.32 (m, 7H), 5.28 (d, J = 12 Hz, 1H), 5.22 (d, J = 12 Hz, 1H), 4.98 (dd, J = 10.5, 8.0 Hz, 1H), 4.67 (t, J = 8.5 Hz, 1H), 4.60 (dd, J = 10.5, 8.5 Hz, 1H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  170.9, 166.3, 135.3, 131.8, 128.49, 128.47, 128.31, 128.26, 128.2, 126.8, 69.5, 68.6, 67.2 ppm.



(*S*)-2-Phenyl-4,5-dihydrooxazole-4-carboxylic acid (3.11). To an Ar-purged solution of 3.10 (1.71 g, 6.07 mmol) in MeOH (50 mL) was added 10 wt% Pd/C catalyst (0.283 g, 17 wt %). The solution was stirred under H<sub>2</sub> (balloon) for 2.5h. The reaction was filtered through celite and the filtrate was concentrated and triturated from EtOAc/hexanes to yield 3.11 as an off-white solid (1.16 g, 99%). mp 128-131 °C (turns red at 125 °C). <sup>1</sup>H NMR (300 MHz, d<sup>6</sup>-DMSO)  $\delta$  7.89 (d, *J* = 7.2 Hz, 2H), 7.58 (m, 1H), 7.49 (m, 2H), 4.86 (m, 1H), 4.62-5.51 (m, 2H) ppm.



*D*-Serine benzyl ester hydrochloride (3.12). Compound 3.12 was prepared following the same procedure used for compound 3.4. Boc-*D*-serine (2.00 g, 9.77 mmol) afforded 3.12 as a white solid (2.10 g, 93%).



(*R*)-Benzyl 2-benzamido-3-hydroxypropanoate (3.13). Compound 3.13 was prepared following the same procedure used for compound 3.9. Benzoic acid (0.250 g, 2.05 mmol) and 3.12 (0.401 g, 1.73 mmol) provided 3.13. Recrystallization from EtOAc/hexanes provided 3.13 as a white solid (0.386 g, 75%). mp = 103-104 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.80 (d, *J* = 7.5 Hz, 1H), 7.49 (m, 1H), 7.34 - 7.41 (m, 5H),

7.24 (d, J = 7.5 Hz, 1H), 5.22 (s, 2H), 4.88 (m, 1H), 4.07 (dd, J = 11.1, 3.6 Hz, 1H), 4.00 (dd, J = 11.4, 3.3 Hz, 1H), 3.05 (br-s, 1H) ppm. <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  170.4, 167.7, 135.1, 133.4, 131.9, 128.60, 128.56, 128.47, 128.1, 127.1, 67.5, 63.3, 55.2 ppm.



(*R*)-Benzyl 2-(2-(benzyloxy)benzamido)-3-hydroxypropanoate (3.14). Compound 3.14 was prepared following the same procedure used for compound 3.5 using EDC-mediated amide coupling. 3.2 (0.476 g, 2.09 mmol) and 3.12 (0.401 g, 1.73 mmol) provided 3.14. Recrystallization from EtOAc/hexanes yielded amide 3.14 as a white solid (0.559 g, 80%). mp = 118-120 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  8.86 (d, *J* = 6.9 Hz, 1H), 8.19 (dd, *J* = 8.1, 1.8 Hz, 1H), 7.32 – 7.46 (m, 10H), 7.02 – 7.10 (m, 2H), 5.17 (s, 2H), 5.13 (m, 2H), 4.88 (m, 1H), 3.91 (m, 2H), 2.25 (br-s, 1H) ppm. <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  170.1, 165.6, 156.9, 135.5, 135.3, 133.2, 132.3, 128.7, 128.3, 128.1, 128.0, 121.5, 112.8, 71.3, 67.2, 63.6, 55.4 ppm.



(*R*)-Benzyl 2-phenyl-4,5-dihydrooxazole-4-carboxylate (3.15). Compound 3.15 was prepared following the same procedure used for oxazoline 3.10. 3.13 (0.204 g, 0.682 mmol) and DAST (0.110 mL, 0.839 mmol) provided oxazoline 3.15 as a colorless oil (0.166 g, 86%).



(*R*)-Benzyl 2-(2-(benzyloxy)phenyl)-4,5-dihydrooxazole-4-carboxylate (3.16). Compound 3.16 was prepared following the same procedure used for oxazoline 3.10. 3.14 (0.301 g, 0.743 mmol) and DAST (0.110 mL, 0.839 mmol) provided oxazoline 3.16. Recrystallization from MeOH yielded 3.16 as a white solid (0.201 g, 70%). mp = 60.5-62.5 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.82 (dd, *J* = 7.8, 1.8 Hz, 1H), 7.50 (m, 2H), 7.27 – 7.43 (m, 9H), 6.99 (m, 2H), 5.25 (m, 2H), 5.19 (s, 2H), 5.01 (dd, *J* = 10.5, 7.8 Hz, 1H), 4.53 – 4.69 (m, 2H) ppm.



**Benzyl 2-(2-(benzyloxy)phenyl)oxazole-4-carboxylate (3.17).** Compound **3.8** (0.101 g, 0.261 mmol) was dissolved in 2.6 mL of  $CH_2Cl_2$  in a flame-dried 10-mL flask under Ar. The solution was cooled in a dry ice/acetone bath maintained at -25 to -20 °C. Bromotrichloromethane (0.186 g, 0.937 mmol) was added dropwise to the reaction followed by 1,8-diazabicyclo[5.4.0]undec-7-ene (0.140 mL, 0.936 mmol). The reaction was stirred for 5 min in the dry ice/acetone bath, then warmed to 4 °C in an ice/H<sub>2</sub>O bath and stirred for 2.5 h under Ar. The reaction was complete by TLC (3:2 hexanes/EtOAC – UV lamp) and the reaction was warmed to RT. The mixture was quenched with saturated NaHCO<sub>3</sub>, then poured into 10 mL of saturated NaHCO<sub>3</sub> and diluted with 12 mL of EtOAc. The layers were separated and the aqueous layer was extracted with EtOAC (3 x 10 mL). The combined organic layers were washed with brine (10 mL), 1M HCl (2 x 10

mL), brine (10 mL), 0.5% NaOCl (2 x 10 mL), and brine (2 x 10 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield an oil residue. Chromatography through 13 g of silica using 100% CH<sub>2</sub>Cl<sub>2</sub> provided oxazole **3.17** as a colorless oil (70.6 mg, 70%). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  8.32 (s, 1H), 8.07 (dd, *J* = 8.1, 1.8 Hz, 1H), 7.53 (m, 2H), 7.29 – 7.49 (m, 9H), 7.06 (m, 2H), 5.41 (s, 2H), 5.22 (s, 2H) ppm. <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  161.4, 161.3, 156.7, 143.9, 135.6, 133.9, 132.4, 131.1, 128.54, 128.46, 128.3, 127.7, 126.8, 121.0, 116.2, 113.5, 70.5, 66.6 ppm.



(*S*)-Methyl 2-(2-(benzyloxy)benzamido)-3-hydroxypropanoate (3.19). Compound 3.19 was prepared following the same procedure used for compound 3.5 using EDC-mediated amide formation. 3.18 (3.15 g, 20.3 mmol), 3.2 (5.55 g, 24.3 mmol), Et<sub>3</sub>N (3.33 mL, 24.0 mmol), and EDC•HCl (4.50 g, 23.5 mmol) provided amide 3.19 as a white solid (6.19 g, 93%). mp = 127-128 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$ 8.78 (d, *J* = 7.0 Hz, 1H), 8.18 (dd, *J* = 7.5, 1.5 Hz, 1H), 7.48 – 7.35 (m, 5H), 7.06 – 7.03 (m, 2H), 5.22 (m, 2H), 4.82 (m, 1H), 3.88 (m, 2H), 3.68 (s, 3H) ppm.



(S)-Methyl 2-(2-(benzyloxy)phenyl)-4,5-dihydrooxazole-4-carboxylate (3.20). Prepared following the same procedure used for oxazoline 3.10. 3.19 (6.78 g, 20.6 mmol) and DAST (3.00 mL, 22.7 mmol) provided oxazoline 3.20. Recrystallization from MeOH provided shiny, off-white crystals (5.57 g, 87%). mp = 121-123 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.81 (dd, J = 13, 2.5 Hz, 1H), 7.50 (m, 2H), 7.42 – 7.28 (m, 4H), 7.01 – 6.95 (m, 2H), 5.18 (s, 2H), 4.97 (dd, J = 17.5, 13.5 Hz, 1H), 4.67 (t, J = 13.5 Hz, 1H), 4.57 (dd, J = 17.5, 14.5 Hz, 1H), 3.80 (s, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  171.7, 165.6, 157.6, 136.8, 132.6, 131.6, 128.3, 127.5, 126.7, 120.7, 117.2, 113.7, 70.6, 69.1, 68.6, 52.5 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>18</sub>H<sub>18</sub>NO<sub>4</sub><sup>+</sup>, 312.1236; obsd, 312.1247.



#### (S)-2-(2-(Benzyloxy)phenyl)-N-hydroxy-4,5-dihydrooxazole-4-carboxamide

(3.21). A methanolic solution of KOH (400 mg, 7.14 mmol) in 5 mL of MeOH (~1.4M) was slowly added to a solution of hydroxylamine hydrochloride (242 mg, 3.48 mmol) in 2.5 mL of MeOH at 0 °C. A solution of **3.20** (509 mg, 1.63 mmol) in 10 mL of MeOH was added to the reaction and the reaction was stirred for 3h at 0 °C and then stored at -10 °C overnight. The reaction was acidified to an apparent pH of 4 through the dropwise addition of 1M HCl. EtOAc and H<sub>2</sub>O were added to the reaction and the biphasic solution was extracted with EtOAc (3 x 30 mL). The combined EtOAc layers were washed with H<sub>2</sub>O (2 x 40 mL) and brine (3 x 40 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield hydroxamate **3.21** as a white solid (108 mg, 22%). mp = 115-120 °C (dec.). <sup>1</sup>H NMR (300 MHz, d<sup>6</sup>-DMSO)  $\delta$  4.45 (t, *J* = 8.1 Hz, 1H), 4.52 (t, *J* = 8.1 Hz, 1H), 4.67 (dd, *J* = 9.5, 8.4 Hz, 1H), 5.22 (s, 2H), 7.02 (t, *J* = 7.3 Hz, 1H), 7.19 (d, *J* = 8.1

Hz, 1H), 7.27-7.52 (m, 6H), 7.72 (dd, J = 7.5, 1.3 Hz, 1H), 9.02 (s, 1H), 10.77 (s, 1H) ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>17</sub>H<sub>17</sub>N<sub>2</sub>O<sub>4</sub><sup>+</sup>, 313.1188; obsd, 313.1198.

General procedure for hydroxamate formation from esters using Me<sub>3</sub>Al and hydroxylamines. Attempted formation of hydroxamate 3.22 from ester 3.20 (Table 3.7, entry 6). *O*-allylhydroxylamine hydrochloride (0.160 g, 1.46 mmol) was suspended in 5 mL of CH<sub>2</sub>Cl<sub>2</sub> and cooled under Ar in an ice/H<sub>2</sub>O bath. Trimethylaluminum (2.0M in heptane, 0.740 mL, 1.48 mmol) was added dropwise and the resultant mixture was warmed to RT and stirred for 1 h, at which time the solid material was fully in solution. The solution was cooled in the ice/H<sub>2</sub>O bath again as a solution of ester 3.20 (0.250 g, 0.800 mmol) in 5 mL of CH<sub>2</sub>Cl<sub>2</sub> was added dropwise to the reaction. The reaction was allowed to reach RT and stirred overnight under Ar. 10% Citric acid (5 mL) was added dropwise to the reaction and the mixture was stirred at RT for 1 h. The layers were separated and the aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 x 15 mL). The combined organic layers were washed with 10% citric acid (2 x 25 mL), H<sub>2</sub>O (20 mL), saturated NaHCO<sub>3</sub> (2 x 30 mL), and brine (2 x 40 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated.



*N*-Benzyloxy-phthalimide (3.25). Compound 3.25 was prepared following the procedure outlined by Welch and Seper.<sup>1</sup> *N*-hydroxyphthalimide (70.0 g, 429 mmol) and benzyl bromide (55.0 mL, 463 mmol) were dissolved in 550 mL of  $CH_3CN$ . The solution

was warmed and Et<sub>3</sub>N (47.3 g, 65.0 mL, 467 mmol) was added slowly to the solution. The mixture was heated at reflux for 6 h until all of the solid material dissolved. The reaction was cooled to RT and poured into 300 mL of H<sub>2</sub>O and extracted with EtOAc (3 x 500 mL). The combined organic layers were dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a yellow solid. Recrystallization from MeOH provided **3.25** as pale yellow crystals (87.4 g, 80%). mp 142-143°C (lit.<sup>1</sup> 141-142°C). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.82 - 7.70 (m, 4H), 7.53 (m, 2H), 7.37 (m, 3H), 5.21 (s, 2H) ppm.



*O*-Benzylhydroxylamine hydrochloride, OBHA+HCl (3.27). EtOH (400 mL) was added to compound 3.25 (20.0 g, 79.0 mmol) in a 1-L round-bottomed flask. The mixture was heated to 40 °C in an oil bath and anhydrous hydrazine (2.90 mL, 92.4 mmol) was added. After 1 min, most of the solid material was in solution. The flask was equipped with a jacketed condenser and the yellow solution was heated at reflux overnight (oil bath temp at 100 °C). After a few min, a white solid started to precipitate out of the solution. After heating to reflux for 18 h, the reaction was allowed to cool to RT and stirred for an additional 7 h. The mixture was filtered under vacuum to afford the by-product 3.26, as a white solid (12.7 g, 99%) and a yellow filtrate. The filtrate was concentrated under vacuum to afford a yellow solid that was partitioned between  $CH_2Cl_2$  (400 mL) and 3M NaOH (300 mL). The layers were separated and the  $CH_2Cl_2$  layer was washed with 3M NaOH (2 x 250 mL), H<sub>2</sub>O (250 mL), and brine (250 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to yield free *O*-benzylhydroxylamine as a thin yellow oil (9.3 g, ~96%).

The free amine was dissolved in 500 mL of anhydrous Et<sub>2</sub>O. HCl gas (prepared by adding conc. HCl to a round-bottomed flask containing anhydrous CaCl<sub>2</sub>, then bubbling through conc. H<sub>2</sub>SO<sub>4</sub> to dry) was bubbled through the solution for 10 min. White solid began to precipitate out of solution immediately. After standing for a few days, HCl was bubbled through the solution for an additional 45 min (more solid precipitated out during this time) in order to saturate the solution with an excess of HCl. The white solid was collected via vacuum filtration. The remaining amount of solid in the filtrate was collected via vacuum filtration through a coarse glass frit and provided **3.27** as a white solid (10.3 g, 82%). mp = 238 °C (sublimes). <sup>1</sup>H NMR (300 MHz, d<sup>6</sup>-DMSO)  $\delta$  11.25 (s, 2H), 7.41 (s, 5H), 5.06 (s, 2H) ppm. <sup>13</sup>C NMR (75 MHz, d<sup>6</sup>-DMSO)  $\delta$  133.6, 129.2, 129.0, 128.6, 75.6 ppm.



General procedure for hydroxamate formation under aqueous conditions using EDC•HCl. (S)-N-(Benzyloxy)-2-(2-hydroxyphenyl)-4,5-dihydrooxazole-4carboxamide (3.28). Carboxylic acid 2.62 (0.105 g, 0.508 mmol) and 3.27 (0.245 g, 1.54 mmol) were dissolved in 5 mL of 3:2 THF/H<sub>2</sub>O. The apparent pH of the mixture was adjusted to 4.5 using a dilute aqueous solution of NaOH. EDC•HCl (0.162 g, 0.845 mmol) was added to the reaction in portions and the mixture was stirred at RT, maintaining an apparent pH of 4.5 by adding aqueous HCl to the reaction mixture. The reaction was complete when adding portions of EDC•HCl no longer had a significant effect on the apparent pH of the reaction mixture. The mixture was diluted with H<sub>2</sub>O (20 mL) and extracted with EtOAc (3 x 15 mL). The combined organic layers were washed with 5% citric acid (3 x 15 mL) and brine (3 x 15 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a white solid (94% crude yield). Chromatography through 15 g of silica using 95:5 CH<sub>2</sub>Cl<sub>2</sub>/EtOAc provided hydroxamate **3.28** as a white solid (130.7 mg, 83%). mp = 178-180 °C (turns red, then melts). Rf = 0.302 (1:1 hexanes/EtOAc – UV lamp). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  11.15 (br-s, 1H), 8.93 (s, 1H), 7.66 (dd, *J* = 7.5, 1.5 Hz, 1H), 7.42 – 7.31 (m, 6H), 6.99 (dd, *J* = 8.5, 0.5 Hz, 1H), 6.90 (m, 1H), 4.93 (s, 2H), 4.88 (dd, *J* = 11, 8.0 Hz, 1H), 4.68 (m, 1H), 4.61 (dd, *J* = 11, 9.0 Hz, 1H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  168.0, 167.7, 159.5, 134.4, 129.3, 128.9, 128.6, 128.5, 119.2, 116.8, 109.8, 78.5, 69.3, 66.8 ppm. HRMS (FAB) *m*/*z* [M+H]<sup>+</sup> calcd for C<sub>17</sub>H<sub>17</sub>N<sub>2</sub>O<sub>4</sub><sup>+</sup>, 313.1188; obsd, 313.1215.



(*S*)-*N*-(**Benzyloxy**)-2-phenyl-4,5-dihydrooxazole-4-carboxamide (3.29). Compound 3.29 was prepared following the same procedure used for the preparation of hydroxamate 3.28. Carboxylic acid 3.11 (0.169 g, 0.882 mmol), 3.27 (0.422 g, 2.64 mmol) and EDC•HCl (0.262 g, 1.37 mmol) provided hydroxamate 3.29. Chromatography through 15 g of silica using 4:1 CH<sub>2</sub>Cl<sub>2</sub>/EtOAc yielded 3.29 as a white solid (0.103 g, 39%). mp = 104-106 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  9.38 (s, 1H), 7.85 (d, *J* = 7.0 Hz, 2H), 7.50 (td, *J* = 7.5, 1.0 Hz, 1H), 7.40 – 7.30 (m, 7H), 4.92 (s, 2H), 4.77 (dd, *J* = 11.0, 8.0 Hz, 1H), 4.68 (t, *J* = 8.0 Hz, 1H), 4.62 (dd, *J* = 11.0, 9.0 Hz, 1H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  168.7, 166.5, 134.8, 132.1, 129.25, 129.21, 128.8, 128.5, 128.4, 128.0, 126.5, 78.4, 70.1, 67.8 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for  $C_{17}H_{17}N_2O_3^+$ , 297.1239; obsd, 297.1219.



*N*-(Benzyloxy)-2-phenyloxazole-4-carboxamide (3.31). Compound 3.31 was prepared following the same procedure used for the preparation of hydroxamate 3.28. Carboxylic acid 3.30 (0.492 g, 2.60 mmol), 3.27 (0.540 g, 3.38 mmol), and EDC•HCl (0.615 g, 3.21 mmol) provided hydroxamate 3.31. Chromatography through 50 g of silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 90% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc yielded 3.31 as a white solid (0.506 g, 66%). mp = 140-142 °C. Rf = 0.21 (95:5 CH<sub>2</sub>Cl<sub>2</sub>/EtOAc – UV lamp). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  9.46 (br-s, 1H), 8.30 (s, 1H), 7.97 (m, 2H), 7.46-7.38 (m, 8H), 5.06 (s, 2H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  161.6, 158.6, 141.4, 135.1, 135.0, 131.2, 129.2, 128.84, 128.77, 128.6, 126.6, 126.2, 78.8 ppm. HRMS (FAB) *m*/*z* [M+H]<sup>+</sup> calcd for C<sub>17</sub>H<sub>15</sub>N<sub>2</sub>O<sub>3</sub><sup>+</sup>, 295.1083; obsd, 295.1073.



#### (S)-N-Hydroxy-2-(2-hydroxyphenyl)-4,5-dihydrooxazole-4-carboxamide

(3.32). All glassware was washed with 6M HCl, then rinsed with  $H_2O$  and acetone before using in order to remove residual metals. MeOH (25 mL) was added to 3.28 (0.487 g, 1.56 mmol) in a 100-mL round-bottomed flask under Ar. The solid did not dissolve completely. EtOAc was added to the mixture until the solid completely

dissolved (~8 mL). The solution was purged with Ar for 10 min and 10 wt% Pd/C (52 mg, ~11 wt%) was added to the solution. The reaction was purged with Ar for 5 min, then H<sub>2</sub> for 5 min, then stirred under H<sub>2</sub> (balloon) for 2 h, after which time a more polar, FeCl<sub>3</sub>-positive spot was observed by TLC analysis. The reaction was purged with Ar for 10 min, then filtered through a small amount of celite. The filtrate was concentrated to yield a white solid (336 mg, 98%). mp = 154-155 °C. Rf = 0.42 (9:1 CH<sub>2</sub>Cl<sub>2</sub>/MeOH-UV lamp, FeCl<sub>3</sub> stain). <sup>1</sup>H NMR (300 MHz, d<sup>6</sup>-DMSO)  $\delta$  11.69 (s, 1H), 11.04 (s, 1H), 9.11 (s, 1H), 7.63 (dd, *J* = 7.8, 1.5 Hz, 1H), 7.46 (ddd, *J* = 9.0, 7.5, 1.8 Hz, 1H), 7.01 – 6.92 (m, 2H), 4.81 (dd, *J* = 9.9, 7.5 Hz, 1H), 4.64 – 4.51 (m, 2H) ppm. <sup>13</sup>C NMR (125 MHz, d<sup>6</sup>-DMSO)  $\delta$  166.2, 165.8, 158.9, 133.9, 127.9, 118.9, 116.4, 109.7, 68.9, 65.3 ppm. HRMS (FAB) *m*/z [M+H]<sup>+</sup> calcd for C<sub>10</sub>H<sub>11</sub>N<sub>2</sub>O<sub>4</sub><sup>+</sup>, 223.0719; obsd, 223.0711.



(*S*)-2-Benzamido-3-hydroxypropanoic acid (3.33). Compound 3.33 was prepared following the same procedure used for the preparation of 2.62. Compound 3.9 (0.725 g, 2.42 mmol) provided 3.33 as a white foam (0.516 g, 99% yield). <sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD)  $\delta$  7.87 (m, 2H), 7.45 – 7.58 (m, 3H), 4.72 (t, *J* = 4.2 Hz, 1H), 3.94 – 4.06 (m, 2H) ppm. <sup>13</sup>C NMR (75 MHz, CD<sub>3</sub>OD)  $\delta$  173.5, 170.2, 135.3, 133.0, 129.6, 128.5, 63.0, 56.8 ppm.



(*S*)-3-Hydroxy-2-(2-hydroxybenzamido)propanoic acid (3.34). Compound 3.34 was prepared following the same procedure used for the preparation of 2.62. Compound 3.5 (1.41 g, 3.47 mmol) provided 3.34 as a light purple foam (0.784 g, 99%). <sup>1</sup>H NMR (500 MHz, d<sup>6</sup>-DMSO, 30 °C)  $\delta$  11.85 (br, 1H), 8.93 (d, *J* = 7.5 Hz, 1H), 7.95 (dd, *J* = 7.5, 1.0 Hz, 1H), 7.39 (m, 1H), 6.93 (m, 2H), 4.53 (dt, *J* = 6.5, 4.5 Hz, 1H), 3.86 (dd, *J* = 16.0, 5.0 Hz, 1H), 3.80 (dd, *J* = 11.0, 4.0 Hz, 1H) ppm. <sup>13</sup>C NMR (125 MHz, d<sup>6</sup>-DMSO, 30 °C)  $\delta$  171.7, 167.1, 158.3, 133.4, 129.2, 118.9, 117.1, 116.5, 61.1, 55.0 ppm. HRMS (FAB) *m*/z [M+H]<sup>+</sup> calcd for C<sub>10</sub>H<sub>12</sub>NO<sub>5</sub><sup>+</sup>, 226.0715; obsd, 226.0717.



(*S*)-N-(1-(Benzyloxyamino)-3-hydroxy-1-oxopropan-2-yl)benzamide (3.35). Compound 3.35 was prepared following the same procedure used for the preparation of hydroxamate 3.28. Carboxylic acid 3.33 (58 mg, 0.28 mmol), 3.27 (51 mg, 0.32 mmol), and EDC•HCl (65 mg, 0.34 mmol) provided 3.35 as a white solid (62 mg, 71%).



(S)-N-(1-(Benzyloxyamino)-3-hydroxy-1-oxopropan-2-yl)-2-

hydroxybenzamide (3.36). Compound 3.36 was prepared following the same procedure

used for the preparation of hydroxamate **3.28**. Carboxylic acid **3.34** (74 mg, 0.33 mmol), **3.27** (57 mg, 0.36 mmol), and EDC•HCl (73 mg, 0.38 mmol) provided **3.36** as a fluffy white solid (77 mg, 71%).



*tert*-Butyl *N*-hydroxycarbamate (3.37). Hydroxylamine hydrochloride (10.00 g, 144.0 mmol) was suspended in 240 mL of THF and 60 mL of H<sub>2</sub>O in a 1-L flask. NaHCO<sub>3</sub> (24.25 g, 288.7 mmol) was added (some bubbling was observed). The biphasic mixture (aq. layer was cloudy) was stirred vigorously at RT for 5-10 min. Di*-tert*-butyldicarbonate (33.0 g, 151 mmol) was added in one portion (bubbling observed), and the mixture was stirred vigorously at RT for 3.75 h. The mixture was diluted with 200 mL of H<sub>2</sub>O and 150 mL of EtOAc (most of the solid dissolved), and the aqueous layer was acidified to an apparent pH of 4 (pH paper) by adding a 10 wt% solution of citric acid (bubbled profusely). The layers were separated and the aqueous layer was extracted with EtOAc (3x 150 mL). The combined organic layers were washed with brine (1x 200 mL), dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated to yield a colorless oil. The oil was dried under vacuum overnight and provided **3.37** as a white solid (18.8 g, 98% yield). mp = 47-50 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.28 (br-s, 1H), 1.44 (s, 9H) ppm. <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  158.9, 82.0, 28.1 ppm.



3-oxa-2-aza-bicyclo[2.2.1]hept-5-ene-2-carboxylate (±)-*tert*-Butyl (3.38). Freshly cracked cyclopentadiene (8.20 mL, 99.0 mmol) was added to a solution of 3.37 (3.24 g, 24.3 mmol) in 4:1 MeOH/H<sub>2</sub>O (200 mL) at 4 °C (internal temperature). A solution of sodium periodate (5.71 g, 26.7 mmol) in  $H_2O$  (60 mL) was added dropwise to the reaction mixture over 35 min. The reaction mixture turned slightly yellow after 2 min, and a considerable amount of white solid was observed. The reaction was stirred for an additional 1.5 h at 4 °C, then filtered to remove the solid material and the volume was reduced with a minimal amount of heating to about 50 mL. The mixture was diluted with H<sub>2</sub>O (30 mL), and brine (50 mL), and extracted with EtOAc (5 x 85 mL). The combined organic layers were washed with brine (3 x 85 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to a brown oil. The oil was chromatographed through 150 g of silica using 4:1 hexanes/EtOAc to yield a yellow oil. Recrystallization from hexanes yielded 3.38 as a white solid (2.76 g, 58%). mp = 43-45 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  6.36 (m, 2H), 5.16 (m, 1H), 4.93 (m, 1H), 1.93 (dt, J = 8.5, 2.0 Hz, 1H), 1.68 (d, J = 8.5 Hz, 1H), 1.41 (s, 9H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 158.5, 134.0, 132.9, 83.5, 82.0, 64.9, 48.0, 28.1 ppm.



(±)-*tert*-Butyl 3-oxa-2-aza-bicyclo[2.2.2]oct-5-ene-2-carboxylate (3.39). Compound 3.39 was prepared following the same procedure used for cycloadduct 3.38. **3.37** (3.23 g, 24.3 mmol), 1,3-cyclohexadiene (4.0 mL, 41.9 mmol), and sodium periodate (5.63 g, 26.3 mmol) provided a crude orange oil. Chromatography through 200 g of silica using 4:1 hexanes/EtOAc yielded **3.39** as an orange oil that solidified to an orange waxy solid upon storage (3.53 g, 69%). Rf = 0.40 (1:1 hexanes/EtOAc – UV lamp, CAM stain). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  6.56 – 6.50 (m, 2H), 4.72 (m, 2H), 2.20 – 2.14 (m, 1H), 2.11 – 2.07 (m, 1H), 1.50 – 1.44 (m, 1H), 1.45 (s, 9H), 1.36 – 1.31 (m, 1H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  157.7, 131.7, 131.5, 81.6, 70.7, 50.1, 28.2, 23.6, 20.5 ppm.



(±)-*tert*-Butyl 8-oxa-7-aza-bicyclo[4.2.2]dec-9-ene-7-carboxylate (1c). *tert*-Butyl hydroxycarbamate (12.1 g, 90.6 mmol) was dissolved in 470 mL of MeOH in a 1-L 3-necked round-bottomed flask equipped with a mechanical stirrer and an addition funnel. The solution was cooled in a crushed ice/H<sub>2</sub>O bath to 3 °C (internal temperature). *cis,cis*-1,3-cyclooctadiene (15.0 mL, 120 mmol) was suspended in the reaction while stirring vigorously and a solution of sodium periodate (20.6 g, 95.3 mmol) in 230 mL of H<sub>2</sub>O was added to the reaction dropwise through the addition funned. After a few minutes, the reaction turned yellow and a lot of white solid formed. After 1.5 h, the addition of the NaIO<sub>4</sub> solution was complete and the reaction was stirred at 25 °C for an additional 5.5 h. The solid material was removed by filtration and washed with EtOAc (150 mL) until all of the yellow color was removed from the solid. The volume of the orange filtrate was reduced by rotary evaporation to about 300 mL (35-40 °C, 21 mm Hg). 250 mL of brine and 200 mL of Et<sub>2</sub>O were added and the layers were separated. The aqueous layer was extracted with Et<sub>2</sub>O (3 x 200 mL) and the combined Et<sub>2</sub>O layers were washed with brine (2 x 200 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated (35-40 °C, 21 mm Hg) to yield an orange solid. The crude material was loaded onto silica and purified in two portions through a Biotage 40M column using a solvent gradient from 90% CH<sub>2</sub>Cl<sub>2</sub>/hexanes to 100% CH<sub>2</sub>Cl<sub>2</sub> to afford **1c** as a light yellow solid (10.2 g, 47% yield). mp = 88-89 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  6.37 (dd, *J* = 9.7, 6.9 Hz, 1H), 5.76 (dd, *J* = 10.1, 4.4 Hz, 1H), 4.90 (br-m, 1H), 4.56 (br-m, 1H), 2.15-1.96 (m, 2H), 1.80-1.53 (m, 6H), 1.46 (s, 9H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  157.9, 131.9, 126.2, 81.1, 75.2, 54.0, 34.1, 31.4, 28.0, 25.7, 22.1 ppm. HRMS (FAB) *m*/*z* [M+H]<sup>+</sup> calcd for C<sub>13</sub>H<sub>22</sub>NO<sub>2</sub><sup>+</sup>, 240.1600; obsd, 240.1606.



(±)-Benzyl 3-oxa-2-aza-bicyclo[2.2.1]hept-5-ene-2-carboxylate (3.42). Compound 3.41 (7.34 g, 43.9 mmol) was dissolved in 460 mL of MeOH in a 3-necked 2-L flask and cooled to 2 °C (internal temperature) in an ice/H<sub>2</sub>O bath. Freshly cracked cyclopentadiene (18.0 mL, 217 mmol) was added to the reaction and sodium periodate (9.92 g, 46.4 mmol) was added dropwise over 75 min. White solid formed in the reaction and the mixture turned yellow in color after a few min. The reaction was stirred for an additional 2 h at 2 °C, and workup of the reaction was completed following the workup procedure for cycloadducts **3.38-3.40**. Chromatography of the crude amber oil through silica using a solvent gradient from 85% hexanes/EtOAc to 70% hexanes/EtOAc yielded **3.42** as a yellow oil (6.31 g, 62%). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 7.33 – 7.29 (m, 5H), 6.35 (m, 2H), 5.21 (m, 1H), 5.17 (d, *J* = 12.5 Hz, 1H), 5.10 (d, *J* = 12.5 Hz, 1H), 5.02 (m, 1H), 1.97 (dt, *J* = 8.5, 2.0 Hz, 1H), 1.72 (d, *J* = 8.5 Hz, 1H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 159.1, 135.4, 134.2, 132.8, 128.3, 128.1, 127.9, 67.6, 64.9, 48.0 ppm.



## (±)-Benzyl 8-oxa-7-aza-bicyclo[4.2.2]dec-9-ene-7-carboxylate (3.43). 3.41

(5.00 g, 29.9 mmol) was dissolved in 200 mL of MeOH in a 1-L flask and cooled to 4 °C in an ice/H<sub>2</sub>O bath. *cis,cis*-1,3-cyclooctadiene (9.5 mL, 76 mmol) was added, followed by a solution of sodium periodate (6.47 g, 30.3 mmol) in H<sub>2</sub>O (0.43 M solution) dropwise over 20 min. A lot of white solid was observed in the reaction. The mixture was stirred vigorously at 4 °C for 1.5 h, then allowed to reach RT slowly. After 6 h, workup of the reaction was completed following the workup procedure for cycloadducts **3.38-3.40**. The crude amber oil was purified through silica using a solvent gradient from 95% CH<sub>2</sub>Cl<sub>2</sub> to 100% CH<sub>2</sub>Cl<sub>2</sub> and yielded **3.43** as an amber oil (3.35 g, 41%). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.35 (m, 5H), 6.36 (dd, *J* = 10.2, 6.9 Hz, 1H), 5.79 (dd, *J* = 10.2, 4.5 Hz, 1H), 5.18 (m, 2H), 4.93 (br-m, 1H), 4.68 (br-m, 1H), 2.20 – 1.60 (m, 8H) ppm. <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  157.9, 136.1, 131.7, 128.4, 128.0, 127.8, 126.4, 76.0, 67.6, 34.2, 31.4, 25.4, 22.2 ppm. HRMS (FAB) *m*/*z* [M+H]<sup>+</sup> calcd for C<sub>16</sub>H<sub>20</sub>NO<sub>3</sub><sup>+</sup>, 274.1443; obsd, 274.1440.

General procedure for acylnitroso HDA reaction using Bu<sub>4</sub>NIO<sub>4</sub> as oxidant. Preparation of cycloadduct 3.40 (Table 3.4, entry 4). Tetrabutylammonium periodate (4.31 g, 9.92 mmol) was dissolved in 30 mL of CH<sub>2</sub>Cl<sub>2</sub> in a flame-dried 200-mL flask under Ar. The solution was cooled in an ice/H<sub>2</sub>O bath and *cis,cis*-1,3-cyclooctadiene (1.50 mL, 12.0 mmol) was added. A solution of **3.37** (1.32 g, 9.89 mmol) in 20 mL of  $CH_2Cl_2$  was added dropwise to the reaction using an addition funnel over a period of ~20 min. The reaction turned yellow, and a white solid was observed after the addition was complete. The reaction was stirred under Ar, allowing the ice/ $H_2O$  bath to melt and the reaction to reach RT gradually. During this time, the reaction became orange. The mixture was stirred overnight under Ar (24 h). Most of the solid had dissolved, and the reaction was reddish-orange in color. The reaction was washed with  $H_2O$  (2 x 50 mL) and brine (50 mL), dried over  $MgSO_4$ , filtered, and concentrated. The crude material was filtered through a plug of silica gel (~30 g), eluting with  $Et_2O$  (300 mL). The yellow filtrate was concentrated by rotary evaporation to yield a light yellow solid (1.54 g, 65.1% yield). 1H-NMR shows residual diene. The solid was dissolved in ~50 mL of toluene and concentrated by rotary evaporation with heating to get remove the excess diene, providing **3.40** as a light yellow solid (1.29 g, 54% yield).

General procedure for acylnitroso HDA reaction using CuCl/tBuOOH as oxidant. Preparation of cycloadduct 3.40 (Table 3.4, entry 6). Compound 3.37 (1.00 g, 7.52 mmol) was dissolved in 15 mL of  $CH_2Cl_2$  in a 50-mL round-bottomed flask under Ar and cooled in a -20 °C dry ice/acetone bath. CuCl (111.3 mg, 1.124 mmol) was added, followed by 5.0 M *tert*-butylhydroperoxide in decane (1.6 mL, 8.0 mmol). After 5 min, the reaction was removed from the dry ice/acetone bath and stirred in an ice/H<sub>2</sub>O bath. The reaction was stirred in the ice/H<sub>2</sub>O bath as it slowly melted. The reaction color changed from colorless to yellow, then green, then dark green, then finally blue over the course of the reaction. The reaction was quenched by adding 10 wt% sodium thiosulfate (20 mL). The layers were separated and the CH<sub>2</sub>Cl<sub>2</sub> layer was washed with 10 wt% sodium thiosulfate (20 mL), H<sub>2</sub>O (2 x 20 mL), brine (2 x 20 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated by rotary evaporation (35°C, 21 torr) to yield a blue liquid. The liquid was stored at 4 °C overnight to afford a light blue solid. Chromatography through 150 g of silica using 100% CH<sub>2</sub>Cl<sub>2</sub> provided **3.40** as a light blue solid, contaminated with Cu-salts (0.587 g, 32.6% yield).

General procedure for performing acylnitroso HDA reaction using FeCl<sub>3</sub>/H<sub>2</sub>O<sub>2</sub> as oxidant. Preparation of cycloadduct 3.43 (Table 3.4, entry 7). Compound 3.41 (5.07 g, 30.3 mmol) was dissolved in 50 mL of CH<sub>2</sub>Cl<sub>2</sub> in a 250-mL round-bottomed flask. *Cis,cis*-1,3-cyclooctadiene (4.0 mL, 32.1 mmol) was added, followed by FeCl<sub>3</sub>•6H<sub>2</sub>O (249.8 mg, 0.924 mmol, 3.0 mol%) and 1,2-ethylenediamine (0.31 mL, 4.63 mmol, 15.2 mol%). The yellow solution turned deep red in color upon the addition of FeCl<sub>3</sub> and 1,2-ethylenediamine. To this blood-red solution was added 30 wt% aqueous H<sub>2</sub>O<sub>2</sub> (22.0 mL, 215 mmol) dropwise over 50 min. Bubbling was observed as the internal temperature of the reaction climbed to 35-40 °C. The reaction was stirred for an additional 1 h at RT. Analysis of the bottom (CH<sub>2</sub>Cl<sub>2</sub>) layer by TLC (1:1 hexanes/EtOAc - CAM stain, UV lamp) indicated a complex mixture. The reaction was allowed to stir for an additional 4 h. CH<sub>2</sub>Cl<sub>2</sub> (50 mL) and H<sub>2</sub>O (100 mL) were added and

the layers were separated. The aqueous layer was extracted with  $CH_2Cl_2$  (3 x 50 mL), and the combined organic layers were washed with  $H_2O$  (50 mL), and brine (50 mL). No product was isolated – complex mixture. NOTE: all layers were quenched by adding 10 wt% sodium thiosulfate until negative to KI-starch test paper.

General procedure for preparing an ethereal solution of diazomethane (CH<sub>2</sub>N<sub>2</sub>) safely from Diazald. [IMPORTANT NOTE: Only the proper glassware should be used for the preparation of diazomethane! All glassware should *not* have ground-glass joints and should be *free* of all cracks and scratched surfaces! A blast shield should always be in place when preparing or using diazomethane until all sources of diazomethane have been quenched!] A solution of Diazald (0.867 g, 4.05 mmol) in 10 mL of Et<sub>2</sub>O (0.4 M) was added dropwise to a solution of KOH (1.46 g, 26.0 mmol) in 9 mL of 5:4 EtOH/H<sub>2</sub>O (2.9 M) that was heated in an oil bath (bath temp 68-70 °C) using the diazomethane glassware kit. It was important that the oil bath temperature was maintained within this range before, during, and after the addition of the Diazald solution. The resultant ethereal diazomethane solution was immediately condensed with a dry ice/acetone-filled condenser into a flask containing the compound to be reacted with diazomethane, or is condensed into a flask for distribution to alternative reaction vessel(s). The yellow diazomethane solution is quenched by the addition of 10 vol% HOAc in Et<sub>2</sub>O until bubbling ceases and the yellow color has completely disappeared.



(±)-Isoxazolidine-2,3,5-tricarboxylic acid 2-tert-butyl ester 3,5-dimethyl ester (3.44). NaIO<sub>4</sub> (6.79 g, 31.7 mmol) was added to a biphasic mixture of 3.38 (1.52 g, 7.70) mmol) in 45 mL of CCl<sub>4</sub>/CH<sub>3</sub>CN/H<sub>2</sub>O (1:1:1). The mixture was cooled to 0 °C and RuCl<sub>3</sub>•H<sub>2</sub>O (35.7 mg, 0.172 mmol) was added. The solution was stirred at 0 °C for 3.5 h. The solid was removed by filtration. The solid was washed with several portions of Et<sub>2</sub>O into the filtrate. The filtrate was separated and the aqueous was extracted with Et<sub>2</sub>O (3 x 50 mL). The combined organic layers were dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a light brown oil. The oil was dissolved in 25 mL Et<sub>2</sub>O and treated with excess CH<sub>2</sub>N<sub>2</sub> (see next procedure below!) at 0 °C. The reaction was quenched with 5% HOAc in H<sub>2</sub>O. The aqueous layer was made basic (pH  $\sim$ 8) by adding solid NaHCO<sub>3</sub> and saturated NaHCO<sub>3</sub>. The aqueous was extracted with  $_{E12O}$  (3 x 100 mL) and the combined Et<sub>2</sub>O layers were dried over MgSO<sub>4</sub>, filtered, and concentrated to yield 3.44 as a white solid (1.40 g, 63% from **3.38**). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 1.49 (s, 9H), 2.77 - 2.83 (m, 2H), 3.77 (s, 3H), 3.78 (s, 3H), 4.59 (t, *J* = 7.2 Hz, 1H), 4.81 (dd, *J* = 8.1, 5.2 Hz, 1H) ppm.



(±)-(3R,6S)-2-(*tert*-Butoxycarbonyl)morpholine-3,6-dicarboxylic acid (3.46). Na<sub>2</sub>CO<sub>3</sub> (0.761 g, 7.18 mmol) was added to a solution of **3.39** (1.51 g, 7.13 mmol) in 44:1 acetone/*tert*-BuOH (60 mL). The solution was cooled to -10 °C (ice/NaCl/H<sub>2</sub>O bath) and KMnO<sub>4</sub> (3.16 g, 20.0 mmol) was added slowly to maintain an internal reaction temperature less than 4 °C. The reaction was stirred at -10 °C for 1 h, and an additional 19 h at 23 °C. The reaction was diluted with 75 mL of EtOAc and quenched with 75 mL of 10% Na<sub>2</sub>S<sub>2</sub>O<sub>5</sub>. The reaction was acidified to an apparent pH of 2 (1M HCl), and the aqueous layer was saturated with NaCl and extracted with EtOAc (2 x 75 mL). The combined EtOAc layers were dried over MgSO<sub>4</sub>, filtered, and concentrated. Crude diacid **3.46** was obtained as an off-white solid (1.65 g, 84.5%) and used directly without further purification.



(±)-(3*R*,6*S*)-2-*tert*-Butyl 3,6-dimethyl morpholine-2,3,6-tricarboxylate (3.45). Crude diacid 3.46 (0.503 g, 1.83 mmol) was stirred as a suspension in 20 mL of Et<sub>2</sub>O and treated with excess  $CH_2N_2$  at 0 °C. The reaction was quenched with 5% HOAc in Et<sub>2</sub>O. The aqueous layer was made basic (pH = 8) with NaHCO<sub>3</sub> and extracted with Et<sub>2</sub>O (3 x 15 mL). The combined Et<sub>2</sub>O layers were dried over MgSO<sub>4</sub>, filtered, and concentrated to

yield a light yellow oil. The crude material was chromatographed through 50 g of silica using a solvent gradient from 80% hexanes/EtOAc to 65% hexanes/EtOAc and provided **3.45** as a colorless oil (0.349 g, 63% yield, 53% from cycloadduct **3.39**). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  4.78 (d, *J* = 2.7 Hz, 1H), 4.39 (dd, *J* = 11.7, 2.1 Hz, 1H), 3.76 (s, 3H), 3.74 (s, 3H), 2.41 (m, 1H), 2.02 – 1.70 (m, 4H), 1.49 (s, 9H) ppm. <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  169.87, 168.93, 82.54, 78.86, 52.66, 52.32, 28.09, 24.29, 23.52, 17.43 ppm. HRMS (FAB) *m*/*z* [M+H]<sup>+</sup> calcd for C<sub>13</sub>H<sub>22</sub>NO<sub>7</sub><sup>+</sup>, 304.1396; obsd, 304.1405.



(±)-(3*R*,6*S*)-3,6-Dibenzyl 2-*tert*-butyl morpholine-2,3,6-tricarboxylate (3.48). Crude diacid 3.46 (0.520 g, 1.89 mmol) and 3.47 (2.95 g, 9.38 mmol) were dissolved in 20 mL of toluene and stirred for 4.5 h at 90–95 °C. The reaction was filtered, diluted with CH<sub>2</sub>Cl<sub>2</sub> (20 mL) and concentrated to yield a yellow oil (2.98 g). The oil was chromatographed through silica using 4:1 hexanes/EtOAc and provided 3.48 as a light yellow oil (0.704 g, 82% yield, 69% from cycloadduct 3.39). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.35 – 7.29 (m, 10H), 5.28 – 5.15 (m, 4H), 4.85 (m, 1H), 4.45 (dd, *J* = 12.0, 2,4 Hz, 1H), 2.47 – 2.42 (m, 1H), 2.00 – 1.70 (m, 3H), 1.43 (s, 9H) ppm. HRMS (FAB) *m*/*z* [M+H]<sup>+</sup> calcd for C<sub>25</sub>H<sub>30</sub>NO<sub>7</sub><sup>+</sup>, 456.2022; obsd, 456.2042.



*O*-Benzyl-*N*, *N*'-dicyclohexylisourea (3.47). *N*, *N*'-dicyclohexylcarbodiimide (10.42 g, 50.50 mmol) was added to a flame-dried 100-mL round-bottomed flask under Ar and placed in a 55 °C oil bath. When the carbodiimide completely melted, benzyl alcohol (5.40 mL, 52.2 mmol) was added, followed by CuCl (151 mg, 1.52 mmol). The mixture was stirred under Ar overnight in the 55 °C oil bath. A small amount of white solid was observed in the reaction. After 16 h of stirring, the brown mixture was cooled to RT under Ar. The mixture was diluted with 80 mL of hexanes and filtered through a plug of neutral alumina (Al<sub>2</sub>O<sub>3</sub>), eluting with 20 mL of hexanes. Green solid remained on top of the alumina. The colorless filtrate was concentrated and dried under vacuum for 1 h and provided **3.47** as a colorless/light yellow oil (14.9 g, 94%). The oil was stored at -10 to 4 °C until ready to use. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 7.40 – 7.28 (m, 5H), 5.13 (s, 2H), 3.57 (br-m, 1H), 3.49 (br-m, 1H), 2.84 (br-m, 1H), 1.94 (m, 2H), 1.77 – 1.69 (m, 6H), 1.60 (m, 2H), 1.32 – 1.07 (m, 10H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 151.1, 138.1, 128.1, 127.4, 127.2, 66.44, 66.41, 54.8, 50.3, 34.4, 25.9, 25.6, 25.2, 24.9 ppm.



(±)-(3R,8S)-2-(*tert*-Butoxycarbonyl)-1,2-oxazocane-3,8-dicarboxylic acid (3.50). Prepared using RuO<sub>4</sub>. Cycloadduct 3.40 (2.06 g, 8.60 mmol) was dissolved in 3:2:2 H<sub>2</sub>O/CH<sub>3</sub>CN/CCl<sub>4</sub> (84 mL) in a 250-mL flask and cooled to 4 °C in an ice/H<sub>2</sub>O
bath. Sodium periodate (7.36 g, 34.4 mmol) was added, followed by ruthenium(III) chloride hydrate (45.7 mg, 0.220 mmol). The mixture immediately turned brown and was stirred vigorously at 4 °C for 30 min, then warmed to RT and stirred for an additional 1.5 h. The white solid was removed by filtration through a pad of celite. The celite was washed with EtOAc (80 mL) and the filtrate was diluted with brine (50 mL). The aqueous layer was acidified to a pH of 2-3 (pH paper) using 1M HCl and saturated with NaCl. The layers were separated and the aqueous layer was extracted with EtOAc (3 x 75 mL) and the combined organic layers were washed with brine (2 x 100 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield crude **3.50** as a brown/purple oil (2.58 g, 99% crude yield). The crude material was used directly without purification.

**Preparation of diacid 3.50 using KMnO<sub>4</sub>.** Cycloadduct **3.40** (3.07 g, 12.8 mmol) was dissolved in 123 mL of 40:1 acetone/*t*BuOH in a 250-mL flask and cooled to -8 °C (internal temp.) in an ice/NaCl/H<sub>2</sub>O bath. Na<sub>2</sub>CO<sub>3</sub> (1.43 g, 13.5 mmol) was added and the suspension was stirred for 2 min. The suspension turned from light yellow to orange, and KMnO<sub>4</sub> (5.67 g, 35.9 mmol) was added to the reaction slowly over a period of 2 min while stirring vigorously. The resultant purple mixture was stirred in the ice/NaCl/H<sub>2</sub>O bath for 30 min, during which time the internal temperature climbed to -2 °C. The purple/brown slurry was then warmed to RT and stirred vigorously overnight (20 h). The reaction was quenched by adding 150 mL of 10 wt% aqueous Na<sub>2</sub>S<sub>2</sub>O<sub>5</sub> (exothermic) and was stirred for 15 min at RT. EtOAc (100 mL) was added to the reaction and the aqueous layer was acidified by adding 6 M HCl slowly until the brown aqueous mixture became colorless (pH of about 2-3 - pH paper). The layers were

separated and the aqueous layer was extracted with EtOAc (3 x 100 mL). The combined organic layers were washed with brine (2 x 75 mL), dried over  $Na_2SO_4$ , filtered, and concentrated to yield crude diacid **3.50** as a yellow foam (3.51 g, 90% crude yield). The material was used directly without purification.



(±)-(3R,8S)-2-tert-Butyl 3,8-dimethyl 1,2-oxazocane-2,3,8-tricarboxylate (3.52). Crude diacid 3.50 (2.58 g, 8.49 mmol), prepared using RuO<sub>4</sub> oxidation, was dissolved in 85 mL of ether and cooled in an ice/H<sub>2</sub>O bath. The solution was treated with excess diazomethane (prepared as an ethereal solution from Diazald). The reaction was stirred for 15 min, then quenched by adding 10 vol% HOAc in ether until the yellow tint disappeared and bubbling ceased. The mixture was poured into saturated NaHCO<sub>3</sub> (150 mL) and the layers were separated. The aqueous layer (pH 7-8) was extracted with ether (2 x 75 mL). The combined organic layers were washed with brine (2 x 100 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a brown oil. The oil was chromatographed through 200 g of silica using a solvent gradient from 98% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to 95% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to yield 3.52 as a thick yellow oil (2.02 g, 71% yield from cycloadduct 3.40).

Preparation of compound 3.52 from cycloadduct 3.40 using ozonolysis:  $O_3$ , KOH, CH<sub>2</sub>Cl<sub>2</sub>/MeOH.<sup>3</sup> Cycloadduct 3.40 (1.20 g, 5.03 mmol) was dissolved in 40 mL of CH<sub>2</sub>Cl<sub>2</sub> and cooled to -78 °C in a dry ice/acetone bath. 10 mL of 2.5M methanolic

KOH (25 mmol) was added, and the colorless solution immediately turned light yellow. The reaction was stirred at -78 °C as ozone was bubbled through the reaction. The reaction immediately turned bright orange and a precipitate formed. After 85 min, the flow of ozone was stopped, the reaction was removed from the dry ice/acetone bath, and the reaction was diluted with  $Et_2O$  (50 mL), then  $H_2O$  (50 mL) until the orange reaction became completely colorless and the solid dissolved. The reaction was stirred for a few min to reach RT, then the layers were separated and the aqueous layer was extracted with Et<sub>2</sub>O (3 x 75 mL). The combined organic layers were washed with brine (2 x 50 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a light yellow oil (1.36 g, 82% crude yield). The crude material was loaded onto silica and chromatographed through a Biotage 40S column (40 g of silica) using 97% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc and provided 3.52 as a light yellow oil (466 mg, 28% yield). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  4.85 – 4.68 (br-m, 1H), 4.44 – 4.28 (br-m, 1H), 3.70 (s, 3H), 3.69 (s, 3H), 2.14 – 2.02 (m, 2H), 1.95 – 1.85 (m, 2H), 1.72 - 1.67 (m, 1H), 1.47 - 1.32 (m, 10H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$ 170.2, 82.3, 81.1, 80.4, 63.5, 61.9, 52.2, 51.9, 29.0, 28.1, 26.8, 25.8, 24.1, 24.03, 23.99, 23.96 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>15</sub>H<sub>26</sub>NO<sub>7</sub><sup>+</sup>, 332.1709; obsd, 332.1685.



(±)-(3R,8S)-Dimethyl 1,2-oxazocane-3,8-dicarboxylate (3.53). MeOH (100 mL) was added to a 250-mL round-bottomed flask under Ar, then cooled in an ice/H<sub>2</sub>O bath. Thionyl chloride (4.00 mL, 55.0 mmol) was added slowly and the solution was stirred for 5 min. A solution of crude 3.50 (4.14 g, 13.6 mmol) in 40 mL of MeOH was

transferred to the flask via cannula, and the reaction was stirred in the ice/H<sub>2</sub>O bath under Ar. After 25 min, the reaction was removed from the ice/H<sub>2</sub>O bath and allowed to stir at After 22 h, the reaction was concentrated to yield a brown oil. The oil was RT. partitioned between 75 mL of CH<sub>2</sub>Cl<sub>2</sub> and 200 mL of saturated aqueous NaHCO<sub>3</sub> (bubbled profusely). The layers were separated and the aqueous layer (pH 8 - pH paper) was extracted with  $CH_2Cl_2$  (3 x 75 mL). The combined  $CH_2Cl_2$  layers were washed with brine (1 x 80 mL), dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated to yield a brown oil (2.27 g, 72% crude yield). The oil was loaded onto silica and chromatographed through a Biotage 40S column (40 g of silica) using a solvent gradient of 100% CH<sub>2</sub>Cl<sub>2</sub> to 95% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc and provided **3.53** as a yellow oil (1.93 g, 61% from cycloadduct **3.40**). Rf = 0.26 (1:1 hexanes/EtOAc – CAM stain). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  6.27 (br-s, 1H), 4.22 (dd, *J* = 10.0, 3.5 Hz, 1H), 3.72 (s, 3H), 3.71 (s, 3H), 2.42 – 2.22 (m, 1H), 2.08 - 1.92 (m, 3H), 1.73 - 1.68 (m, 3H), 1.57 - 1.52 (m, 1H) ppm. <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 172.4, 171.4, 80.7, 62.0, 51.8, 51.6, 28.0, 26.9, 26.0, 24.1 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>10</sub>H<sub>18</sub>NO<sub>5</sub><sup>+</sup>, 232.1185; obsd, 232.1198.

Attempted formation of compounds 3.56 and 3.57 by ozonolysis of cycloadduct 3.40. Cycloadduct 3.40 (501 mg, 1.83 mmol) was dissolved in 18 mL of  $5:1 \text{ CH}_2\text{Cl}_2/\text{MeOH}$  in a single-necked 50-mL round-bottomed flask. The light yellow solution was cooled to -78 °C in a dry ice/acetone bath and NaHCO<sub>3</sub> (602 mg, 7.17 mmol) was added. Ozone was bubbled into the solution until the color changed to deep blue (5 min). The solution was purged with Ar until the blue color disappeared. The mixture was allowed to warm to RT and the solid material was removed by filtration.

The solid was rinsed with portions of  $CH_2Cl_2$  (10 mL), and the filtrate was concentrated to yield a white foam. The foam was dissolved in anhydrous 1:1  $CH_2Cl_2$ /pyridine (20 mL) in a single-necked 100-mL round-bottomed flask under Ar. Distilled acetic anhydride (0.70 mL, 7.5 mmol) was added to the reaction and the colorless solution was stirred under Ar at RT. The solution gradually turned from colorless to yellow, then yellow/orange. After 2 days, the reaction was concentrated by passing a stream of air over the solution for ~2 h. The orange residue was partitioned between 1M HCl (50 mL) and EtOAc (30 mL). The layers were separated and the aqueous layer was extracted with EtOAc (3 x 20 mL). The combined organic layers were washed with 1M HCl (2 x 20 mL), H<sub>2</sub>O (20 mL), saturated NaHCO<sub>3</sub> (2 x 20 mL), and brine (2 x 20 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a reddish-orange, sweet-smelling oil (~0.432 g crude yield). Analysis of the crude material indicated a complex mixture of products.



(±)-(2S,6aR)-2-(Hydroxymethyl)-hexahydrooxazolo[3,4-b][1,2]oxazocin-

**9(2H)-one (3.59).** Cycloadduct **3.43** (1.18 g, 4.32 mmol) was dissolved in 42 mL of 5:1 CH<sub>2</sub>Cl<sub>2</sub>/MeOH in a single-necked 100-mL round-bottomed flask and cooled to -78 °C in a dry ice/acetone. Ozone was bubbled through the light yellow solution until the blue color of ozone was observed in the solution (~5 min). The solution was purged with Ar until the blue color disappeared then the reaction was removed from the dry ice/acetone bath and transferred to a crushed ice/NaCl/H<sub>2</sub>O bath. NaBH<sub>4</sub> (1.02 g, 27.0 mmol) was

added to the reaction in portions over 60 min. Bubbling was observed during this time. After the addition was complete, the cloudy mixture was allowed to slowly reach RT over 45 min. The volume of the mixture was reduced by rotary evaporation (40-45 °C, 21 mm Hg) to 5 mL, at which time a white precipitate was observed. The mixture was partitioned between H<sub>2</sub>O (50 mL) and EtOAc (50 mL) and the layers were separated. The aqueous layer was extracted with EtOAc (2 x 30 mL) and the combined organic layers were washed with brine (2 x 30 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a colorless oil. The oil was chromatographed through 150 g of silica gel and yielded **3.59** as a tan-colored solid (423.3 mg, 48% yield). mp = 64-65 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  4.38 (dd, *J* = 8.4, 7.5 Hz, 1H), 4.27 (dd, *J* = 11.1, 3.9 Hz, 1H), 3.98 – 3.88 (m, 1H), 3.88 (dd, *J* = 11.4, 8.4 Hz, 1H), 3.76 – 3.65 (m, 1H), 3.51 – 3.32 (m, 1H), 1.92 – 1.31 (m, 8H) ppm. <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  162.1, 85.7, 67.1, 63.4, 60.9, 27.1, 24.8, 22.7, 22.0 ppm. MS (FAB) m/z [M+H]<sup>+</sup> at 202.



**2,2,2-Trifluoroethyl** *N*-hydroxycarbamate (3.63). *N*,*N*'-Carbonyldiimidazole (2.468 g, 15.22 mmol) was dissolved in 50 mL of anhydrous THF in a flame-dried single-necked 250-mL round-bottomed flask under Ar. To this solution was added 2,2,2-trifluoroethanol (1.00 mL, 13.9 mmol). The resultant solution was stirred under Ar at RT for 2 h. In a separate flame-dried single-necked 100-mL round-bottomed flask, hydroxylamine hydrochloride (1.111 g, 15.99 mmol) was dissolved in about 40 mL of anhydrous pyridine and transferred via cannula to the flask containing the acylimidazole. The light yellow solution immediately became cloudy and warmed slightly. After

stirring for about 2 h, the reaction was analyzed by TLC and a FeCl<sub>3</sub>-positive spot was observed. The reaction mixture was concentrated and the yellow oil/solid mixture was partitioned between EtOAc (50 mL) and H<sub>2</sub>O (50 mL). 6M HCl (10 mL) was added and the layers were separated. The organic layer was washed with portions of 3M HCl (2 x 25 mL) and brine (20 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, concentrated and provided **3.63** as a yellow oil (1.29 g, 58% yield). Rf = 0.30 (1:1 hexanes/EtOAc – FeCl<sub>3</sub> stain). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.72 (s, 1H), 4.52 (qd, *J* = 8.0, 0.5 Hz, 2H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  157.1, 122.6 (q, *J* = 276 Hz), 61.5 (q, *J* = 37 Hz) ppm. MS (FAB) *m*/*z* [M+H]<sup>+</sup> at 160.



(±)-2,2,2-Trifluoroethyl 8-oxa-7-aza-bicyclo[4.2.2]dec-9-ene-7-carboxylate (3.64). Compound 3.63 (1.289 g, 8.104 mmol) was dissolved in 40 mL of methanol in a 250-mL round-bottomed flask. *cis,cis*-1,3-cyclooctadiene (1.30 mL, 10.4 mmol) was added. The reaction was stirred vigorously as a solution of sodium periodate (1.925 g, 8.910 mmol) in 20 mL of H<sub>2</sub>O was added dropwise over about 30 min. A yellow color was observed immediately, and a white solid was observed after a few minutes into the addition. After the addition was complete, the mixture was stirred vigorously at RT. After 2 h, no 3.63 remained by TLC. The white solid was removed by filtration and washed with EtOAc (10 mL). The volume of the orange filtrate was reduced to half by rotary evaporation and partitioned between EtOAc (50 mL) and brine (50 mL). The layers were separated and the aqueous was extracted with EtOAc (4 x 25 mL).

combined organic layers were washed with brine (25 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to yield a brown oil (1.93 g). The oil was chromatographed through 200 g of silica using a solvent gradient from 90% CH<sub>2</sub>Cl<sub>2</sub>/hexanes to 100% CH<sub>2</sub>Cl<sub>2</sub> to yield **3.64** as a yellow oil (1.103 g, 51.3% yield). The NMR spectra for compound **3.64** indicated a significant amount of impurities; however, the MS spectrum confirmed the presence of a 1:1 adduct. MS (FAB) m/z [M+H]<sup>+</sup> at 266.



## (±)-(2S,6aR)-9-Oxo-octahydrooxazolo[3,4-b][1,2]oxazocine-2-carbaldehyde

(3.66). Oxalyl chloride (0.12 mL, 1.4 mmol) was added to 5 mL of anhydrous  $CH_2Cl_2$  in a flame-dried, single-necked 50-mL round-bottomed flask cooled to -78 °C (dry ice/acetone) under Ar. In a separate flame-dried single-necked 10-mL round-bottomed flask was added anhydrous dimethyl sulfoxide (0.17 mL, 2.4 mmol) to 5 mL of anhydrous  $CH_2Cl_2$ . The DMSO solution was added to the oxalyl chloride solution dropwise over about 1 min and the resultant solution was stirred at -78 °C for 10 min under Ar. In a separate flame-dried single-necked 25-mL round-bottomed flask under Ar was dissolved **3.59** (0.151 g, 0.750 mmol) in 5 mL of anhydrous  $CH_2Cl_2$ . This solution was transferred slowly to the flask that contained the oxidant via cannula. The resultant solution was stirred at -78 °C for 45 min, then triethylamine (0.60 mL, 4.3 mmol) was added. The reaction was stirred at -78 °C for 20 min, then removed from the dry ice/acetone bath and allowed to reach RT while stirring under Ar. After 2 h, the reaction was diluted with  $CH_2Cl_2$  (20 mL) and  $H_2O$  (50 mL) and the layers were separated. The aqueous layer was extracted with  $CH_2Cl_2$  (2 x 20 mL) and the combined organic layers were washed with brine (20 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a yellow oil (149 mg). The crude material was analyzed by TLC and was composed of mostly one spot with the same Rf as the starting material but that stained darker to CAM stain. The crude material was chromatographed through 20 g of silica using a solvent gradient from 100%  $CH_2Cl_2$  to 90%  $CH_2Cl_2/EtOAc$  and provided **3.66** as a yellow oil (80.5 mg, 54% yield). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  10.14 (s, 1H), 4.40 (dd, *J* = 8.4, 7.2 Hz, 1H), 4.21 (ddd, *J* = 11.4, 2.7, 0.6 Hz, 1H), 3.89 (dd, *J* = 11.4, 8.4 Hz, 1H), 3.82 – 3.71 (m, 1H), 2.19 – 2.05 (m, 1H), 2.00 – 1.86 (m, 3H), 1.81 – 1.69 (m, 3H), 1.58 – 1.43 (m, 1H) ppm. <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  204.0, 166.8, 89.0, 66.5, 60.9, 27.4, 24.7, 22.5, 22.0 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>9</sub>H<sub>14</sub>NO<sub>4</sub><sup>+</sup>, 200.1923; obsd, 200.0935.



(±)-(2*R*,7*S*)-Dimethyl 2-(*tert*-butoxycarbonylamino)-7-hydroxyoctanedioate (3.69). General procedure for N-O bond reduction using samarium diiodide (Table 3.5, entry 4). To a flame-dried 50-mL round-bottomed flask under Ar was added samarium metal (160 mg, 1.06 mmol, 40 mesh) and 9 mL of anhydrous THF. The mixture was stirred vigorously and cooled in an ice/H<sub>2</sub>O bath to 4°C under Ar.

Diiodomethane (0.072 mL, 0.89 mmol) was added slowly and the mixture was stirred in the ice/H<sub>2</sub>O bath under Ar. After 15 min, the solution had turned from colorless to light yellow to a deep blue-green color (indicating formation of SmI<sub>2</sub>). After 1.5 h of stirring, **3.52** (144 mg, 0.435 mmol) was dissolved in 1 mL of anhydrous THF in a flame-dried 5mL flask under Ar and transferred to the  $SmI_2$  solution via cannula. After a few min, the blue-green color turned to a dark green color. No further color change was observed. Reaction progress was monitored by TLC (1:1 hexanes/EtOAc - CAM stain). After 5 h, no starting material was observed and a new, more polar spot (Rf ~0.21) was observed by TLC. The reaction was diluted with CH<sub>2</sub>Cl<sub>2</sub> (30 mL) and 10 wt% aqueous Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> (20 mL) was added to quench the reaction. An emulsion formed and  $H_2O$  (~50-60 mL) was added. 1M HCl (~20 mL) was added and the layers were separated. The aqueous layer was extracted with  $CH_2Cl_2$  (2 x 20 mL). The combined organic layers were washed with 1M HCl (1 x 30 mL), dried over MgSO<sub>4</sub>, filtered through a pad of celite, and concentrated by rotary evaporation (30°C, 21 mm Hg) to afford a pungent-smelling yellow/white solid/oil. The crude material was chromatographed through 50 g of silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 80% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc and yielded **3.69** as a colorless oil (57.0 mg, 39% yield). Rf = 0.21 (1:1 hexanes/EtOAc – CAM stain). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  5.04 (br-d, J = 7.5 Hz, 1H), 4.26 (m, 1H), 4.15 (dd, J = 7.5, 4.0 Hz, 1H), 3.76 (s, 3H), 3.71 (s, (3H), 2.52 (br, 1H), 1.79 - 1.73 (m, 2H), 1.64 - 1.57 (m, 2H), 1.41 (s, 9H), 1.45 – 1.33 (m, 4H) ppm.  $^{13}$ C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  175.6, 173.3, 155.3, 79.8, 70.1, 53.3, 52.5, 52.2, 34.0, 32.5, 28.2, 24.9, 24.3 ppm. MS (FAB) m/z [M+H]<sup>+</sup> at 334, 278, 234 (base peak), 174.

General procedure for N-O bond reduction using  $Mo(CO)_6$ . Attempted reduction of 3.52 (Table 3.5, entry 2). Compound 3.52 (0.101 g, 0.303 mmol) was dissolved in 15:1 CH<sub>3</sub>CN/H<sub>2</sub>O (4.4 mL) in a 25-mL flask. Molybdenum hexacarbonyl (59 mg, 0.22 mmol) was added and the solution was heated at reflux (oil bath temperature 85-100 °C) overnight. The color of the solution changed to yellow, then deep brown. The mixture was cooled to RT, then cooled at 0 °C for 1 h and filtered through a pad of celite, rinsing the solid residue with portions of EtOAc. The filtrate was concentrated to yield a pale yellow residue. Analysis of the crude material did not indicate the presence of the desired product, compound **3.69**.

General procedure for N-O bond reduction using Zn/HOac. Attempted reduction of 3.52 (Table 3.5, entry 3). Zinc dust (0.329 g, 5.03 mmol), previously activated by rinsing with conc. HCl and drying in the oven, was added to 3.52 (0.100 g, 0.303 mmol) in 2.5 mL of glacial acetic acid in a 10-mL flask. The mixture was stirred at RT overnight. Analysis of the mixture only indicated the presence of 3.52 and the mixture was heated in an oil bath maintained at 75-80 °C overnight. No change was observed during this time.

General procedure for N-O bond reduction using Na-Hg. Attempted reduction of 3.52 (Table 3.5, entry 5). Compound 3.52 (0.122 g, 0.367 mmol) was added to a flame-dried 25-mL flask and dissolved in 2:1 MeOH/THF (6 mL) under Ar. The solution was cooled to 4 °C in an ice/H<sub>2</sub>O bath and Na<sub>2</sub>HPO<sub>4</sub> (0.264 g, 1.86 mmol) was added. 5% sodium-mercury amalgam (0.868 g, 1.90 mmol sodium) was added and the resultant cloudy mixture was stirred for 2 h. Liquid mercury was observed in the bottom of the flask and the mixture was warmed to RT. The mixture was stirred at RT for 2 h and the solution was decanted from the liquid mercury and filtered through celite. Multiple portions of EtOAc and MeOH were added to the reaction flask, decanted from the residual mercury, and filtered through celite. The filtrate was concentrated to yield a white solid. Analysis of the solid material did not indicate the presence of the desired compound.

General procedure for N-O bond reduction using hydrogenolysis. Attempted reduction of 3.53 (Table 3.5, entry 7). Acetyl chloride (0.075 mL, 1.1 mmol) was added to 5 mL of anhydrous methanol under Ar in a 250-mL Parr bomb. After 30 min, 3.53 (0.204 g, 0.882 mmol) in 5 mL of methanol was added and the Parr bomb was evacuated and purged with Ar (2x). 10% Pd/C (33 mg, 16 wt%) was added and the reaction flask was evacuated and purged with Ar (2x), then evacuated and purged with H<sub>2</sub> (2x). The mixture was shaken under H<sub>2</sub> (2.8 bar, 30 psi) using a Parr hydrogenation apparatus for 6 h. The mixture was evacuated and purged with Ar (3x), then filtered through celite. The filtrate was concentrated to yield a pale yellow residue. Analysis of the residue indicated mostly 3.53 remained in the mixture in addition to a small amount of baseline ninhydrin-positive material (presumably amine 3.70) and a less polar product. Transesterification may have occurred in the reaction, but this was not confirmed.



(±)-*tert*-Butyl (1R,4S)-4-hydroxycyclopent-2-enylcarbamate (3.71). Cycloadduct 3.38 (1.11 g, 5.63 mmol) was dissolved in 20 mL of 4:1 CH<sub>3</sub>CN/H+O.  $Mo(CO)_6$  (598 mg, 2.27 mmol) was added, followed by NaBH= (660 mg, 17.4 mmol). The reaction bubbled and turned bright yellow. The reaction was heated to 70 °C with a condenser and stirred under mild reflux. The reaction turned brown after 30 min. The reaction was monitored by TLC (1:1 hexanes/EtOAc). After 12 h, the reaction was cooled to room temperature and then cooled to 0 °C for 30 min. The reaction was filtered through a pad of celite to remove the solid. The celite was washed with EtOAc and CH<sub>3</sub>CN and the yellow filtrate was concentrated to yield a dark yellow oil (1.29 g). The oil was chromatographed through 100 g of silica using 1:1 hexanes/EtOAc to yield 3.71 as a yellow oil (898 mg, 80%). Rf = 0.20 (1:1 hexanes/EtOAc – UV lamp, CAM stain). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 5.94 (m, 1H), 5.80 (m, 1H), 5.01 (m, 1H), 4.65 (m, 1H), 4.42 (m, 1H), 3.37 (br, 1H), 2.69 (m, 1H), 1.49 (d, J = 14.5 Hz, 1H), 1.41 (s, 10H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 155.3, 136.0, 134.1, 79.5, 75.0, 54.7, 41.3, 28.4 ppm.



(±)-*tert*-Butyl (1*R*,4*S*)-4-hydroxycyclohex-2-enylcarbamate (3.72). Prepared following the same procedure used for 3.71. Cycloadduct 3.39 (1.05 g, 4.97 mmol),  $Mo(CO)_6$  (525 mg, 1.99 mmol), and NaBH<sub>4</sub> (600 mg, 15.9 mmol) provided a dark yellow oil (1.12 g). The oil was chromatographed through 75 g of silica using 1:1

hexanes/EtOAc to yield **3.72** as a yellow oil (903 mg, 85%). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  5.861 (dq, J = 9.9 Hz, 1H), 5.741 (dd, J = 10.2, 2.7 Hz, 1H), 4.596 (br-m, 1H), 4.151 (br-m, 1H), 1.855 (m, 2H), 1.675 (m, 2H), 1.443 (s, 9H) ppm.

(±)-Acetic acid 4-tert-butoxycarbonylamino-cyclopent-2-enyl ester (3.73). Alcohol 3.71 (852 mg, 4.28 mmol) was dissolved in 30 mL of CH<sub>2</sub>Cl<sub>2</sub>. Imidazole (501 mg, 7.36 mmol) was added to the reaction and the reaction was cooled to 0  $^{\circ}$ C. Acetyl chloride (0.460 mL, 6.45 mmol) was added slowly to the reaction. A white precipitate formed. The reaction was stirred overnight and monitored by TLC (1:1 hexanes/EtOAc). After 18 h, 1 equiv more acetyl chloride was added to the reaction. No change was observed after 4 h. The reaction was diluted and partitioned between  $CH_2Cl_2$  and  $H_2O$ . The aqueous layer (pH 2) was extracted with  $CH_2Cl_2$ . The combined organic layers were washed with 1M HCl, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to yield an orange oil (867 mg). The oil was chromatographed through 87 g of silica using 4:1 to 3:1 hexanes/EtOAc to yield a colorless oil. Recrystalization from hexanes yielded 3.73 as white needlelike crystals (673 mg, 65%). mp = 48-51 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$ 5.95 (d, J = 5.5 Hz, 1H), 5.89 (d, J = 5.5 Hz, 1H), 5.49 (m, 1H), 4.65 (m, 1H), 2.79 (m, 1H), 2.00 (s, 3H), 1.49 (dt, J = 14.5, 4.0 Hz, 1H), 1.41 (m, 10H) ppm. <sup>13</sup>C NMR (125) MHz, CDCl<sub>3</sub>) δ 170.5, 154.9, 136.9, 132.0, 79.5, 77.4, 54.2, 38.5, 28.3, 21.1 ppm.



(±)-Acetic acid 4-tert-butoxycarbonylamino-cyclohex-2-enyl ester (3.74). Compound 3.74 was prepared following the same procedure for compound 3.73. Alcohol 3.72 (903 mg, 4.23 mmol), imidazole (489 mg, 7.18 mmol), and acetyl chloride (0.450 mL, 6.31 mmol) provided a yellow oil (900 mg). The oil was chromatographed through 90 g of silica using 4:1 to 3:1 hexanes/EtOAc to yield 3.74 as a white solid (630 mg, 59%). mp = 82-83 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  5.81 (dd, *J* = 10, 2.5 Hz, 1H), 5.76 (d, *J* = 10 Hz, 1H), 5.14 (m, 1H), 4.60 (d, *J* = 7.0 Hz, 1H), 4.11 (m, 1H), 2.00 (s, 3H), 1.87 – 1.69 (m, 3H), 1.63 – 1.54 (m, 1H), 1.40 (s, 9H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  170.4, 155.1, 133.5, 127.9, 79.4, 68.1, 66.8, 45.8, 28.3, 25.7, 21.2 ppm.



(±)-*tert*-Butyl (1*R*,4*S*)-4-(tert-butyldimethylsilyloxy)cyclopent-2-

**enylcarbamate** (3.75). Compound 3.71 (201 mg, 1.01 mmol) was dissolved in 20 mL of anhydrous DMF in a flame-dried 50-mL round-bottomed flask under Ar. Imidazole (277 mg, 4.07 mmol) was added, followed by TBSCl (303 mg, 2.01 mmol) and the solution was stirred at RT under Ar. After 4 h, TLC of the reaction (1:1 hexanes/EtOAc - CAM stain) did not show any starting material and the reaction was complete. The reaction was concentrated to 10 mL by rotary evaporation.  $H_2O$  (75 mL) and EtOAc (50 mL) were added and the layers were separated. The aqueous layer was extracted with EtOAc (3 x 50 mL), and the combined organic layers were washed with brine (2 x 35 mL), dried

over MgSO<sub>4</sub>, filtered, and concentrated to yield a yellow oil. The oil was chromatographed through 30 g of silica using 100% CH<sub>2</sub>Cl<sub>2</sub> and provided **3.75** as a colorless oil (0.288 g, 91%). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  5.85 – 5.78 (m, 2H), 4.69 (t, J = 6.0 Hz, 1H), 4.61 (br-m, 1H), 4.57 (br-m, 1H), 2.71 (m, 1H), 1.40 (s, 9H), 1.36 (m, 1H), 0.89 (s, 9H), 0.07 (s, 6H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  155.1, 136.6, 133.6, 75.4, 54.4, 42.8, 28.4, 25.9, 18.2, -4.7 ppm.

(±)-*tert*-Butyl (1R,4S,Z)-4-hydroxycyclooct-2-enylcarbamate (3.76). Cycloadduct 3.40 (1.50 g, 6.27 mmol) was dissolved in 64 mL of 15:1 CH<sub>3</sub>CN/H<sub>2</sub>O in a 250-mL round-bottomed flask and heated in a 55 °C oil bath. Molybdenum hexacarbonyl (1.19 g, 4.52 mmol) was added to the reaction in one portion. A jacketed condenser was attached to the flask and the reaction was heated to reflux (oil temp = 95°C). The reaction turned yellow after 15 min of heating, then turned brown and finally black overnight. After 19 h, the TLC analysis (1:1 hexanes/EtOAc - CAM stain), the of the reaction showed clean, but incomplete conversion to the product. Additional  $Mo(CO)_6$  (0.462 g, 1.75 mmol) was added (total  $Mo(CO)_6 = 1$  equivalent) and the reaction was heated to reflux for an additional 2 h. The reaction was cooled to RT, then filtered through a pad of celite. The brown filtrate was concentrated to yield a brown oil and a glassy solid. The mixture was partitioned between EtOAc (75 mL) and  $H_2O$  (75 mL) and the layers were separated. The aqueous layer was extracted with EtOAc (3 x 60 mL) and the combined organic layers were washed with brine (2 x 75 mL), dried over

MgSO<sub>4</sub>, filtered, and concentrated to yield a tan solid. The crude material was loaded onto silica and chromatographed through a Biotage 40M column (90 g of silica) using a solvent gradient from 75% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to 60% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to yield **3.76** as a white solid (1.44 g, 95%). mp = 137-139 °C. Rf = 0.25 (1:1 hexanes/EtOAc – CAM stain). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  5.57 (dd, *J* = 11, 7.5 Hz, 1H), 5.23 (ddd, *J* = 11.5, 10.5, 1.5 Hz, 1H), 4.63 (br, 1H), 4.34 (br, 1H), 2.44 (br, 1H), 1.91 – 1.81 (m, 2H), 1.61 – 1.53 (m, 2H), 1.48 – 1.41 (m, 3H), 1.41 (s, 9H), 1.33 – 1.25 (m, 1H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  155.2, 134.7, 130.0, 79.4, 69.3, 48.8, 38.2, 36.5, 28.4, 24.0, 23.4 ppm. HRMS (FAB) *m*/*z* [M+H]<sup>+</sup> calcd for C<sub>13</sub>H<sub>24</sub>NO<sub>3</sub><sup>+</sup>, 242.1756; obsd, 242.1754.

**Preparation of 3.76 using SmI<sub>2</sub>.** Compound **3.76** was prepared following the general procedure for N-O bond reduction using samarium diiodide. Cycloadduct **3.40** (102 mg, 0.426 mmol) was treated with 2.5 eq of samarium diiodide prepared from samarium metal (0.161 g, 1.07 mmol) and diiodomethane (0.070 mL, 0.87 mmol) and provided **3.76** as a white foam after chromatography through 10 g of silica (81 mg, 79%).



 $(\pm)$ -(1*S*,4*R*,*Z*)-4-(*tert*-Butoxycarbonylamino)cyclooct-2-enyl acetate (3.77). Alcohol 3.76 (2.59 g, 10.7 mmol) was dissolved in 50 mL of anhydrous pyridine in a flame-dried single-necked 250-mL round-bottomed flask under Ar. Acetic anhydride (2.50 mL, 26.5 mmol) was added and the light yellow solution was stirred at RT under Ar. After 25 h, the reaction was concentrated by rotary evaporation to yield a light yellow solid. The solid was partitioned between 1M HCl (100 mL) and EtOAc (100 mL). The layers were separated and the aqueous layer was extracted with EtOAc (2 x 40 mL). The combined organic layers were washed with 1M HCl (2 x 35 mL), H<sub>2</sub>O (35 mL), saturated NaHCO<sub>3</sub> (2 x 35 mL), and brine (35 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a white solid (2.99 g, 98% yield). Further purification was not necessary, but could be carried out by chromatography through silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 80% CH<sub>2</sub>Cl<sub>2</sub>. mp = 117-119 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  5.60 (m, 1H), 5.54 (dd, *J* = 10.2, 7.6 Hz, 1H), 5.33 (t, *J* = 9.3 Hz, 1H), 4.57 (br-m, 1H), 4.37 (br-m, 1H), 2.00 (s, 3H), 1.92 (m, 2H), 1.62-1.44 (m, 4H), 1.41 (s, 9H), 1.36-1.29 (m, 2H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  170.0, 154.9, 131.2, 129.7, 79.2, 72.1, 49.0, 36.2, 35.1, 28.3, 23.8, 23.0, 21.2 ppm. HRMS (FAB) *m/z* [M+H]<sup>+</sup> calcd for C<sub>15</sub>H<sub>26</sub>NO<sub>4</sub><sup>+</sup>, 284.1862; obsd, 284.1847.



( $\pm$ )-*tert*-Butyl (1*R*,4*S*,*Z*)-4-(benzyloxy)cyclooct-2-enylcarbamate (3.78). Alcohol 3.76 (0.504 g, 2.09 mmol) was dissoved in 20 mL of anhydrous THF under Ar in a flame-dried 100-mL round-bottomed flask. Benzyl bromide (0.50 mL, 4.5 mmol) was added and the light yellow solution was cooled in an ice/H<sub>2</sub>O bath under Ar. Sodium hydride (0.214 g, 50% dispersion in mineral oil, 4.46 mmol) was added to the reaction in one portion. The reaction immediately turned cloudy. The cloudy suspension was stirred in the ice/H<sub>2</sub>O bath, slowly reaching RT under Ar overnight. Monitored reaction by TLC (4:1 hexanes/EtOAc - CAM stain). Alcohol 3.76 remained after stirring overnight and over the weekend (3 days). An additional 20 mL of anhydrous THF was added to the white suspension and the reaction was stirred under Ar at RT. After 2 days (5 days total) no **3.76** was observed by TLC of the reaction mixture. The reaction was cooled in an ice/H<sub>2</sub>O bath and quenched by adding H<sub>2</sub>O dropwise until all of the white solid dissolved completely (~15 mL - some bubbling observed). Et<sub>2</sub>O (50 mL) was added and the layers were separated. The aqueous layer was extracted with  $Et_2O$  (2 x 50 mL), and the combined organic layers were washed with brine (2 x 50 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a light yellow oil. The crude material was loaded onto silica and chromatographed through a 40S Biotage column (40 g of silica) using a solvent gradient from 100% hexanes to 80% hexanes/EtOAc and provided **3.78** as a light yellow oil (553 mg, 80% yield). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 7.41 – 7.28 (m, 5H), 5.67 (ddd, J = 11, 7.0, 1.0 Hz, 1H), 5.46 (ddd, J = 11, 8.0, 1.5 Hz, 1H), 4.73 (d, J = 7.0 Hz, 1H), 4.65 (d, J = 11.5 Hz, 1H), 4.49 (d, J = 11.5 Hz, 1H), 4.39 (br, 2H), 2.07 - 2.01 (m, 1H), 1.91(m, 1H), 1.63 - 1.51 (m, 13H), 1.51 (s, 9H), 1.38 - 1.29 (m, 1H) ppm. <sup>13</sup>C NMR (125) MHz, CDCl<sub>3</sub>) δ 155.0, 138.4, 133.1, 132.0, 128.2, 127.7, 127.3, 79.1, 76.2, 70.8, 49.0, 36.7, 35.8, 28.3, 24.2, 23.3 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>20</sub>H<sub>30</sub>NO<sub>3</sub><sup>+</sup>, 332.2226; obsd, 332.2200.



(±)-*tert*-Butyl (1*R*,4*S*,*Z*)-4-(tert-butyldimethylsilyloxy)cyclooct-2enylcarbamate (3.79). Alcohol 3.76 (1.44 g, 5.96 mmol) was dissolved in 60 mL of DMF in a single-necked 250-mL round-bottomed flask under Ar. Imidazole (1.76 g, 25.9 mmol) was added, followed by TBSCl (1.81 g, 12.0 mmol). The solution was stirred at RT under Ar. After 2.5 days, the reaction was complete by TLC analysis of the reaction mixture (4:1 hexanes/EtOAc - CAM stain). The yellow solution was concentrated by rotary evaporation (55 °C, 20 mm Hg) to about 10 mL, then partitioned between  $H_2O$  (50 mL) and EtOAc (100 mL). The layers were separated and the aqueous layer was extracted with EtOAc (2 x 75 mL). Some brine was added to the biphasic mixture during separation to break the emulsion that formed. The combined EtOAc layers were washed with H<sub>2</sub>O (50 mL), and brine (2 x 50 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated by rotary evaporation (45 °C, 20 mm Hg) to yield 2.8 g of a yellow oil. The oil was chromatographed through 200 g of silica using a solvent gradient from 90%  $CH_2Cl_2$ /hexanes to 100%  $CH_2Cl_2$  and yielded **3.79** as a white solid (2.02 g, 95% yield). mp = 84-85 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  5.54 (dd, J = 11.0, 7.0 Hz, 1H), 5.17 (ddd, J = 11.0, 8.5, 1.5 Hz, 1H), 4.53 (br, 2H), 4.34 (br, 1H), 1.89 – 1.79 (m, 2H), 1.55 – 1.44 (m, 5H), 1.43 (s, 9H), 1.31 – 1.24 (m, 1H), 0.88 (s, 9H), 0.063 (s, 3H), 0.057 (s, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 155.0, 135.7, 128.5, 79.2, 70.2, 49.0, 39.3, 36.9, 28.4, 25.8, 24.2, 23.4, 18.2, -4.6, -4.8 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for  $C_{19}H_{38}NO_3Si^+$ , 356.2621; obsd, 356.2633.



(±)-Benzyl (1*R*,4*S*,*Z*)-4-hydroxycyclooct-2-enylcarbamate (3.80). Compound
3.80 was prepared following the same procedure for 3.76. Cycloadduct 3.43 (5.73 g, 21.0 mmol) and molybdenum hexacarbonyl (5.55 g, 21.0 mmol) provided a brown solid.

Chromatography through 180 g of silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 60% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc provided **3.80** as a white solid (5.01 g, 87%). mp = 125-126 °C. Rf = 0.18 (1:1 hexanes/EtOAc – CAM stain). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.35 (m, 5H), 5.61 (dd, J = 11.0, 7.5 Hz, 1H), 5.27 (ddd, J = 11.0, 8.5, 1.5 Hz, 1H), 5.08 (s, 2H), 4.85 (br, 1H), 4.65 (br, 1H), 4.42 (br, 1H), 2.12 (br, 1H), 1.90 (m, 2H), 1.61 (m, 2H), 1.49 (m, 2H), 1.38 – 1.25 (m, 1H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  155.6, 136.4, 134.9, 129.7, 128.5, 128.1, 69.4, 66.7, 49.3, 38.3, 36.5, 24.0, 23.3 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>16</sub>H<sub>22</sub>NO<sub>3</sub><sup>+</sup>, 276.1600; obsd, 276.1592.



## (±)-Benzyl (1*R*,4*S*,*Z*)-4-(tert-butyldimethylsilyloxy)cyclooct-2-enylcarbamate (3.81). Compound 3.81 was prepared following the same procedure for 3.79. Alcohol 3.80 (3.00 g, 10.9 mmol), TBDMSC1 (3.32 g, 22.0 mmol), and imidazole (2.98 g, 43.8 mmol) provided 4.5 g of crude material. Chromatography through 200 g of silica using a solvent gradient from 90% CH<sub>2</sub>Cl<sub>2</sub>/hexanes to 100% CH<sub>2</sub>Cl<sub>2</sub> yielded 3.81 as a colorless oil that solidified upon standing (3.96 g, 93%). mp = 80-82 °C. Rf = 0.50 (2:1 hexanes/EtOAc – CAM stain). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) $\delta$ 7.35 (m, 5H), 5.58 (m, 1H), 5.21 (ddd, *J* = 10.5, 8.0, 1.5 Hz, 1H), 5.11 (m, 2H), 4.81 (br, 1H), 4.57 (br, 1H), 4.43 (br, 1H), 1.90 – 1.82 (m, 2H), 1.60 – 1.48 (m, 5H), 1.32 (m, 1H), 0.90 (s, 9H), 0.10 (s, 3H), 0.09 (s, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) $\delta$ 155.4, 136.1, 128.5, 128.3, 128.0, 70.2, 66.6, 49.4, 39.2, 36.7, 25.8, 24.1, 23.4, 18.2, -4.6, -4.9 ppm. HRMS (FAB) *m*/*z* [M+H]<sup>+</sup> calcd for C<sub>22</sub>H<sub>36</sub>NO<sub>3</sub>Si<sup>+</sup>, 390.2464; obsd, 390.2458.

Attempted Kornblum-DeLaMare rearrangement of 3.40. Cycloadduct 3.40 (0.108 g, 0.451 mmol) was dissolved in 5 mL of anhydrous  $CH_2Cl_2$  in a flame-dried single-necked 25-mL round-bottomed flask under Ar. Dry triethylamine (0.020 mL, 0.14 mmol) was added and the solution was stirred at RT under Ar. After 2 days at RT, only 3.40 was observed by TLC. No change was observed after stirring at RT for a few additional days or after the addition of excess triethylamine.



(±)-(2*S*,7*R*)-Dimethyl 2-acetoxy-7-(*tert*-butoxycarbonylamino)octanedioate

(3.86). Compound 3.77 (507.8 mg, 1.792 mmol) was dissolved in 17.5 mL of  $CCl_4/CH_3CN/H_2O$  (2:2:3) in a 50-mL round-bottomed flask, and cooled to 4°C in an ice/H<sub>2</sub>O bath. NaIO<sub>4</sub> (1.542 g, 7.21 mmol) was added to the reaction, followed by RuCl<sub>3</sub> hydrate (15 mg, 0.072 mmol). The reaction immediately turned light brown. After 1 h the reaction mixture was still brown and was stirred at RT. After 3.5 h, the reaction had become jet black in color. 20 mL of 10% Na<sub>2</sub>S<sub>2</sub>O<sub>5</sub> was added to the reaction and the mixture was stirred for 10 min before filtering through a pad of celite and rinsing the reaction flask with 50 mL of EtOAc. The layers in the filtrate were separated and the aqueous layer (dark green in color, pH 2-3 by pH paper) was extracted with EtOAc (3 x 50 mL). The combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated

to yield a brown oil (720 mg). The crude material was dissolved in  $Et_2O$  (15 mL) in a 50mL flask and treated with excess diazomethane, prepared from Diazald as described in the general procedure, at 4 °C (ice/H<sub>2</sub>O bath). Bubbling was observed in the reaction flask during the addition of fresh diazomethane solution. The reaction was stirred for an 15 min at 4 °C, then was quenched by adding 10 vol% HOAc in Et<sub>2</sub>O slowly until no bubbles were observed in the reaction. The light brown solution was poured into 50 mL of saturated NaHCO<sub>3</sub> and the layers were separated. The aqueous layer (pH 8 - pH paper) was extracted with Et<sub>2</sub>O (2 x 20 mL), and the combined Et<sub>2</sub>O layers were washed with brine (20 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a brown oil ( $\sim$ 640 mg). The crude material was chromatographed through 75 g of silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 80% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to yield **3.86** as a yellow oil (503.5 mg, 74.8% yield) as well as minor amounts of two additional products (24.5 mg and 18 mg, respectively). The byproducts were not fully characterized. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 5.04 (m, 1H), 4.92 (m, 1H), 4.23 (m, 1H), 3.67 (s, 6H), 2.07 (s, 3H), 1.80 – 1.72 (m, 3H), 1.63 – 1.53 (m, 2H), 1.42 – 1.28 (m, 12H) ppm. <sup>13</sup>C NMR (125 MHz. CDCl<sub>3</sub>) δ 173.1, 170.5, 170.3, 155.2, 82.2, 79.7, 77.9, 71.9, 53.1, 52.1, 32.3, 30.7, 28.2, 24.7, 24.6, 20.4 ppm. MS (FAB) m/z [M+H]<sup>+</sup> at 376; 320, 276 (base), 260, 216.

*N*'-Benzylidene-4-methylbenzenesulfonohydrazide (3.87). *p*-Toluenesulfonylhydrazide (14.24 g, 76.5 mmol) was suspended in 25 mL of MeOH in a 125-mL Erlenmayer flask. Benzaldehyde (7.0 mL, 68.9 mmol), purified from the benzaldehyde-bisulfite adduct, was added rapidly to the slurry while stirring. The solid material dissolved after 1 min and a mildly exothermic reaction was observed. The yellow solution was left to stand and reach RT. After a few minutes, the product began to crystallize out of the solution. After 40 min, the mixture was filtered under vacuum and the white crystals were washed with a small amount of cold methanol. The crystals were dried under the aspirator vacuum to yield **3.87** as white needle-like crystals (15.675 g, 83% yield). mp = 126-128 °C (lit. 124-125 °C).

General procedure for preparation of phenyldiazomethane from 3.87. A freshly prepared solution of 1.05 M sodium methoxide in MeOH (5.30 mL, 5.57 mmol) was added to 3.87 (1.48 g, 5.39 mmol) in a single-necked 25-mL round-bottomed flask. The mixture was swirled until the solid dissolved and the resultant yellow solution was concentrated by rotary evaporation (45 °C, 21 mm Hg) to yield a white solid. The solid was dried under vacuum (<1 mm Hg) for 30 min. The flask was fitted with a short-path distillation apparatus and a receiver flask, then evacuated (<1 mm Hg). The receiver flask was cooled to -50 to -60 °C in a dry ice/acetone bath and a safety sheild was set in front of the apparatus. The flask containing the hydrazone salt was heated in an oil bath. The oil temperature was raised slowly to  $\sim 200 \,^{\circ}$ C over 45-60 min. At an oil temperature of  $\sim 110 \,^{\circ}$ C, red phenyldiazomethane (3.88) was observed collecting in the receiver flask. At an oil temperature of  $\sim 200$  °C, phenyldiazomethane was not observed collecting in the receiver flask anymore and the oil bath was allowed to cool to < 100 °C before the apparatus was disassembled. **3.88** was used immediately without warming to RT (it has been reported to be explosive at room temperature). Workup procedures for reactions utilizing phenyldiazomethane follow the same procedure used for diazomethane

reactions. All remaining and excess diphenyldiazomethane was quenched by adding a dilute solution of acetic acid in ether.

General procedure for oxidative alkene cleavage reactions using OsO<sub>4</sub>/Oxone. Attempted oxidative cleavage of 3.77 (Table 3.6, entry 10). Compound 3.77 (148.1 mg, 0.523 mmol) was dissolved in 3 mL of anhydrous DMF in a flame-dried 10-mL round-bottomed flask under Ar. OsO<sub>4</sub> (0.07 mL, 2.5 wt% in *t*BuOH, 0.006 mmol) was added to the solution (the reaction turned slightly yellow). After 5 min, Oxone (1.285 g, 2.09 mmol) was added to the reaction in one portion and the mixture was stirred vigorously at RT under Ar. The reaction was monitored by TLC (1:1 hexanes/EtOAc - CAM stain). After 4 h, some products were observed, but not the desired dicarboxylic acid products (Rf was too high, just below **3.77**), as well as remaining **3.77**. After stirring overnight at RT, no change was observed. The reaction was quenched by adding 6 equiv of Na<sub>2</sub>SO<sub>3</sub> and stirring at RT. Reactions using added sodium bicarbonate followed the same procedure except for the addition of 7 equivalents of NaHCO<sub>3</sub> to the reaction mixture.

General procedure for attempted dihydroxylation of an olefin using OsO<sub>4</sub>/NMO. Attempted dihydroxylation of 3.77. Compound 3.77 (117.3 mg, 0.414 mmol) was dissolved in 4 mL of THF in a single-necked 25-mL round-bottomed flask. NMO (50 wt% in H<sub>2</sub>O, 0.18 mL, 0.87 mmol) was added to the reaction followed by 2 drops of OsO<sub>4</sub> (2.5 wt% in *t*BuOH). The solution was stirred at RT. TLC analysis (1:1

hexanes/EtOAc - CAM stain) of the solution after 4 h showed only starting material. After stirring at RT for 2 days, no change was observed.

General procedure for attempted dihydroxylation of an olefin using RuCl<sub>3</sub>/NaIO<sub>4</sub>/CeCl<sub>3</sub>. Attempted dihydroxylation of 3.79. H<sub>2</sub>O (0.10 mL) was added to NaIO<sub>4</sub> (65.2 mg, 0.305 mmol) and CeCl<sub>3</sub>•7H<sub>2</sub>O (10.5 mg, 0.028 mmol) in a 1-dram vial. The mixture was heated until the color changed to bright yellow, then was cooled in a crushed ice/H<sub>2</sub>O bath. CH<sub>3</sub>CN (0.30 mL) was added, followed by a 0.1 M aqueous solution of RuCl<sub>3</sub> (0.01 mL, 0.001 mmol). After stirring for 2 min, **3.79** (71 mg, 0.20 mmol) was added to the brown suspension as a solution in EtOAc (0.30 mL). The mixture was stirred in the ice/H<sub>2</sub>O bath and analyzed by TLC (1:1 hexanes/EtOAc - CAM stain) for the disappearance of **3.79**. After 15 min, TLC analysis indicated the reaction was progressing and two new, more polar spots were observed by TLC, but **3.79** still remained. After 45 min, the initial two new spots had not increased in intensity, but about 4 new spots started to emerge and a complex mixture was obtained.



(±)-Dimethyl 2-(*tert*-butoxycarbonylamino)heptanedioate (3.90). Compound 3.77 (734.5 mg, 2.592 mmol) was dissolved in 21 mL of  $CH_2Cl_2$ . 5.2 mL of 2.5M methanolic KOH (13 mmol) was added and the colorless solution immediately became light yellow. The solution was cooled in a dry ice/acetone bath and  $O_3/O_2$  was bubbled

through the solution. The reaction immediately turned a deep tangerine-orange color and a yellow/orange precipitate formed in the reaction. After 1 h, the ozone bubbling was stopped and the reaction was diluted with Et<sub>2</sub>O (15 mL) then H<sub>2</sub>O until the reaction became colorless and the solid material dissolved completely. The tangerine/orange suspension became a colorless solution as the reaction was warmed to RT. The layers were separated and the aqueous layer was extracted with  $Et_2O$  (3 x 40 mL), and the combined organic layers were washed with brine (2 x 30 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a light yellow oil (708 mg). TLC analysis (1:1 hexanes/EtOAc - CAM stain) shows clean conversion to a mixture of 2 compounds. The crude material was chromatographed through a 40S Biotage column (40g of silica) using a solvent gradient from 70% hexanes/EtOAc to 50% hexanes/EtOAc to yield 3.69 as a residue (74 mg, 8%) and **3.90** as a light yellow oil (309 mg, 39%). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  5.00 (d, J = 8.0 Hz, 1H), 4.29 (dd, J = 13.5, 8.0 Hz, 1H), 3.73 (s, 3H), 3.66 (s, 3H), 2.30 (t, J = 7.5 Hz, 2H), 1.86 – 1.76 (m, 1H), 1.66 – 1.59 (m, 4H), 1.44 (s, 9H), 1.40 - 1.34 (m, 1H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 173.8, 173.2, 155.3, 79.8, 53.2, 52.2, 51.5, 33.7, 32.4, 28.2, 24.8, 24.4 ppm. MS (FAB) m/z [M+H]<sup>+</sup> at 304, 248, 204 (base), 144.

General procedure for removal of an acetate using  $K_2CO_3$ . Compound 3.69 from saponification of 3.86. Compound 3.86 (0.398 g, 1.06 mmol) was dissolved in 11 mL of anhydrous methanol in a 100-mL round-bottomed flask under Ar. Potassium carbonate (0.208 g, 1.51 mmol) was added and the yellow mixture was stirred at RT under Ar. The reaction was monitored by TLC (1:1 hexanes/EtOAc - CAM stain). After 30 min, only a small amount of **3.86** was observed and a more polar spot was observed. After 1.5 h, the reaction was partitioned between  $H_2O$  (25 mL), brine (25 mL), and EtOAc (30 mL). The layers were separated and the aqueous layer was extracted with EtOAc (2 x 30 mL). The combined organic layers were washed with brine (25 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield **3.69** as a yellow oil (0.277 g, 78% yield). See above for characterization data.



(±)-(2*S*,7*R*)-Dimethyl 2-(benzyloxy)-7-(*tert*-butoxycarbonylamino) octanedioate (3.91). Compound 3.91 was prepared following the same procedure used for compound 3.90. The use of 3.78 (469 mg, 1.41 mmol) provided a colorless oil (471 mg). The oil was loaded onto silica and chromatographed through a Biotage 40S column (40 g of silica) using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 90% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to yield 3.91 as a yellow oil (118.5 mg, 20%). The product was not pure by NMR; however, spectral information for the appropriate NMR signals are presented here from the impure sample: <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.34 (m, 5H), 5.05 (d, *J* = 8.1 Hz, 1H), 4.69 (d, *J* = 11.7 Hz, 1H), 4.38 (d, *J* = 11.7 Hz, 1H), 4.28 (m, 1H), 3.93 (t, *J* = 6.3 Hz, 1H), 3.75 (s, 3H), 3.72 (s, 3H), 2.36 – 2.29 (m, 1H), 1.79 – 1.60 (m, 6H), 1.44 (s, 9H), 1.43 – 1.31 (m, 2H) ppm.



## $(\pm)$ -(2R,7S)-Dimethyl

## 2-(tert-butoxycarbonylamino)-7-(tert-

**butyldimethylsilyloxy)octanedioate (3.92).** Compound **3.92** was prepared following the same procedure used for **3.90**. The use of **3.79** (514.7 mg, 1.45 mmol) provided a yellow oil. The oil was chromatographed through 65 g of silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 90% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to afford **3.92** as a pale yellow oil (351.9 mg, 54%). The product contained TBS-containing impurities by NMR. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  4.99 (br, 1H), 4.26 (br, 1H), 4.16 (t, *J* = 6.0 Hz, 1H), 3.70 (s, 3H), 3.68 (s, 3H), 2.29 (t, *J* = 7.0 Hz, 1H), 1.77 (m, 1H), 1.69 – 1.64 (m, 2H), 1.63 – 1.55 (m, 1H), 1.41 (s, 9H), 1.38 – 1.29 (m, 2H), 0.90 (s, impurity), 0.87 (s, 9H), 0.23 (s, impurity), 0.04 (s, 3H), 0.02 (s, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  174.1, 173.3, 155.3, 79.8, 72.0, 53.4, 52.1, 51.7, 35.6, 34.8, 32.6, 32.4, 28.3, 25.7, 25.5, 25.0, 24.8, 24.7, 24.6, 18.2, 17.5, -4.9, -5.0, - 5.4 ppm.



(±)-*tert*-Butyl (2*R*,7*S*)-7-(*tert*-butyldimethylsilyloxy)-1,8-dioxooctan-2ylcarbamate (3.93). Compound 3.79 (149.6 mg, 0.421 mmol) was dissolved in 5 mL of

4:1 CH<sub>2</sub>Cl<sub>2</sub>/MeOH in a single-necked 25-mL round-bottomed flask. The solution was cooled in a dry ice/actone bath to -78 °C and a stream of ozone was bubbled through the solution. The blue color of ozone was observed in the solution after  $\sim 5$  min, and the ozone flow was stopped after 10 min. The solution was purged of ozone by bubbling Ar through the solution until the blue color disappated (5 min), and polymer-supported triphenylphosphine (0.8 mmol/g, 1.2 g, 0.96 mmol) was added to the reaction. The mixture was removed from the dry ice/acetone bath and stirred at RT for 1.5 h. The PS-PPh<sub>3</sub> was removed by filtration and washed with  $CH_2Cl_2$  (5-10 mL). The filtrate was concentrated to yield a light yellow oil (146 mg crude). The crude material was analyzed by TLC (2:1 hexanes/EtOAc - CAM stain) and was composed of one major spot plus some baseline material. The oil was chromatographed through 20 g of silica gel using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 85% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to afford 3.93 as a colorless oil (113.0 mg, 69.3%). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  9.56 (d, J = 1.0 Hz, 1H), 9.55 (s, 1H), 5.07 (d, *J* = 6.5 Hz, 1H), 4.19 (dd, *J* = 7.0, 6.0 Hz, 1H), 3.99 (m, 1H), 1.88 (m, 1H), 1.61 (m, 2H), 1.43 (s, 9H), 1.37 (m, 4H), 0.90 (s, 9H), 0.06 (s, 3H), 0.05 (s, 3H) ppm.  $^{13}C$  NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  204.1, 199.7, 155.5, 80.0, 77.3, 59.7, 32.3, 29.0, 28.2, 25.7, 25.0, 24.4, 18.1, -4.7, -5.0 ppm. HRMS (FAB) m/z [M]<sup>+</sup> calcd for C<sub>19</sub>H<sub>37</sub>NO<sub>5</sub>Si<sup>+</sup>, 387.2441; obsd, 387.2446.



(2R,7S)-7-(tert-butyldimethylsilyloxy)-1,8-dihydroxyoctan-2-(±)-*tert*-Butyl ylcarbamate (3.94). Compound 3.79 (3.035 g, 8.535 mmol) was dissolved in 100 mL of 4:1 CH<sub>2</sub>Cl<sub>2</sub>/MeOH in a single-necked 250-mL round-bottomed flask. The solution was cooled in a dry ice/actone bath to -78 °C and a stream of ozone was bubbled through the solution. The blue color of ozone was observed in the solution after ~10 min, and the ozone flow was stopped. The solution was purged of ozone by bubbling Ar through the solution until the blue color disappated (20 min). The reaction was removed from the dry ice/acetone bath and placed in a crushed ice/NaCl/H<sub>2</sub>O bath, at which time NaBH<sub>4</sub> (2.03, 53.7 mmol) was added to the reaction in portions over 60 min. Bubbling was observed and the reaction was allowed to stir in the ice/NaCl/H2O bath for 1 h, then warmed to RT and stirred for an additional 1 h. The reaction was diluted with H<sub>2</sub>O (100 mL) and EtOAc (200 mL). The layers were separated and the aqueous layer was extracted with EtOAc (3 x 100 mL). The combined organic layers were washed with brine (2 x 100 mL). mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a colorless oil. The oil was chromatographed through 250 g of silica gel using a solvent gradient from 50% hexanes/EtOAc to 30% hexanes/EtOAc to yield 3.94 as a colorless oil (3.30 g, 99%). Rf = 0.13 (1:1 hexanes/EtOAc - CAM stain). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  4.77 (d, J = 8.0 Hz, 1H), 3.68 (m, 1H), 3.58 (m, 2H), 3.50 (m, 2H), 3.41 (m, 1H), 3.08 (br-s, 1H), 2.24 (br-s, 1H), 1.46 (m, 3H), 1.40 (s, 9H), 1.33-1.22 (m, 5H), 0.86 (s, 9H), 0.04 (s, 6H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 156.4, 79.4, 72.6, 66.0, 65.5, 52.5, 33.7, 31.4, 28.3,

26.1, 25.8, 25.0, 18.0, -4.6, -4.7 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for  $C_{19}H_{42}NO_5Si^+$ , 392.2832; obsd, 392.2838.



(±)-Benzyl (2R,7S)-7-(tert-butyldimethylsilyloxy)-1,8-dihydroxyoctan-2ylcarbamate (3.95). Compound 3.95 was prepared following the procedure for 3.94. 3.81 (1.949 g, 5.003 mmol) was treated with ozone at -78 °C, then NaBH<sub>4</sub> (1.23 g, 32.5 mmol) at -10 °C. The crude material was purified through 100 g of silica gel using a solvent gradient from 50% EtOAc/hexanes to 60% EtOAc/hexanes to afford 3.95 as a colorless oil (2.02 g, 95%). Rf = 0.11 (1:1 hexanes/EtOAc – CAM stain). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.35 (m, 5H), 5.09 (s, 2H), 4.91 (d, *J* = 7.5 Hz, 1H), 3.0 (m, 3H), 3.56 (m, 2H), 3.43 (m, 1H), 2.33 (br-s, 1H), 1.96 (t, *J* = 6.5 Hz, 1H), 1.49-1.25 (m, 8H), 0.89 (s, 9H), 0.07 (s, 6H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  156.7, 136.3, 128.5, 128.2, 128.1, 72.6, 66.8, 66.1, 65.5, 53.1, 33.7, 31.3, 26.1, 25.9, 25.1, 18.1, -4.5, -4.6 ppm.



(±)-*tert*-Butyl (2*R*,7*S*)-1,7,8-trihydroxyoctan-2-ylcarbamate (3.96). Compound 3.94 (0.418 g, 1.07 mmol) was dissolved in 5 mL of anhydrous THF in a flame-dried

single-necked 25-mL round-bottomed flask under Ar. TBAF (1.0 M, 1.60 mL, 1.60 mmol) was added dropwise and the resultant light-yellow solution was stirred at RT under Ar. After 4.5 h, the reaction was complete by TLC analysis (9:1 EtOAc/MeOH - CAM stain). The reaction was concentrated by rotary evaporation (40-45 °C, 21 mm Hg), and the resultant yellow oil was chromatographed through 10 g of silica gel using a solvent gradient from 100% EtOAc to 90% EtOAc/MeOH to afford 80 as a colorless oil (285.5 mg, 87%). The oil solidified upon standing for ~1 week. <sup>1</sup>H NMR (500 MHz, d<sup>6</sup>-DMSO)  $\delta$  6.32 (d, *J* = 8.0 Hz, 1H), 4.45 (t, *J* = 5.5 Hz, 1H), 4.32 (t, *J* = 5.5 Hz, 1H), 4.22 (d, *J* = 5.0 Hz, 1H), 3.33 – 3.28 (m, 2H), 3.26 – 3.19 (m, 3H), 1.51 – 1.45 (m, 1H), 1.40 – 1.37 (m, 11H), 1.31 – 1.27 (m, 1H), 1.25 – 1.17 (m, 4H) ppm. <sup>13</sup>C NMR (125 MHz, d<sup>6</sup>-DMSO)  $\delta$  155.3, 77.1, 70.9, 65.8, 63.5, 52.1, 33.3, 30.9, 28.1, 25.6, 25.1 ppm. HRMS (FAB) *m*/z [M+H]<sup>+</sup> calcd for C<sub>13</sub>H<sub>28</sub>NO<sub>5</sub><sup>+</sup>, 278.1967; obsd, 278.1952.



(±)-Benzyl (2R,7S)-1,7,8-trihydroxyoctan-2-ylcarbamate (3.97). Compound 3.97 was prepared following the procedure for 3.96. Compound 3.95 (1.00 g, 2.35 mmol) was treated with 3.0 mL of 1.0 M TBAF (3.0 mmol). The crude material was purified through 75 g of silica gel using a solvent gradient from 100% EtOAc to 90% EtOAc/MeOH to yield 3.97 as a colorless oil that solidified upon standing (0.641 g, 88%). mp = 81-83 °C. Rf = 0.26 (9:1 EtOAc/MeOH – CAM stain). <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD)  $\delta$  7.35 – 7.28 (m, 5H), 5.07 (s, 2H), 3.58 (br-m, 2H), 3.49 – 3.39 (m, 4H), 1.62 – 1.30 (m, 8H) ppm. <sup>13</sup>C NMR (125 MHz, CD<sub>3</sub>OD) δ 158.9, 138.5, 129.5, 129.0, 128.8, 73.2, 67.40, 67.36, 65.4, 54.4, 34.4, 32.3, 27.2, 26.6 ppm.



(±)-tert-Butyl (R)-6-((S)-2,2-dimethyl-1,3-dioxolan-4-yl)-1-hydroxyhexan-2ylcarbamate (3.98). Compound 3.96 (1.499 g, 5.405 mmol) was dissolved in 55 mL of anhydrous THF in a flame-dried single-necked 250-mL round-bottomed flask under Ar. 2,2-Dimethoxypropane (0.70 mL, 5.7 mmol) was added to the reaction, followed by anhydrous p-toluenesulfonic acid (48.1 mg, 0.279 mmol). The reaction was stirred at RT under Ar for 6 h, at which time TLC analysis of the reaction mixture (1:1 hexanes/EtOAc and 9:1 EtOAc/MeOH - CAM stain) only shows a small amount of starting material and mostly one product. About 300 mg of solid sodium carbonate was added to the reaction and the mixture was concentrated by rotary evaporation (30-35 °C, 21 mm Hg) to a yellow oil. Saturated NaHCO<sub>3</sub> (50 mL), H<sub>2</sub>O (20 mL), and EtOAc (75 mL) were added to the oil and the layers were separated. The combined organic layers were washed with H<sub>2</sub>O (50 mL) and brine (50 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to yield a yellow oil. The oil was purified through 150 g of silica gel using a solvent gradient from 75% hexanes/EtOAc to 33% hexanes/EtOAc to yield a light yellow oil. The oil was dried under vacuum (1.5 mm Hg) overnight to afford **3.98** as a waxy white solid (1.64 g, 96%). Rf = 0.14 (1:1 hexanes/EtOAc – CAM stain). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  4.73 (d, J =

7.0 Hz, 1H), 4.10 – 3.98 (m, 2H), 3.62 – 3.54 (br-m, 2H), 3.51 – 3.47 (m, 1H), 3.46 (t, J = 7.0 Hz, 1H), 2.83 (br, 1H), 1.61 – 1.57 (m, 1H), 1.52 -1.35 (m, 7H), 1.41 (s, 9H), 1.37 (s, 3H), 1.32 (s, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  156.4, 108.6, 79.5, 75.9, 69.4, 65.7, 52.6, 33.3, 31.3, 28.3, 26.9, 25.9, 25.7, 25.6 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>16</sub>H<sub>32</sub>NO<sub>5</sub><sup>+</sup>, 318.2280; obsd, 318.2259.



(±)-Benzyl (*R*)-6-((*S*)-2,2-dimethyl-1,3-dioxolan-4-yl)-1-hydroxyhexan-2ylcarbamate (3.99). Compound 3.99 was prepared following the same procedure for 3.98. The use of 3.97 (0.641 g, 2.06 mmol), 2,2-dimethoxypropane (0.35 mL, 2.85 mmol), and *p*-toluenesulfonic acid hydrate (27.8 mg, 0.146 mmol) provided a yellow oil. The oil was chromatographed through 75 g of silica using a solvent gradient from 75% hexanes/EtOAc to 33% hexanes/EtOAc and yielded **3.99** as an oil that solidified to a waxy solid upon standing (0.614 g, 85%). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.35 – 7.30 (m, 5H), 5.09 (s, 2H), 4.95 (d, *J* = 7.5 Hz, 1H), 4.06 – 3.99 (m, 2H), 3.67 (br-m, 2H), 3.55 (br-m, 1H), 3.47 (t, *J* = 7.5 Hz, 1H), 2.41 (br, 1H), 1.64 -1.24 (m, 8H), 1.39 (s, 3H), 1.34 (s, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  156.7, 136.3, 128.5, 128.14, 128.08, 108.7, 75.9, 69.4, 66.8, 65.4, 53.1, 33.3, 31.2, 26.9, 25.9, 25.7, 25.6 ppm.



(±)-*tert*-Butyl (R)-6-((S)-2,2-dimethyl-1,3-dioxolan-4-yl)-1-oxohexan-2ylcarbamate (3.100). Alcohol 3.98 (0.5158 g, 1.625 mmol) was dissolved in 10 mL of anhydrous CH<sub>2</sub>Cl<sub>2</sub> in a flame-dried single-necked 50-mL round-bottomed flask under Ar. Dess-Martin periodinane (15 wt% in CH<sub>2</sub>Cl<sub>2</sub>, 5.10 mL, 2.40 mmol) was added and the resultant cloudy yellow mixture was stirred at RT. The reaction was monitored by TLC (1:1 hexanes/EtOAc - CAM stain). After 4 h, starting material remained and additional DMP was added (1 mL, ~0.3 eq). After stirring for 45 min, 40 mL of 1:1 saturated NaHCO<sub>3</sub>/10 wt% Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> was added to the reaction along with 10 mL of EtOAc. The mixture was stirred at RT for 30 min until the solid material had completely dissolved. The layers were separated and the aqueous layer was extracted with EtOAc (3 x 50 mL). The combined organic layers were washed with saturated NaHCO<sub>3</sub> (3 x 50 mL),  $H_2O$  (50 mL) and brine (50 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to yield a yellow oil. The oil was purified through 50 g of silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 80% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to afford **3.100** as a light yellow oil that solidified upon standing to a waxy solid (0.489 g, 95%). Rf = 0.40 (1:1 hexanes/EtOAc - CAM stain). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  9.57 (br-s, 1H), 5.08 (d, J = 6.0 Hz, 1H), 4.21 (dd, J = 6.5, 6.0 Hz, 1H), 4.06-4.00 (m, 2H), 3.48 (t, J = 7.0 Hz, 1H), 1.88 (br-m, 1H), 1.60-1.46 (m, 5H), 1.44 (s, 9H), 1.39 (s, 3H), 1.34 (s, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 199.8,
155.5, 108.7, 80.1, 75.8, 69.4, 59.7, 33.2, 29.0, 28.3, 26.9, 25.7, 25.6, 25.1 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>16</sub>H<sub>30</sub>O<sub>5</sub>N<sup>+</sup>, 316.2124; obsd, 316.2119.

Aldehyde 3.100 from alcohol 3.98 by Swern oxidation. Oxalyl chloride (0.20 mL, 2.3 mmol) was added to 5 mL of anhydrous CH<sub>2</sub>Cl<sub>2</sub> in a flame-dried, single-necked 50-mL round-bottomed flask cooled to -78 °C (dry ice/acetone) under Ar. In a separate flame-dried single-necked 10-mL round-bottomed flask was dissolved anhydrous dimethyl sulfoxide (0.35 mL, 4.9 mmol) in 5 mL of anhydrous CH<sub>2</sub>Cl<sub>2</sub>. The DMSO solution was added to the oxalyl chloride solution slowly via cannula and the resultant solution was stirred at -78 °C for 20 min under Ar. In a separate flame-dried singlenecked 25-mL round-bottomed flask under Ar was dissolved 3.98 (0.507 g, 1.60 mmol) in 10 mL of anhydrous  $CH_2Cl_2$ . This solution was transferred dropwise to the reaction flask via cannula. The resultant mixture was stirred at -78 °C for 1 h, then Et<sub>3</sub>N (1.25 mL, 8.89 mmol) was added. The mixture was stirred for 1 h at -78 °C, then the dry ice/acetone bath was removed and the mixture was allowed to slowly reach RT under Ar. After 1 h, the reaction mixture was diluted with H<sub>2</sub>O (25 mL) and CH<sub>2</sub>Cl<sub>2</sub> (10 mL) and the layers were separated. The aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (2 x 25 mL) and the combined  $CH_2Cl_2$  layers were washed with brine (30 mL), dried over MgSO<sub>4</sub>, and concentrated to yield an orange oil. The crude material was chromatographed through 50 g of silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 75% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc and yielded **3.100** as a yellow oil (0.460 g, 91.3% yield).



(±)-(*R*)-2-(*tert*-Butoxycarbonylamino)-6-((*S*)-2,2-dimethyl-1,3-dioxolan-4yl)hexanoic acid (3.101). Oxidation using KMnO<sub>4</sub>. Compound 3.100 (0.1592 g, 0.5048 mmol) was dissolved in 10 mL of MeOH in a single-necked 50-mL roundbottomed flask. А solution of  $KMnO_4$  (0.185 g, 1.17 mmol) and benzyltriethylammonium chloride (5 mg, 0.02 mmol) in 15 mL of H<sub>2</sub>O was added and the mixture was stirred vigorously at RT. The temperature of the mixture increased slightly and the color of the mixture changed from deep purple to a brownish/purple color. After 5 h, the reaction was filtered. EtOAc (20 mL) and H<sub>2</sub>O (10 mL) were added to the murky brown filtrate. The aqueous layer was acidified to pH 3 using 6M HCl (3) drops). 10 wt% sodium meta bisulfite (about 5 mL) was added and the brown mixture turned colorless. The layers were separated and the aqueous was quickly extracted with EtOAc ( $3 \times 10 \text{ mL}$ ). The combined organic layers were dried over MgSO<sub>4</sub>, filtered, and concentrated to afford a colorless oil. The oil was dried under vacuum (1.5 mm Hg) to afford a colorless oil (152.5 mg, 91.2% yield). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  9.93 (br-s, 1H), 6.31 (m, 1H), 5.11 (d, J = 8.1 Hz, 1H), 4.27 (m, 1H), 4.07-3.99 (m, 3H), 3.47 (t, J =6.9 Hz, 1H), 1.82 (m, 1H), 1.66-1.57 (m, 3H), 1.41-1.32 (m, 2H), 1.42 (s, 9H), 1.38 (s, 3H), 1.32 (s, 3H) ppm. <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 176.8, 155.5, 108.7, 80.0, 75.8, 69.3, 53.1, 33.2, 33.1, 32.2, 28.2, 26.8, 25.6, 25.3, 25.2 ppm.

Oxidation conditions for aldehyde 3.100 using NaClO<sub>2</sub> (Table 3.7, entry 2). Aldenhyde 3.100 (55.0 mg, 0.174 mmol) was dissolved in 5 mL of 1:1 *t*BuOH/H<sub>2</sub>O in a 20-mL scintillation vial. 2-Methyl-2-butene (0.40 mL, 3.8 mmol) was added followed by sodium chlorite (51.9, 0.574 mmol) and monobasic sodium phosphate (63.9 mg, 0.463 mmol). The mixture was stirred at RT for about 6 h, then partitioned between H<sub>2</sub>O (5 mL) and EtOAc (5 mL). The aqueous layer was carefully acidified to a pH of 2-3 (pH paper) using 1M HCl and the layers were separated. The aqueous layer was extracted with EtOAc (2 x 5 mL) and the combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated.

Oxidation conditions for aldehyde 3.100 using Oxone (Table 3.7, entry 3). Aldehyde 3.100 (72.8 mg, 0.231 mmol) was dissolved in 4 mL of DMF in a 20-mL scintillation vial. Oxone (160 mg, 0.26 mmol) was added in one portion and the mixture was stirred at RT for 5 h. The reaction was partitioned between  $H_2O$  (10 mL) and EtOAc (10 mL). The aqueous layer was a pH of 2-3 (pH paper). The layers were separated and the aqueous layer was extracted with EtOAc (2 x 10 mL). The combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to afford a residue.

Oxidation conditions for aldehyde 3.100 using Ag<sub>2</sub>O (Table 3.7, entry 5). Aldehyde 3.100 (59.7 mg, 0.189 mmol) was dissolved in 3 mL of EtOH in a 20 mL scintillation vial. A solution of silver nitrate (82 mg, 0.48 mmol) in 0.5 mL of H<sub>2</sub>O was added, followed by the dropwise addition of a solution of KOH (62 mg, 1.1 mmol) in 1.5 mL of H<sub>2</sub>O. A brown solid precipitated out of solution immediately. The mixture was stirred at RT for about 4 h. The brown solid was removed by filtration and the filtrate was diluted with EtOAc (5 mL) and  $H_2O$  (5 mL). The aqueous layer was carefully acidified to a pH of about 2-3 (pH paper) using 1M HCl. The layers were separated and the aqueous layer was extracted with EtOAc (2 x 5 mL). The combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to afford a residue.



(±)-(*R*)-Methyl 2-(*tert*-butoxycarbonylamino)-6-((*S*)-2,2-dimethyl-1,3dioxolan-4-yl)hexanoate (3.102). Crude 3.101, prepared as described above from oxidation of 3.100 (79.8 mg, 0.253 mmol) using KMnO<sub>4</sub>, was dissolved in about 10 mL of Et<sub>2</sub>O and 2 mL of THF and treated with an excess of diazomethane in Et<sub>2</sub>O. The resultant yellow solution was quenched with a 10 vol% acetic acid/Et<sub>2</sub>O solution until the yellow color disappeared. Saturated NaHCO<sub>3</sub> (20 mL) was added and the layers were separated. The aqueous layer was extracted with EtOAc (2 x 5 mL) and the combined organic layers were washed with brine (5 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. The crude material was chromatographed through 5 g of silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 90% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to afford **3.102** as a yellow oil (63.2 mg, 72.3% yield). Rf = 0.46 (1:1 hexanes/EtOAc – CAM stain). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  5.03 (d, *J* = 8.5 Hz, 1H), 4.26 (dd, *J* = 13.0, 7.5 Hz, 1H), 4.06-3.99 (m, 2H), 3.71 (s, 3H), 3.47 (t, *J* = 7.0 Hz, 1H), 1.90-1.75 (m, 1H), 1.70-1.55 (m, 2H), 1.501.30 (m, 3H), 1.42 (s, 9H), 1.38 (s, 3H), 1.32 (s, 3H) ppm.  $^{13}$ C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  173.3, 155.3, 108.6, 79.8, 75.8, 69.4, 53.3, 52.2, 33.2, 32.5, 28.3, 26.9, 25.7, 25.3, 25.2 ppm. HRMS (FAB) *m*/*z* [M+H]<sup>+</sup> calcd for C<sub>17</sub>H<sub>32</sub>NO<sub>6</sub><sup>+</sup>, 346.2230; obsd, 346.2219.



(±)-(R)-2-(tert-butoxycarbonylamino)-6-((S)-2,2-dimethyl-1,3-dioxolan-4-

yl)hexyl acetate (3.105). Compound 3.105 was prepared following the same procedure used for 3.77. The use of 3.98 (0.503 g, 1.58 mmol) and acetic anhydride (0.40 mL, 4.2 mmol) provided a yellow oil. The oil was chromatographed through a Biotage 40S column using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 85% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc and yielded 3.105 as a colorless oil. The oil was dried under vacuum overnight to provide 3.105 (0.529 g, 92.9% yield). Rf = 0.44 (1:1 hexanes/EtOAc – CAM stain). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 4.53 (d, *J* = 9.0 Hz, 1H), 4.04 (m, 4H), 3.83 (br, 1H), 3.49 (t, *J* = 6.9 Hz, 1H), 2.07 (s, 3H), 1.70 – 1.35 (m, 8H), 1.45 (s, 9H), 1.41 (s, 3H), 1.35 (s, 3H) ppm. <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 170.9, 155.4, 108.6, 75.9, 69.4, 66.3, 49.4, 33.3, 31.7, 28.3, 26.9, 25.8, 25.7, 25.6, 20.8 ppm. HRMS (FAB) *m*/*z* [M+H]<sup>+</sup> calcd for C<sub>18</sub>H<sub>34</sub>NO<sub>6</sub><sup>+</sup>, 360.2386; obsd, 360.2388.



## (±)-(2R,7S)-2-(tert-Butoxycarbonylamino)-7,8-dihydroxyoctyl acetate (3.106).

Compound 3.105 (0.253 g, 0.703 mmol) was dissolved in 6 mL of MeOH in a 20-mL scintillation vial. Pyridinium para-toluenesulfonate (21.4 mg, 0.0852 mmol) was added After 1.5 h, the distinct aroma of 2,2and the mixture was stirred at RT. dimethoxypropane was detected in the reaction mixture. TLC of the reaction indicated a new more polar spot (just above baseline, 1:1 hexanes/EtOAc - CAM stain), but 3.105 remained (Rf = 0.44). After 24 h, no change was observed. The reaction was concentrated (to remove the 2,2-dimethoxypropane), and the resultant colorless oil was dissolved in 5 mL of MeOH and stirred at RT. Only a small amount of 3.105 remained after stirring for 1 day. The reaction was concentrated and the colorless oil was partitioned between EtOAc (20 mL), and H<sub>2</sub>O (10 mL). The layers were separated and the organic layer was washed with 1M HCl (3 x 5 mL),  $H_2O$  (5 mL), saturated NaHCO<sub>3</sub> (2 x 5 mL), and brine (5 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to yield a colorless oil. The oil was chromatographed through 15 g of silica using 100% EtOAc to yield **3.106** as a light yellow oil (171.1 mg, 76.2% yield). Rf = 0.2 (100% EtOAc – CAM stain). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  4.74 (d, J = 7.8 Hz, 1H), 4.04 (m, 2H), 3.83 (br, 1H), 3.69 - 3.60 (m, 2H), 3.42 (dd, J = 11.1, 7.5 Hz, 1H), 3.32 (br, 2H), 2.07 (s, 3H), 1.52 – 1.32 (m, 8H), 1.44 (s, 9H) ppm. <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 171.0, 155.6, 79.4, 72.0, 66.6, 49.3, 32.7, 28.3, 25.6, 25.2, 20.8 ppm.

Attempted oxidation of 3.106 using trichloroisocyanuric acid/TEMPO. Compound 3.106 (59.7 mg, 0.187 mmol) was dissolved in 2 mL of  $CH_2Cl_2$  in a 20-mL scintillation vial and cooled in an ice/H<sub>2</sub>O bath. Trichloroisocyanuric acid (47.0 mg, 0.202 mmol) was added, followed by TEMPO (1 mg, 0.006 mmol). The yellow reaction was stirred in the ice/H<sub>2</sub>O bath. A complex mixture was observed after 1 h.

Enzymatic resolution of ( $\pm$ )-3.76 using Novozyme 435. CH<sub>2</sub>Cl<sub>2</sub> (40 mL) and wet *n*-heptane (50 mL) were added to ( $\pm$ )-3.76 (2.268 g, 9.40 mmol) in a 250-mL erlenmayer flask. To the slurry was added vinyl acetate (2.90 mL, 31.5 mmol), followed by immobilized *C. antarctica* B lipase enzyme (Novozyme 435, 255 mg). The mixture was not completely in solution. The mixture was shaken at 37 °C overnight. After 15 h, an aliquot was removed, concentrated, and the resultant solid was analyzed by <sup>1</sup>H NMR. The reaction was determined to have proceeded to 50% conversion. The immobilized enzyme was removed from the mixture by filtration and washed with portions of EtOAc (10 mL) and CH<sub>2</sub>Cl<sub>2</sub> (10 mL). The filtrate was concentrated to yield a white solid. The solid was chromatographed through 250 g of silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 80% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to elute (+)-**3.77** and 50% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to elute (-)-**3.76**. Total recovery was 97%.



(+)-(1*R*,4*S*,*Z*)-4-(*tert*-Butoxycarbonylamino)cyclooct-2-enyl acetate (+)-3.77. Compound (+)-3.77 was isolated as above as an off-white crystalline solid (1.24 g, 46%

yield). NMR spectra matches racemic **3.77**. mp = 125-126 °C.  $[\alpha]^{23}$  = +69.6 (c = 0.53 in MeOH). HRMS (FAB) *m/z* [M+H]<sup>+</sup> calcd for C<sub>15</sub>H<sub>26</sub>NO<sub>4</sub><sup>+</sup>, 284.1862; obsd, 284.1885.



(-)-*tert*-Butyl (1*R*,4*S*,Z)-4-hydroxycyclooct-2-enylcarbamate (-)-3.76. Compound (-)-3.76 was isolated as above as a white amorphous solid (1.14 g, 50% yield). NMR spectra matches racemic 3.76. mp = 151-152 °C.  $[\alpha]^{23} = -84.1$  (c = 0.52 in MeOH). HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>13</sub>H<sub>24</sub>NO<sub>3</sub><sup>+</sup>, 242.1756; obsd, 242.1739.



(-)-(1*S*,4*R*,*Z*)-4-(*tert*-Butoxycarbonylamino)cyclooct-2-enyl acetate (-)-3.77.
Compound (-)-3.77 was prepared following the same procedure for (±)-3.77. Compound
(-)-3.76 (0.153 g, 0.632 mmol) and acetic anhydride (0.15 mL, 1.59 mmol) provided (-)3.77 as an off-white solid (0.167 g, 93% yield). NMR spectra matches racemic 3.77.



(+)-tert-Butyl (1S,4R,Z)-4-hydroxycyclooct-2-enylcarbamate (+)-3.76. General procedure for acetate removal using  $K_2CO_3$ . A 100-mL round-bottomed flask was charged with potassium carbonate (0.731 g, 5.29 mmol) and flame-dried under Ar. (+)-3.77 (1.05 g, 3.70 mmol) was added and dissolved in 20 mL of anhydrous methanol. The mixture was stirred at RT under Ar. After 2.5 h, the reaction was complete by TLC (1:1 hexanes/EtOAc - CAM stain) analysis. The yellow mixture was partitioned between EtOAc (50 mL), H<sub>2</sub>O (25 mL), and brine (25 mL) and the layers were separated. The aqueous layer was extracted with EtOAc (2 x 25 mL) and the combined organic layers were washed with H<sub>2</sub>O (25 mL) and brine (2 x 25 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a white solid. The solid was dried under vacuum overnight (1.5 mm Hg) to yield (+)-**3.76** as a white solid (0.884 g, 99% yield). NMR spectra matches racemic **3.76**. mp = 150.5-151.5 °C.  $[\alpha]^{23} = +109$  (c = 0.50 in MeOH).



(*R*)-1-(2-Nitrophenyl)pyrrolidine-2-carboxylic acid, *R*-2-NPP (3.108). *D*-Proline (1.06 g, 9.18 mmol) was dissolved in 100 mL of EtOH/H<sub>2</sub>O (1:1) in a 250-mL round-bottomed flask fitted with a stir bar and condenser. Sodium bicarbonate (2.08 g, 24.8 mmol) was added, followed by 1-fluoro-2-nitrobenzene (0.96 mL, 9.1 mmol). The reaction was heated to reflux in an oil bath (oil temperature = 95-100 °C). After about 10 min, the color of the solution changed from yellow to red. After 5 h, the mixture was cooled and stirred at RT overnight. The volume of the red solution was reduced to about 30-40 mL by rotary evaporation and the red mixture was acidified to pH 2-3 (pH paper) by adding 1M HCl (25-30 mL). Yellow solid precipitated out of the solution. The mixture was diluted with EtOAc (75 mL) and the layers were separated. The aqueous layer was extracted with EtOAc (3 x 30 mL) and the combined organic layers were dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a sticky orange oil. The oil was dried

under vacuum (~2 torr). After the sticky oil was scratched with a spatula and left to stand overnight at 4 °C, the oil solidified to yield **3.108** as a yellow solid (2.2 g, ~100% yield).  $[\alpha]^{23} = +933$  (c = 1.0 in MeOH); lit.<sup>6</sup> = +1020 (c = 1.0 in MeOH). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  10.28 (br, 1H), 7.73 (dd, *J* = 8.0, 1.5 Hz, 1H), 7.39 (tdd, *J* = 7.5, 1.5, 0.5 Hz, 1H), 6.86 (m, 2H), 4.43 (t, *J* = 7.5 Hz, 1H), 3.55 (td, *J* = 9.5, 7.5 Hz, 1H), 3.07 (ddd, *J* = 9.5, 7.5, 4.0 Hz, 1H), 2.49 (dtd, *J* = 12, 7.0, 5.0 Hz, 1H), 2.20 – 2.07 (m, 2H), 1.98 – 1.89 (m, 1H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  177.9, 141.2, 138.9, 133.2, 126.6, 118.2, 116.8, 61.7, 51.9, 30.9, 24.8 ppm.



(S)-1-(2-Nitrophenyl)pyrrolidine-2-carboxylic acid, S-2-NPP (3.109). Compound 3.109 was prepared following the same procedure used for 3.108. *L*-proline (1.05 g, 9.12 mmol), 1-fluoro-2-nitrobenzene (0.96 mL, 9.08 mmol), and sodium bicarbonate (2.0 g, 24 mmol) provided 3.109 as a sticky orange oil (2.10 g, 98% yield). NMR spectra is identical to 3.108.  $[\alpha]^{23} = -941$  (c = 1.0 in MeOH); lit. = -1080 (c = 1.0 in MeOH).



(1S,4R,Z)-4-((R)-1-(2-Nitrophenyl)pyrrolidine-2-carboxamido)cyclooct-2-enyl acetate (3.110). Compound (-)-3.77 (126 mg, 0.445 mmol) was dissolved in 5 mL of

 $CH_2Cl_2$  in a 25-mL round-bottomed flask. Triethylsilane (0.080 mL, 0.50 mmol) was added, followed by trifluoroacetic acid (0.50 mL, 6.7 mmol). The light yellow solution was stirred at RT under Ar. The reaction was monitored by TLC (1:1 hexanes/EtOAc -CAM stain, ninhydrin stain) for the disappearance of (-)-**3.77**. After 1.5 h, the reaction was still incomplete and an additional 0.5 mL (6.7 mmol) TFA was added. After 2 h, the reaction was complete and the reaction was diluted with  $CH_2Cl_2$  (10 mL) and poured into 50 mL of saturated NaHCO<sub>3</sub>. The pH of the aqueous layer was adjusted to 8 (pH paper) by adding portions of solid Na<sub>2</sub>CO<sub>3</sub>. The layers were separated and the aqueous layer was extracted with  $CH_2Cl_2$  (3 x 15 mL). The combined organic layers were dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a yellow oil (59.3 mg, 73% yield of free amine).

The free amine intermediate (28.2 mg, 0.154 mmol) and HOBt (26.2 mg, 0.194 mmol) were dissolved in 2 mL of CH<sub>3</sub>CN in a 25-mL round-bottomed flask under Ar. **3.108** (44.4 mg, 0.188 mmol) was added as a solution in 3 mL of dry CH<sub>3</sub>CN. EDC•HCl (44 mg, 0.23 mmol) was added to the yellow solution and the mixture was stirred at RT under Ar. After 2 days, the reaction was analyzed by TLC (1:1 hexanes/EtOAc - UV lamp). The mixture was concentrated and the yellow residue was partitioned between H<sub>2</sub>O (10 mL), and EtOAc (10 mL). The layers were separated and the aqueous layer was extracted with EtOAc (2 x 15 mL). The combined organic layers were washed with H<sub>2</sub>O (20 mL), saturated NaHCO<sub>3</sub> (2 x 20 mL), H<sub>2</sub>O (20 mL), 10 wt% citric acid (2 x 20 mL), H<sub>2</sub>O (20 mL), and brine (20 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. The crude material was chromatographed through 15 g of silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 60% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to yield **3.110** as a bright yellow oil (49.8 mg,

80.6% yield). Rf = 0.31 (3:2 CH<sub>2</sub>Cl<sub>2</sub>/EtOAc – UV lamp). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$ 7.76 (dd, J = 8.4, 1.8 Hz, 1H), 7.41 (ddd, J = 8.7, 7.2, 1.5 Hz, 1H), 7.03 (d, J = 8.4 Hz, 1H), 6.96 – 6.87 (m, 2H), 5.59 (m, 1H), 5.44 (ddd, J = 10.8, 7.2, 1.2 Hz, 1H), 5.05 (ddd, J= 11.1, 8.4, 1.5 Hz, 1H), 4.63 (m, 1H), 4.41 (t, J = 7.5 Hz, 1H), 3.64 (m, 1H), 2.82 (m, 1H), 2.58 (m, 1H), 2.03 – 1.85 (m, 6H), 1.62 – 1.25 (m, 6H) ppm. <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  171.4, 169.9, 142.0, 140.1, 133.6, 130.2, 130.1, 126.2, 119.4, 117.5, 72.0, 63.2, 53.1, 47.6, 35.8, 34.9, 31.6, 25.5, 23.8, 23.0, 21.2 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>21</sub>H<sub>28</sub>N<sub>3</sub>O<sub>5</sub><sup>+</sup>, 402.2029; obsd, 402.2003.



(1*S*,4*R*,*Z*)-4-((*S*)-1-(2-Nitrophenyl)pyrrolidine-2-carboxamido)cyclooct-2-enyl acetate (3.111). Compound 3.111 was prepared following the same procedure used for 3.110. Used the free amine intermediate (28.9 mg, 0.158 mmol), HOBt (25.0 mg, 0.185 mmol), 3.109 (46.0 mg, 0.195 mmol), and EDC•HCl (43 mg, 0.22 mmol) to provide 3.111 as a bright yellow oil (62.6 mg, 98.9% yield). Rf = 0.31 (3:2 CH<sub>2</sub>Cl<sub>2</sub>/EtOAc – UV lamp). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.76 (dd, *J* = 8.4, 2.1 Hz, 1H), 7.41 (ddd, *J* = 8.7, 7.2, 1.5 Hz, 1H), 7.03 (d, *J* = 8.4 Hz, 1H), 6.93 (td, *J* = 8.4, 1.2 Hz, 1H), 6.85 (d, *J* = 7.2 Hz, 1H), 5.66 – 5.27 (m, 2H), 5.33 (m, 1H), 4.62 (m, 1H), 4.41 (t, *J* = 7.5 Hz, 1H), 3.64 (m, 1H), 2.81 (m, 1H), 2.56 (m, 1H), 2.10 – 1.80 (m, 6H), 1.70 – 1.30 (m, 6H) ppm. <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  171.4, 170.0, 141.9, 140.0, 133.4, 130.4, 130.0, 126.1, 119.3,

117.3, 72.0, 63.4, 53.0, 47.6, 35.5, 34.9, 31.5, 25.6, 23.7, 22.9, 21.2 ppm. HRMS (FAB)  $m/z \,[M+H]^+ \,\text{calcd for } C_{21}H_{28}N_3O_5^+, 402.2029; \,\text{obsd}, 402.2009.$ 



(-)-*tert*-Butyl (1*R*,4*S*,*Z*)-4-(*tert*-butyldimethylsilyloxy)cyclooct-2-

enylcarbamate (1*R*,4*S*)-3.79. The title compound was prepared following the same procedure used for racemic 3.79. The use of (-)-3.76 (0.916 g, 3.80 mmol), imidazole (1.05 g, 15.5 mmol), and *tert*-butyldimethylsilyl chloride (1.03 g, 6.84 mmol) provided (-)-3.79 as a white solid (1.148 g, 85.1% yield). mp = 85.5-86.5 °C. NMR spectra matches racemic 3.79.  $[\alpha]^{23} = -12.5$  (c = 0.52 in MeOH).



(+)-*tert*-Butyl (1*S*,4*R*,*Z*)-4-(*tert*-butyldimethylsilyloxy)cyclooct-2-

enylcarbamate (1*S*,4*R*)-3.79. The title compound was prepared following the same procedure used for racemic 3.79. The use of (+)-3.76 (1.36 g, 5.65 mmol), imidazole (1.54 g, 22.6 mmol), and *tert*-butyldimethylsilyl chloride (1.54 g, 10.2 mmol) provided (+)-3.79 as a white solid (1.91 g, 95.1% yield). mp = 85.5-86.5 °C. NMR spectra matches racemic 3.79.  $[\alpha]^{23} = +17.1$  (c = 0.53 in MeOH).



*tert*-Butyl (2*R*,7*S*)-7-(*tert*-butyldimethylsilyloxy)-1,8-dihydroxyoctan-2ylcarbamate (2*R*,7*S*)-3.94. The title compound was prepared following the same procedure used for racemic 3.94. The use of (-)-3.79 (2.45 g, 6.89 mmol) and sodium borohydride (1.41 g, 37.3 mmol) provided (2*R*,7*S*)-3.94 as a light yellow sticky oil (2.71 g, 99% yield). NMR spectra matches racemic 3.94.  $[\alpha]^{23} \sim 0$  (c = 0.5 in MeOH). HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>19</sub>H<sub>42</sub>NO<sub>5</sub>Si<sup>+</sup>, 392.2832; obsd, 392.2812.



*tert*-Butyl (2*S*,7*R*)-7-(*tert*-butyldimethylsilyloxy)-1,8-dihydroxyoctan-2ylcarbamate (2*S*,7*R*)-3.94. The title compound was prepared following the same procedure used for racemic 3.94. The use of (+)-3.79 (1.84 g, 5.19 mmol) and sodium borohydride (1.01 g, 26.7 mmol) provided (2*S*,7*R*)-3.94 as a light yellow sticky oil (1.63 g, 80.2% yield). NMR spectra matches racemic 3.94.  $[\alpha]^{23} \sim 0$  (c = 0.5 in MeOH). HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>19</sub>H<sub>42</sub>NO<sub>5</sub>Si<sup>+</sup>, 392.2832; obsd, 392.2841.



*tert*-Butyl (2*R*,7*S*)-1,7,8-trihydroxyoctan-2-ylcarbamate (2*R*,7*S*)-3.96. The title compound was prepared following the same procedure used for racemic 3.96. The use of (2*R*,7*S*)-3.94 (2.56 g, 6.54 mmol) and 1.0M TBAF in THF (8.0 mL, 8.0 mmol) provided (2*R*,7*S*)-3.96 as a white solid (1.64 g, 90.3% yield). mp = 75-77 °C. NMR spectra matches racemic 3.96.  $[\alpha]^{23} = -1.1$  (c = 0.47 in MeOH).



*tert*-Butyl (2*S*,7*R*)-1,7,8-trihydroxyoctan-2-ylcarbamate (2*S*,7*R*)-3.96. The title compound was prepared following the same procedure used for racemic 3.96. The use of (2*S*,7*R*)-3.94 (1.55 g, 3.96 mmol) and 1.0M TBAF in THF (4.8 mL, 4.8 mmol) provided (2*S*,7*R*)-3.96 as a white solid (0.982 g, 89.3% yield). mp = 74-76 °C. NMR spectra matches racemic 3.96.  $[\alpha]^{23} = -5.2$  (c = 0.33 in MeOH).



*tert*-Butyl (*R*)-6-((*S*)-2,2-dimethyl-1,3-dioxolan-4-yl)-1-hydroxyhexan-2ylcarbamate (2*R*,7*S*)-3.98. The title compound was prepared following the same procedure used for racemic 3.98. The use of (2*R*,7*S*)-3.96 (1.52 g, 5.49 mmol), 2,2dimethoxypropane (0.70 mL, 5.7 mmol), and *p*-toluenesulfonic acid (45 mg, 0.26 mmol) provided (2*R*,7*S*)-3.98 as a yellow oil that solidified upon standing (1.64 g, 95.5% yield). NMR spectra matches racemic 3.98.  $[\alpha]^{23} \sim 0$  (c = 0.5 in MeOH). HRMS (FAB) *m/z*  $[M+H]^+$  calcd for C<sub>16</sub>H<sub>32</sub>NO<sub>5</sub><sup>+</sup>, 318.2280; obsd, 318.2259.



*tert*-Butyl (*S*)-6-((*R*)-2,2-dimethyl-1,3-dioxolan-4-yl)-1-hydroxyhexan-2ylcarbamate (2*S*,7*R*)-3.98. The title compound was prepared following the same procedure used for racemic 3.98. The use of (2*S*,7*R*)-3.96 (0.837 g, 3.02 mmol), 2,2dimethoxypropane (0.38 mL, 3.1 mmol), and *p*-toluenesulfonic acid (25 mg, 0.15 mmol) provided (2*S*,7*R*)-3.98 as a yellow oil that solidified upon standing (0.683 g, 71.4% yield). NMR spectra matches racemic 3.98.  $[\alpha]^{23} \sim 0$  (c = 0.5 in MeOH). HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>16</sub>H<sub>32</sub>NO<sub>5</sub><sup>+</sup>, 318.2280; obsd, 318.2268.



(-)-*tert*-Butyl (*R*)-6-((*S*)-2,2-dimethyl-1,3-dioxolan-4-yl)-1-oxohexan-2ylcarbamate (2*R*,7*S*)-3.100. The title compound was prepared following the same procedure used for racemic 3.100 by Swern oxidation. (2*R*,7*S*)-3.98 (0.560 g, 1.76 mmol), oxalyl chloride (0.23 mL, 2.7 mmol), DMSO (0.33 mL, 4.6 mmol), and triethylamine (1.20 mL, 8.54 mmol) provided (2*R*,7*S*)-3.100 as a yellow oil that solidified (0.447 g, 80.3% yield). mp = 42-45 °C. NMR spectra matches racemic 3.100.  $[\alpha]^{23} = -7.7$  (c = 0.59 in CHCl<sub>3</sub>). HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>16</sub>H<sub>30</sub>NO<sub>5</sub><sup>+</sup>, 316.2124; obsd, 316.2131.



(+)-*tert*-Butyl (S)-6-((R)-2,2-dimethyl-1,3-dioxolan-4-yl)-1-oxohexan-2ylcarbamate (2S,7R)-3.100. The title compound was prepared following the same procedure used for racemic 3.100 by Swern oxidation. (2S,7R)-3.98 (0.411 g, 1.30 mmol), oxalyl chloride (0.18 mL, 2.1 mmol), DMSO (0.33 mL, 4.6 mmol), and triethylamine (0.90 mL, 6.4 mmol) provided (2S,7R)-3.100 as a yellow oil that solidified (0.353 g, 86.3% yield). mp = 45-47 °C. NMR spectra matches racemic **3.100**.  $[\alpha]^{23}$  = +36.2 (c = 0.54 in CHCl<sub>3</sub>). HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>16</sub>H<sub>30</sub>NO<sub>5</sub><sup>+</sup>, 316.2124; obsd, 316.2131.



(R)-2-(tert-Butoxycarbonylamino)-6-((S)-2,2-dimethyl-1,3-dioxolan-4-

yl)hexanoic acid (2*R*,7*S*)-3.101. The title compound was prepared following the same procedure used for racemic 3.101 by KMnO<sub>4</sub> oxidation. (2*R*,7*S*)-3.100 (0.458 g, 1.45 mmol), KMnO<sub>4</sub> (0.506 g, 3.20 mmol), and benzyltriethylammonium chloride (17.4 mg, 0.0764 mmol) provided (2*R*,7*S*)-3.101 as a yellow oil (0.435 g, 90.5% yield). NMR spectra matches racemic 3.101.  $[\alpha]^{23} = -5.2$  (c = 0.35 in CHCl<sub>3</sub>). HRMS (FAB) *m/z* [M+H]<sup>+</sup> calcd for C<sub>16</sub>H<sub>30</sub>NO<sub>6</sub><sup>+</sup>, 332.2073; obsd, 332.2054.



(S)-2-(tert-Butoxycarbonylamino)-6-((R)-2,2-dimethyl-1,3-dioxolan-4-

yl)hexanoic acid (2S,7R)-3.101. The title compound was prepared following the same procedure used for racemic 3.101 by KMnO<sub>4</sub> oxidation. (2S,7R)-3.100 (0.372 g, 1.18

mmol), KMnO<sub>4</sub> (0.412 g, 2.61 mmol), and benzyltriethylammonium chloride (15.7 mg, 0.0689 mmol) provided (**2***S***,7***R***)-3.101** as a yellow oil (0.307 g, 78.6% yield). NMR spectra matches racemic **3.101**.  $[\alpha]^{23} = -2.1$  (c = 0.39 in CHCl<sub>3</sub>). HRMS (FAB) *m/z* [M+H]<sup>+</sup> calcd for C<sub>16</sub>H<sub>30</sub>NO<sub>6</sub><sup>+</sup>, 332.2073; obsd, 332.2048.



Procedure for diazotization of L-lysine. Preparation of intermediate carboxylic acids 3.113 and 3.114. A 3-necked 500-mL round-bottomed flask was charged with Cbz-L-Lys-OH (3.112, 10.34 g, 36.88 mmol) and 150 mL of H<sub>2</sub>O. The flask was fitted with a temperature probe and a pH probe and the solution was heated to 60 °C (internal temperature) in an oil bath. The pH of the solution was adjusted to 9.5 through the addition of 3M NaOH (or 6M HCl if too basic). Sodium nitroferricyanide dihydrate (13.20 g, 44.30 mmol) was added to the solution in portions while the solution was stirred vigorously at 60 °C. As the reaction became more acidic, the pH was maintained at 9.5 by the addition of 3M NaOH. Bubbles were observed and the color of the solution chanbed from colorless to yellow to a brick red suspension. After the addition was complete, the reaction was stirred vigorously while the internal temperature was maintained at 60 °C and the pH was maintained at 9.5 for 4 h. The mixture was cooled to RT, then filtered through a pad of celite. The filter cake was washed with H<sub>2</sub>O (200 mL) and the reddish-orange filtrate was extracted with EtOAc (3 x 200 mL). The combined organic layers were washed with brine (2 x 100 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a yellow oil (11.7 g crude material). The crude

material was used directly for the preparation of compounds **3.115-3.118** without purification.



(S)-Benzyl 1-(benzyloxyamino)-6-hydroxy-1-oxohexan-2-ylcarbamate (3.115). The crude mixture of **3.113** and **3.114** (36.9 mmol) was dissolved in 300 mL of THF and 225 mL of H<sub>2</sub>O in a 1-L beaker equipped with a stir bar and pH probe. 0-Benzylhydroxylamine hydrochloride (8.90 g, 55.8 mmol) was added and the pH of the yellow solution was adjusted to 4.5 by adding 3M NaOH (~10-15 mL). To the yellow solution was added EDC•HCl (30.1 g, 157 mmol) in portions over 80 min. The pH of the reaction was re-adjusted to 4.5 by adding 6M HCl as the pH of the solution increased. The mixture became cloudy after  $\sim$ 1-2 equivalents of EDC was added. After the addition of EDC•HCl was complete, the mixture was stirred for an additional 50 min (~2 h total reaction time) as the pH was maintained at 4.5 by adding 6M HCl. EtOAc (150 mL) was added and the layers were separated. The aqueous layer was extracted with EtOAc (3 x 150 mL), and the combined organic layers were washed with brine (2 x 100 mL), dried over Na<sub>2</sub>SO, filtered, and concentrated to yield a light yellow solid. The crude material was purified through 500 g of silica using 80% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to afford the alkene product, **3.116** (see below), then 100% EtOAc to afford the alcohol product, **3.115** as a fluffy white solid (6.14 g, 43.1% yield from **3.112**). mp = 115-117 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  9.89 (s, 1H), 7.31 (m, 10H), 5.76 (d, J = 8.5 Hz, 1H), 5.02 (d, J = 12.5Hz, 1H), 4.95 (d, J = 12.5 Hz, 1H), 4.87 (d, J = 11 Hz, 1H), 4.84 (d, J = 11 Hz, 1H), 4.04

(q, J = 7.5 Hz, 1H), 3.50 (m, 2H), 2.58 (br, 1H), 1.76 – 1.68 (m, 1H), 1.63 – 1.55 (m, 1H), 1.50 – 1.48 (m, 2H), 1.35 – 1.30 (m, 2H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  169.3, 156.3, 135.9, 135.0, 129.2, 128.7, 128.5, 128.4, 128.2, 127.9, 78.1, 67.1, 62.0, 52.3, 32.1, 31.7, 21.7 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>21</sub>H<sub>27</sub>N<sub>2</sub>O<sub>5</sub><sup>+</sup>, 387.1920; obsd, 387.1927.



(*S*)-2-(Benzyloxycarbonylamino)hex-5-enoic acid (3.116). Compound 3.116 was isolated as described above as a yellow oil that solidified upon standing (3.96 g, 29.2% yield from 3.112). Still contains impurities by NMR. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  9.43 (s, 1H), 7.36 – 7.30 (m, 10H), 5.73 (td, *J* = 16.8, 6.5 Hz, 1H), 5.56 (d, *J* = 8.5 Hz, 1H), 5.05 – 4.95 (m, 5H), 4.88 (s, 2H), 4.05 (q, *J* = 7.5 Hz, 1H), 2.10 – 1.95 (m, 2H), 1.89 – 1.81 (m, 1H), 1.73 – 1.65 (m, 1H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$ 169.1, 156.2, 136.8, 135.8, 135.0, 129.2, 128.7, 128.5, 128.2, 128.0, 127.9, 115.8, 78.2, 67.1, 51.8, 31.2, 29.4, 17.9 ppm. MS (FAB) *m/z* [M+H]<sup>+</sup> at 369 (base peak), 325, 246, 214.



(S)-Methyl 2-(benzyloxycarbonylamino)-6-hydroxyhexanoate (3.118). The crude mixture of 3.13 and 3.14 (12 mmol) was dissolved in about 100 mL of anhydrous

 $Et_2O$ . The blue-green solution was cooled in a crushed ice/H<sub>2</sub>O bath and a solution of excess CH<sub>2</sub>N<sub>2</sub> in ether was added dropwise to the mixture. The reaction was stirred in the ice/H<sub>2</sub>O bath for an additional 20 min then quenched by adding 10% acetic acid in Et<sub>2</sub>O (some bubbling was observed). The mixture was poured into 100 mL of saturated NaHCO<sub>3</sub> and the layers were separated. The aqueous layer was extracted with  $Et_2O$  (2 x 75 mL) and the combined organic layers were washed with brine (75 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to yield a light orange oil. TLC analysis of the crude mixture was performed with 1:1 hexanes/EtOAc and 2:1 CH2Cl2/EtOAc (UV lamp, CAM stain). The crude material was loaded onto silica and purified through a Biotage 40M column using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 50% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc and provided 3.117 (see below) and 3.118 as a cloudy oil (1.346 g, 38% yield from 3.112). <sup>1</sup>NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.35 – 7.30 (m, 5H), 5.47 (d, J = 8.0 Hz, 1H), 5.09 (s, 2H), 4.37 (td, J = 8.0, 5.5 Hz, 1H), 3.72 (s, 3H), 3.60 (t, J = 6.5 Hz, 2H), 2.00 (br, 1H), 1.84 (m, 1H), 1.67 (m, 1H), 1.54 (m, 2H), 1.41 (m, 2H) ppm.  $^{13}$ C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$ 173.0, 155.9, 136.1, 128.4, 128.1, 128.0, 66.9, 62.2, 53.7, 52.3, 32.3, 31.9, 21.5 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>15</sub>H<sub>22</sub>NO<sub>5</sub><sup>+</sup>, 296.1498; obsd, 296.1518.



(*S*)-Methyl 2-(benzyloxycarbonylamino)hex-5-enoate (3.117). Compound 3.117 was isolated as described above as a cloudy oil (0.665 g, 20% yield from 3.112). Contains impurities by NMR. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.36 – 7.32 (m, 5H), 5.78 (tdd, *J* = 13.0, 10.5, 6.5 Hz, 1H), 5.33 (d, *J* = 8.5 Hz, 1H), 5.01 (m, 2H), 4.40 (td, *J* = 8.5,

5.5 Hz, 1H), 3.75 (s, 3H), 3.68 (m, 1H), 2.17 – 2.05 (m, 2H), 1.98 – 1.91 (m, 1H), 1.80 – 1.69 (m, 1H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  172.9, 155.8, 136.7, 128.5, 128.2, 128.15, 128.07, 115.8, 67.0, 53.3, 52.3, 31.8, 29.3 ppm. HRMS (FAB) *m*/*z* [M+H]<sup>+</sup> calcd for C<sub>15</sub>H<sub>20</sub>NO<sub>4</sub><sup>+</sup>, 278.1392; obsd, 278.1411.



(S)-6-(Benzyloxyamino)-5-(benzyloxycarbonylamino)-6-oxohexyl

methanesulfonate (3.119). Compound 3.115 (2.376 g, 6.15 mmol) was dissolved in 45 mL of anhydrous pyridine in a flame-dried single-necked 200-mL round-bottomed flask under Ar. The solution was cooled in an ice/H<sub>2</sub>O bath and methanesulfonyl chloride (0.58 mL, 7.5 mmol) was added dropwise over 10 min. The solution was stirred for 1 h at 0 °C under Ar. The reaction was partitioned between EtOAc (200 mL) and 1M HCl (200 mL). The layers were separated and the organic layer was washed with 1M HCl (3 x 100 mL), H<sub>2</sub>O (100 mL), and brine (100 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to yield a white solid (2.714 g, 95.0% yield). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 9.19 (s, 1H), 7.35 – 7.32 (m, 10H), 5.45 (d, *J* = 8.4 Hz, 1H), 5.04 (d, *J* = 12 Hz, 1H), 4.99 (d, *J* = 12 Hz, 1H), 4.88 (s, 2H), 4.15 (m, 2H), 4.01 (q, *J* = 7.2 Hz, 1H), 2.96 (s, 3H), 1.80 – 1.55 (m, 4H), 1.45 – 1.35 (m, 2H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 169.0, 156.2, 135.8, 134.9, 129.2, 128.7, 128.49, 128.47, 128.2, 127.9, 78.2, 69.5, 67.1, 52.1, 37.2, 31.5, 28.4, 21.3 ppm.



(S)-Benzyl 1-(benzyloxy)-2-oxoazepan-3-ylcarbamate (3.120). Mesylate 3.119 (6.32 g, 13.6 mmol) was dissolved in 350 mL of reagent grade acetone in a single-necked 1-L round-bottomed flask. Potassium carbonate (5.80 g, 42.0 mmol) was added, the flask was fitted with a water condenser and heated to reflux (oil bath temperature = 85-95 °C) overnight. After 23 h, the reaction was cooled slowly to RT and allowed to stir for an additional 2 days before diluting with EtOAc (150 mL) and H<sub>2</sub>O (150 mL). The layers were separated and the aqueous layer was extracted with EtOAc (3 x 150 mL). The combined organic layers were washed with brine (2 x 75 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to yield a yellow oil. The oil was chromatographed through a Biotage 40M column using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 80% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to afford 3.121 (see below), 3.120 as a colorless oil, and a mixture of 3.121 and 3.120 (4.64 g total, 92.6% total yield). Compound **3.120** was recovered as a colorless oil that solidified upon standing (3.33 g, 67% yield total). mp = 62-64 °C. Rf = 0.36 (9:1 CH<sub>2</sub>Cl<sub>2</sub>/EtOAc – CAM stain). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 7.43-7.30 (m, 10H), 6.15 (d, J = 6.0 Hz, 1H), 5.11 (m, 2H), 4.98 (d, J = 10.5 Hz, 1H), 4.88 (d, J = 10.5 Hz, 1H),4.27 (ddd, J = 11.0, 6.5, 1.5 Hz, 1H), 3.58 (dd, J = 16.0, 11.5 Hz, 1H), 3.46 (dd, J = 16.0, 5.0 Hz, 1H), 2.03 (d, J = 12.0 Hz, 1H), 1.92-1.85 (m, 1H), 1.69-1.63 (m, 2H), 1.56-1.48 (m, 1H), 1.43-1.35 (m, 1H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 170.0, 155.4, 136.4, 135.0, 129.6, 128.8, 128.5, 128.4, 128.0, 127.9, 76.8, 66.6, 53.2, 52.6, 31.8, 27.5, 26.1 ppm.



(*S*)-Benzyl 2-(benzyloxyimino)oxepan-3-ylcarbamate (3.121). Compound 3.121 was isolated as a glassy solid (1.31 g, 26% yield) as described above. Rf = 0.54 (9:1 CH<sub>2</sub>Cl<sub>2</sub>/EtOAc - CAM stain). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.38-7.28 (m, 10H), 5.93 (d, *J* = 6 Hz, 1H), 5.10 (s, 2H), 5.00 (s, 2H), 4.45 (dd, *J* = 9.5, 7.0 Hz, 1H), 4.37 (dd, *J* = 12.5, 4.5 Hz, 1H), 4.10 (m, 1H), 2.21 (d, *J* = 11.5 Hz, 1H), 1.90 (m, 2H), 1.72-1.58 (m, 4H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  155.6, 155.2, 137.5, 136.3, 128.5, 128.4, 128.2, 128.12, 128.07, 127.8, 76.4, 70.2, 66.8, 50.7, 35.0, 29.6, 26.2 ppm.



(*S*)-1-(Benzyloxy)-2-oxoazepan-3-amonium bromide (3.122). Hydroxamate 3.120 (2.128 g, 5.776 mmol) was dissolved in 60 mL of anhydrous  $CH_2Cl_2$  in a flamedried 250-mL round-bottomed flask under Ar. 33% HBr in acetic acid (6.0 mL, 35 mmol) was added to the solution slowly and the Ar inlet needle was removed. The reaction was stirred at RT under Ar. White solid was observed after 2 min. After 3 h, no starting material was observed by TLC (4:1  $CH_2Cl_2/EtOAc - CAM$  stain, ninhydrin stain) and a ninhydrin-positive baseline spot was observed. The mixture was diluted with  $CH_2Cl_2$  (20 mL), and concentrated by rotary evaporation (40 °C, 21 mm Hg) to yield a light orange slurry. The slurry was concentrated from toluene (3 x 50 mL) to remove traces of acetic acid and water to afford a tan/orange solid. The solid was dried under vacuum (1.5 mm Hg) overnight to afford HBr salt **3.122** as an off-white solid (1.80 g, 99% yield). <sup>1</sup>H NMR (500 MHz, d<sup>6</sup>-DMSO)  $\delta$  8.25 (br, 3H), 7.47 – 7.38 (m, 5H), 4.93 (d, *J* = 10.0 Hz, 1H), 4.90 (d, *J* = 10.0 Hz, 1H), 4.24 (d, *J* = 5.0 Hz, 1H), 3.88 (dd, *J* = 16.0, 11.5 Hz, 1H), 3.60 (dd, *J* = 16.0, 5.0 Hz, 1H), 1.86 (m, 2H), 1.73 – 1.50 (m, 3H), 1.33 – 1.25 (m, 1H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  167.1, 135.0, 129.3, 128.7, 128.4, 75.7, 51.3, 51.2, 28.2, 26.1, 25.8 ppm.



(*R*)-*N*-((*S*)-1-(Benzyloxy)-2-oxoazepan-3-yl)-3-hydroxybutanamide (3.124). Anhydrous CH<sub>3</sub>CN (30 mL) was added to a 100-mL round-bottomed flask containing 3.122 (1.00 g, 3.17 mmol), (*R*)-3-hydroxybutyric acid (0.3906 g, 3.752 mmol), and HOBt (0.501 g, 3.71 mmol). To the mixture was added triethylamine (0.50 mL, 3.6 mmol), followed by EDC•HCl (0.761 g, 3.97 mmol). The mixture was stirred at RT under Ar. After 1 h, the solid material had fully dissolved. The reaction was stirred overnight at RT under Ar. The cloudy peach-colored solution was concentrated and the residue was partitioned between H<sub>2</sub>O (25 mL) and EtOAc (20 mL). The layers were separated and the aqueous layer was extracted with EtOAc (2 x 25 mL). The combined organic layers were washed with H<sub>2</sub>O (30 mL), saturated NaHCO<sub>3</sub> (2 x 30 mL), H<sub>2</sub>O (30 mL), and brine (30 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to yield a white solid. The solid was chromatographed through a 40S Biotage column using 100% EtOAc to yield **3.124** as a white solid (0.891 g, 87.7% yield). mp = 120-123 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.42-7.34 (m, 5H), 7.09 (d, *J* = 6.5 Hz, 1H), 4.97 (d, *J* = 10.5 Hz, 1H), 4.88 (d, *J* = 10.5 Hz, 1H), 4.48 (ddd, J = 11.0, 6.5, 1.5 Hz, 1H), 4.17 (m, 1H), 3.94 (m, 1H), 3.60 (dd, J = 16.0, 11.5 Hz, 1H), 3.47 (dd, J = 16.0, 5.0 Hz, 1H), 2.43 (dd, J = 15.5, 3.0 Hz, 1H), 2.32 (dd, J = 15.5, 9.0 Hz, 1H), 1.98 (d, J = 12.5 Hz, 1H), 1.91 (m, 1H), 1.71-1.65 (m, 2H), 1.49-1.30 (m, 2H), 1.21 (d, J = 6.0 Hz, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  171.5, 170.0, 134.9, 129.6, 128.9, 128.6, 76.8, 64.7, 52.7, 51.6, 43.6, 31.3, 27.5, 26.1, 22.6 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>17</sub>H<sub>25</sub>N<sub>2</sub>O<sub>4</sub><sup>+</sup>, 321.1814; obsd, 321.1810.



(*S*)-*N*-((*S*)-1-(Benzyloxy)-2-oxoazepan-3-yl)-3-hydroxybutanamide (3.125). Compound 3.125 was prepared following the same procedure used for 3.124. The use of 3.122 (1.20 g, 3.81 mmol), (*S*)-3-hydroxybutyric acid (0.479 g, 4.60 mmol), HOBt (0.617 g, 4.57 mmol), EDC•HCl (0.938 g, 4.89 mmol), and triethylamine (0.60 mL, 4.27 mmol) provided 3.125 as a white solid (0.863 g, 70.7% yield). Rf = 0.14 (100% EtOAc – CAM stain). mp = 135-137 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.41-7.40 (m, 2H), 7.37-7.35 (m, 3H), 6.99 (d, *J* = 6.5 Hz, 1H), 4.98 (d, *J* = 10.5 Hz, 1H), 4.89 (d, *J* = 10.5 Hz, 1H), 4.47 (ddd, *J* = 11.5, 7.0, 2.0 Hz, 1H), 4.18 (m, 1H), 3.61 (dd, *J* = 16.0, 11.5 Hz, 1H), 3.48 (dd, *J* = 16.0, 5.0 Hz, 1H), 2.40 (dd, *J* = 15.0, 3.0 Hz, 1H), 2.32 (dd, *J* = 15.0, 9.0 Hz, 1H), 1.98 (m, 1H), 1.92 (m, 1H), 1.71-1.65 (m, 2H), 1.54-1.35 (m, 2H), 1.23 (d, *J* = 6.5 Hz, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  171.8, 170.1, 134.9, 129.6, 128.9, 128.6, 76.8, 64.9, 52.7, 51.8, 44.1, 31.2, 27.5, 26.1, 22.7 ppm. HRMS (FAB) *m*/z [M+H]<sup>+</sup> calcd for C<sub>17</sub>H<sub>25</sub>N<sub>2</sub>O<sub>4</sub><sup>+</sup>, 321.1814; obsd, 321.1802. 8.3 Experimental procedures for chapter 4



**Dimethyl** 2-hydroxy-7-((*S*)-2-(2-hydroxyphenyl)-4,5-dihydrooxazole-4carboxamido)octanedioate (4.5a and 4.5b). Compound 3.69 (130.8 mg, 0.392 mmol) was dissolved in 5 mL of anhydrous  $Et_2O$  in a 25-mL round-bottomed flask. The solution was cooled in an ice/H<sub>2</sub>O bath and HCl gas (generated by adding conc. H<sub>2</sub>SO<sub>4</sub> dropwise to conc. HCl/NaCl then drying by bubbling through conc. H<sub>2</sub>SO<sub>4</sub>) was bubbled through the solution for 30 min. The solution was brownish in color, and no significant precipitate was observed. Bubbling was ceased and the reaction was warmed to RT and stirred for an additional 4 h. The reaction was concentrated by rotary evaporation to yield the amine•HCl salt as a brown semi-solid residue. The crude material was used immediately without purification.

Anhydrous CH<sub>3</sub>CN (5 mL) was added to the crude amine (105.8 mg, 0.452 mmol) and **2.62** (102.4 mg, 0.494 mmol) in a 25-mL round-bottomed flask under Ar. Distilled triethylamine (0.10 mL, 0.71 mmol) was added and most of the solid material dissolved. To the resultant yellow solution was added HOBt (67.4 mg, 0.499 mmol) and EDC•HCl (108.1 mg, 0.564 mmol). The reaction was stirred at RT under Ar. After 4 days, the reaction was concentrated by rotary evaporation (35°C, 21 mm Hg). The orange residue was partitioned between EtOAc (20 mL) and H<sub>2</sub>O (30 mL). The layers were separated and the aqueous layer was extracted with EtOAc (2 x 20 mL). The combined organic layers were washed with saturated aqueous NaHCO<sub>3</sub> (2 x 20 mL), H<sub>2</sub>O (1 x 20 mL), 5 wt% aqueous citric acid (2 x 20 mL), H<sub>2</sub>O (1 x 20 mL), and brine (2 x 20 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated by rotary evaporation (35°C, 21 mm Hg) to yield an orange oil (200 mg). The oil was purified through 20 g of silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 50% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to yield a mixture of **4.5a** and **4.5b** as a cloudy pink residue (59.7 mg, 31.3% yield) and **4.6** (see below). Recovered as a mixture of diastereomers by NMR: <sup>1</sup>H NMR (500MHz, CDCl<sub>3</sub>)  $\delta$  7.67 (m, 2H), 7.41 (m, 2H), 7.02 (d, *J* = 8.5 Hz, 2H), 6.91 - 6.87 (m, 2H), 4.98 - 4.93 (m, 2H), 4.70 - 4.55 (m, 6H), 4.17 (dd, *J* = 7.6, 4.0 Hz, 1H), 4.11 (dd, *J* = 7.7, 4.0 Hz, 1H), 3.77 (s, 3H, CH3), 3.75 (s, 3H), 3.72 (s, 3H), 3.69 (s, 3H), 1.93 - 1.24 (m, 16H) ppm. MS (FAB) *m/z* [M+H]<sup>+</sup> at 423; 363, 162.



**Diacylated byproduct (4.6).** Compound **4.6** was isolated as described above as a cloudy yellow oil (63.6 mg, 23.0% yield). Recovered as a mixture of diastereomers by NMR: <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 7.69 – 7.62 (m, 2H), 7.46 – 7.53 (m, 2H), 7.07 – 7.00 (m, 2H), 6.91 – 6.82 (m, 2H), 5.12 – 5.04 (m, 3H), 4.78 – 4.55 (m, 4H), 3.75 (s, 3H), 3.73 (s, 3H), 3.72 (s, 3H), 3.70 (s, 3H), 1.95 – 1.70 (m, 4H), 1.47 – 1.30 (m, 4H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 172.2, 172.0, 169.99, 169.92, 169.87, 169.8, 169.7, 168.2, 168.1, 168.0, 159.9, 159.8, 134.7, 134.2, 128.7, 128.4, 119.2, 118.8, 116.93, 116.90, 72.9,

69.9, 69.0, 67.0, 66.3, 52.5, 52.1, 31.9, 31.8, 31.54, 30.47, 29.6, 24.8, 24.6 ppm. MS (FAB) *m*/*z* [M+H]<sup>+</sup> at 612 (base peak), 307, 208.



**Dimethyl** 2-acetoxy-7-((*S*)-2-(2-hydroxyphenyl)-4,5-dihydrooxazole-4carboxamido)octanedioate (mixture of diastereomers 4.7a and 4.7b). Compound 3.86 (194 mg, 0.517 mmol) was dissolved in 10 mL of anhydrous  $Et_2O$  in a 50-mL round-bottomed flask. Anhydrous HCl gas (generated by adding conc. H<sub>2</sub>SO<sub>4</sub> dropwise to NaCl/HCl, then passed through conc. H<sub>2</sub>SO<sub>4</sub>) was bubbled through the solution. After 30 min, the solution appeared slightly cloudy. After 1.5 h, TLC of the solution (1:1 hexanes/EtOAc - CAM stain, ninhydrin stain) indicated no starting material remained and a new, ninhydrin positive spot appeared on the baseline. The solution was concentrated to afford a yellow oil (mass recovery was >100%). The oil was used immediately without purification.

Anhydrous CH<sub>3</sub>CN (5 mL) was added to the yellow oil in a 50-mL roundbottomed flask under Ar. Anhydrous triethylamine (0.10 mL, 0.712 mmol) was added and the solid dissolved. To the light orange/yellow solution was added **2.62** (114 mg, 0.550 mmol) and the color of the solution changed to deep orange/red. EDC•HCl (122 mg, 0.636 mmol) was added and the solution was stirred under Ar at RT overnight. The reaction was concentrated by rotary evaporation ( $35^{\circ}$ C, 21 mm Hg). The orange residue was partitioned between EtOAc (20 mL) and H<sub>2</sub>O (30 mL). The layers were separated and the aqueous layer was extracted with EtOAc (2 x 20 mL). The combined organic layers were washed with saturated aqueous NaHCO<sub>3</sub> (2 x 20 mL), H<sub>2</sub>O (1 x 20 mL), 5 wt% aqueous citric acid (2 x 20 mL), H<sub>2</sub>O (1 x 20 mL), and brine (2 x 20 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated by rotary evaporation ( $35^{\circ}$ C, 21 mm Hg) to yield a yellow oil. The oil was purified through 20 g of silica using CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to afford the mixture of diastereomers **4.7a** and **4.7b** as a yellow oil (75.8 mg, 31.6% yield). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 11.33 (br-s, 2H), 7.68 - 7.65 (m, 2H), 7.42 - 7.37 (m, 2H), 7.02 -7.00 (m, 2H), 6.90 - 6.81 (m, 4H), 4.99 - 4.90 (m, 4H), 4.68 - 4.53 (m, 6H), 3.75 (s, 3H),3.72 (s, 3H), 3.69 (s, 3H), 3.68 (s, 3H), 2.11 (s, 3H), 2.07 (s, 3H), 1.90 – 1.64 (m, 10H), 1.42 - 1.23 (m, 12H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  172.06, 171.97, 170.50, 170.46, 170.38, 170.33, 170.26, 167.99, 167.75, 159.70 (2), 134.23, 128.49, 128.46, 119.02, 116.89, 116.86, 109.90, 109.88, 71.88, 71.84, 69.44, 69.42, 67.94, 67.88, 52.43, 52.40, 52.17, 52.11, 51.91, 31.98, 31.92, 30.72, 30.64, 29.58, 24.82, 24.77, 24.57, 25.50, 20.53, 20.48 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd. for C<sub>22</sub>H<sub>29</sub>N<sub>2</sub>O<sub>8</sub><sup>+</sup>, 465.1873; obsd, 465.1855.



(±)-(2*S*,7*R*)-Dimethyl 2-acetoxy-7-benzamidooctanedioate (4.8). Compound 3.86 (138 mg, 0.366 mmol) was dissolved in 1 mL of  $CH_2Cl_2$  in a 10-mL round-bottomed flask and cooled in a crushed ice/H<sub>2</sub>O bath under Ar. TFA (0.20 mL, 2.7 mmol) was

added to the reaction and the resultant light pink solution was stirred in the crushed ice/ $H_2O$  bath for 60 min. TLC of the reaction (1:1 hexanes/EtOAc - CAM stain) indicates a complex mixture, but some amine present on the baseline. The reaction was warmed to RT and allowed to stir for an additional 60 min, then concentrated and used immediately without purification.

The crude material was dissolved in  $CH_2Cl_2$  (1.5 mL). Benzoyl chloride (0.090 mL, 0.77 mmol) and Et<sub>3</sub>N (0.75 mL, 5.3 mmol) were added and the reaction was stirred at RT overnight. The reaction was diluted with H<sub>2</sub>O (10 mL) and CH<sub>2</sub>Cl<sub>2</sub> (10 mL) and the layers were separated. The aqueous layer was washed with  $CH_2Cl_2$  (5 mL) and the combined  $CH_2Cl_2$  layers were washed with saturated NaHCO<sub>3</sub> (2 x 15 mL), H<sub>2</sub>O (15 mL), 10 wt% citric acid (2 x 15 mL), H<sub>2</sub>O (15 mL), and brine (15 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a yellow oil. The oil was chromatographed through 20 g of silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 85% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc and provided **4.8** as a yellow oil (117 mg, 84% yield). Evidence of epimerization is observed from the <sup>13</sup>C NMR spectrum. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.77 (d, J = 7.0 Hz, 2H), 7.46 (m, 1H), 7.39 (t, J = 7.5 Hz, 2H), 6.83 (d, J = 7.5 Hz, 1H), 4.93 (t, J = 6.0 Hz, 1H), 4.76 (td, J = 8.0, 6.0 Hz, 1H), 3.73 (s, 3H), 3.67 (s, 3H), 2.06 (s, 3H), 1.80 -1.74 (m, 4H), 1.43 – 1.34 (m, 4H) ppm.  $^{13}$ C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  172.9, 171.0, 170.5, 170.4, 166.9, 133.7, 131.6, 128.4, 126.9, 71.9, 60.2, 52.33, 52.27, 52.1, 32.1, 30.7, 24.8, 24.6, 20.9, 20.5 ppm. MS (FAB) m/z [M+H]<sup>+</sup> found at 380.

Attempted saponification of compound 4.8. Compound 4.8 (22 mg, 0.058 mmol) was dissolved in 0.6 mL of 2:1 THF/H<sub>2</sub>O in a 1-dram screw-cap vial. LiOH (6

mg, 0.25 mmol) was added and the mixture was stirred at RT. Monitored by TLC (1:1 hexanes/EtOAc and 100% EtOAc - UV lamp, CAM stain). After 45 min the reaction was nearly complete, and after 1 h, no **4.8** was observed by TLC. 1 mL of 1M HCl was added to the vial and the acidic mixture (pH ~2-3 - pH paper) was extracted with EtOAc (10 x 1 mL). The combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to yield a light yellow residue (18.9 mg, ~100% yield). Residue did not dissolve in CDCl<sub>3</sub>, but dissolved in EtOAc. MS data did not show evidence of the desired product. No identifiable peaks were found in the LC-MS traces of the aqueous layer.



(4S)-N-(6-(2,2-Dimethyl-1,3-dioxolan-4-yl)-1-hydroxyhexan-2-yl)-2-(2-

hydroxyphenyl)-4,5-dihydrooxazole-4-carboxamide (mixture of diastereomers 4.10a and 4.10b). Compound ( $\pm$ )-3.99 (74.1 mg, 0.211 mmol) was dissolved in 10 mL of methanol in a flame-dried single-necked 25-mL round-bottomed flask. The solution was purged with Ar and 10 wt% Pd/C (17.7 mg) was added. The solution was purged with Ar, then H<sub>2</sub>, and stirred under 1 atm of H<sub>2</sub> (balloon). After 4 h, TLC analysis of the reaction mixture (1:1 hexanes/EtOAc - CAM stain, ninhydrin stain) indicated the starting material was consumed and a new, ninhydrin/CAM stain positive spot was observed on the baseline. The reaction was purged with Ar, then filtered through a pad of celite. The

celite was rinsed with MeOH (5 mL) and the filtrate was concentrated to yield a colorless residue. The material was used immediately without purification.

The residue (41 mg, 0.20 mmol) and HOBt (25 mg, 0.19 mmol) were dissolved in 5 mL of dry acetonitrile in a single-necked 25-mL round-bottomed flask under Ar. EDC•HCl (43 mg, 0.22 mmol) was added and the peach-colored solution was stirred at RT under Ar overnight.  $H_2O$  (30 mL) and EtOAc (20 mL) were added to the solution and the layers were separated. The aqueous layer was extracted with EtOAc (3 x 15 mL) and the combined organic layers were washed successively with saturated NaHCO<sub>3</sub> (2 x 20 mL), H<sub>2</sub>O (20 mL), 10 wt% citric acid (2 x 20 mL), and brine (2 x 20 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a light yellow residue. The residue was chromatographed through 15 g of silica gel using a solvent gradient from 2:1 CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to 100% EtOAc to afford one of the diastereomers of 4.10 pure (14.0 mg, 19.7% yield), and mixtures of diastereomers 4.10a and 4.10b (46.6 mg, 65.8% yield). Total yield of 4.10 was 60.6 mg (85.5% yield). Analysis of the pure diastereomer isolated: mp = 112-115 °C. Rf = 0.15 (1:2 CH<sub>2</sub>Cl<sub>2</sub>/EtOAc – UV lamp; other isomer Rf =0.09). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 11.37 (s, 1H), 7.69 (m, 1H), 7.43 (m, 1H), 7.02 (d, J = 8.5 Hz, 1H), 6.91 (td, J = 8.0, 0.5 Hz, 1H), 6.48 (d, J = 8.0 Hz, 1H), 4.95 (dd, J = 10.0 Hz, 1H), 4.0 Hz, 1H), 4.0 Hz, 1H, 1 11.0, 8.0 Hz, 1H), 4.70 (dd, J = 11.0, 9.0 Hz, 1H), 4.62 (t, J = 8.5 Hz, 1H), 4.06 (m, 1H), 4.02 (dd, *J* = 7.5, 6.0 Hz, 1H), 3.95 (m, 1H), 3.65 (d, *J* = 10.5 Hz, 1H), 3.56 (dd, *J* = 11.0, 5.5 Hz, 1H), 3.50 (t, J = 7.5 Hz, 1H), 2.46 (br-s, 1H), 1.66-1.40 (m, 8H), 1.40 (s, 3H), 1.35 (s, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 171.4, 168.0, 159.6, 134.4, 128.6, 119.2, 110.0, 108.7, 75.9, 69.8, 69.4, 68.0, 65.2, 51.9, 33.4, 30.9, 29.7, 26.9, 26.1, 25.7, 25.6 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>21</sub>H<sub>31</sub>N<sub>2</sub>O<sub>6</sub><sup>+</sup>, 407.2182; obsd, 407.2176.



(*R*)-Methyl 6-amino-2-(benzyloxycarbonylamino)hexanoate (4.15). Methanol (10 mL) was cooled under Ar to 4 °C in an ice/H<sub>2</sub>O bath. Thionyl chloride (0.35 mL, 4.8 mmol) was added and the resultant solution was transferred via cannula to a stirred suspension of 4.14 (1.03 g, 3.67 mmol) in 30 mL of MeOH under Ar at 4 °C. The mixture was stirred for 1 h, then warmed to RT and stirred for an additional 7.5 h. The reaction was concentrated to yield a yellow oil. The oil was dissolved in saturated NaHCO<sub>3</sub> (100 mL) and extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 x 20 mL). The combined organic layers were dried over MgSO<sub>4</sub>, filtered, and concentrated to yield 4.15 as a reddish-brown oil (1.05 g, 97% yield). The oil was stored at -10 °C and used immediately, otherwise decomposition was observed.



## (R)-13,13-Dimethyl-3,11-dioxo-1-phenyl-2,12-dioxa-4,10-diazatetradecane-5-

**carboxylic acid (4.16).** Sodium bicarbonate (0.343 g, 4.08 mmol) was added to **4.14** (0.454 g, 1.62 mmol) in 12 mL of 2:1 THF/H<sub>2</sub>O, followed by di-*tert*-butyldicarbonate (0.411 g, 1.88 mmol). The mixture was stirred at RT overnight. The THF was removed *in vacuo* and the reaction was diluted with H<sub>2</sub>O (50 mL) and CH<sub>2</sub>Cl<sub>2</sub> (20 mL). The

aqueous layer was acidified to a pH of 2-3 (pH paper) using 1M HCl. The layers were separated and the aqueous was extracted with  $CH_2Cl_2$  (3 x 15 mL). The combined organic layers were washed with brine (2 x 20 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to a pale yellow oil. The oil was dried under vacuum to yield **4.16** as a white foam (0.577 g, 94% yield). The crude material was used directly without further purification.



2S-Benzyloxycarbonylamino-6-*tert*-butoxycarbonylamino-hexanoic acid benzyl ester (4.17). Crude 4.16 (651 mg, 1.71 mmol) was dissolved in 10 mL of CH<sub>3</sub>CN. K<sub>2</sub>CO<sub>3</sub> (548 mg, 3.96 mmol) was added, followed by benzyl bromide (0.30 mL, 2.53 mmol). The reaction was stirred for 24 h and monitored by TLC (3:1 hexanes/EtOAc – ninhydrin stain). The solid was removed by filtration and the filtrate was concentrated to yield a cloudy oil (1.20 g). Chromatographed through 40 g of silica using 3:1 hexanes/EtOAc to yield 4.17 as a colorless oil (747 mg, 93%). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.347 (s, 10H), 5.386 (d, *J* = 7.2 Hz, 1H), 5.167 (d, *J* = 6.0 Hz, 2H), 5.101 (s, 2H), 4.507 (br-s, 1H), 4.399 (m, 1H), 3.047 (m, 1H), 1.826 – 1.626 (m, 2H), 1.422 (s, 9H) ppm.


6-Benzyloxycarbonylamino-2*S*-*tert*-butoxycarbonylamino-hexanoic acid benzyl ester (4.20). Compound 4.20 was prepared following the procedure used for 4.17. The use of 4.19 (801 mg, 2.10 mmol), K<sub>2</sub>CO<sub>3</sub> (735 mg, 5.32 mmol), and BrBr (0.38 mL, 3.20 mmol) afforded a colorless oil (1.20 g). Chromatographed through 50 g of silica using 3:1 hexanes/EtOAc provided 4.20 as a light yellow oil (986 mg, 100%). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 7.36 – 7.34 (m, 10H), 5.20 (d, *J* = 12 Hz, 1H), 5.11 (d, *J* = 12 Hz, 1H), 5.09 (br-s, 4H), 4.77 (m, 1H), 4.33 (m, 1H), 3.14 (m, 2H), 1.80 (m, 1H), 1.65 (m, 1H), 1.42 (s, 9H), 1.45 – 1.29 (m, 4H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 172.6, 156.4, 155.4, 136.5, 135.3, 128.5, 128.44, 128.39, 128.3, 128.1, 128.0, 79.8, 66.9, 66.5, 53.2, 40.5, 32.2, 29.3, 28.2, 22.3 ppm.



(*R*)-Benzyl 6-amino-2-(benzyloxycarbonylamino)hexanoate (4.18). HCl gas, prepared by adding conc.  $H_2SO_4$  dropwise to NaCl/conc. HCl and dried by bubbling through conc.  $H_2SO_4$ , was bubbled through a solution of 4.17 (0.634 g, 1.35 mmol) in Et<sub>2</sub>O at 0 °C for 2.5 h. The reaction was monitored by TLC for the disappearance of 4.17. The solvent was removed to yield a pale yellow oil. The oil was dissolved in saturated NaHCO<sub>3</sub> (50 mL) and CH<sub>2</sub>Cl<sub>2</sub> (20 mL) was added. The layers were separated and the aqueous layer was extracted with  $CH_2Cl_2$  (2 x 20 mL). The combined organic layers were dried over MgSO<sub>4</sub>, filtered, and concentrated to yield **4.18** as a pale yellow oil (0.461 g, 92% yield). The material was used immediately without purification.



(S)-Benzyl 2-amino-6-(benzyloxycarbonylamino)hexanoate (4.21). Compound 4.21 was prepared following the same procedure used for 4.18. 4.20 (0.980 g, 2.08 mmol) provided 4.21 as a pale yellow oil (0.697 g, 90% yield). The material was used immediately without purification.



(S)-Methyl 2-(benzyloxycarbonylamino)-6-((S)-2-phenyl-4,5-dihydrooxazole-4-carboxamido)hexanoate (4.22). Compound 3.11 (135 mg, 0.708 mmol) and 4.15 (250 mg, 0.848 mmol) were dissolved in CH<sub>3</sub>CN (20 mL). EDC•HCl (166 mg, 0.865 mmol) was added and the reaction was stirred at 23°C for 15 h. The reaction was monitored by TLC (2:3 hexanes/EtOAc). The reaction was diluted with H<sub>2</sub>O (50 mL) and extracted with EtOAc (4 x 30 mL). The combined EtOAc layers were washed with saturated NaHCO<sub>3</sub> (3 x 20 mL), 5% citric acid (3 x 20 mL), and brine (2 x 30 mL), dried (MgSO4), filtered, and concentrated to yield a pale orange oil (170 mg). Chromatographed through 17 g of silica using 3:2 CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to yield **4.22** as an orange oil (135 mg, 41%). Rf = 0.111 (2:3 hexanes/EtOAc). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.96 (dd, *J* = 8.1, 0.9 Hz, 2H), 7.52 (t, *J* = 7.3 Hz, 1H), 7.42 (t, *J* = 7.5 Hz, 2H), 7.34 (m, 5H), 6.72 (br-t, 1H), 5.44 (d, *J* = 8.1 Hz, 1H, NHCbz), 5.09 (s, 2H), 4.821 (dd, *J* = 10.8, 8.7 Hz, 1H), 4.68 – 4.54 (m, 2H), 4.34 (m, 1H), 3.71 (s, 3H), 3.34 – 3.15 (m, 2H), 1.88 – 1.60 (m, 2H), 1.53 – 1.33 (m, 4H) ppm. HRMS (FAB) *m*/*z* [M+H]<sup>+</sup> calcd for C<sub>25</sub>H<sub>30</sub>N<sub>3</sub>O<sub>6</sub><sup>+</sup>, 468.2135; obsd, 468.2144.



(S)-Methyl 2-(benzyloxycarbonylamino)-6-((S)-2-(2-hydroxyphenyl)-4,5dihydrooxazole-4-carboxamido)hexanoate (4.23). Compound 4.23 was prepared following the same procedure used for 4.22. The use of oxazoline 2.62 (0.104 g, 0.502 mmol), amine 4.15 (0.174 g, 0.591 mmol), and EDC•HCl (0.125 g, 0.653 mmol) provided a pale orange oil (0.149 g, 61%). Chromatography through 20 g of silica using 4:1 CH<sub>2</sub>Cl<sub>2</sub>/EtOAc afforded 4.23 as a white solid (0.127 g, 52%). mp = 142-143 °C. Rf = 0.189 (2:3 hexanes/EtOAc). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.68 (dd, *J* = 7.8, 1.3 Hz, 1H), 7.42 (td, *J* = 8.7, 1.5 Hz, 1H), 7.35 (m, 5H), 7.02 (d, *J* = 8.4 Hz, 1H), 6.91 (t, *J* = 7.8 Hz, 1H), 6.42 (br-t, 1H, NHBoc), 5.41 (d, *J* = 8.1 Hz, 1H, NHCbz), 5.10 (s, 2H), 4.90 (t, *J* = 9.5 Hz, 1H), 4.67 – 4.58 (m, 2H), 4.34 (m, 1H), 3.72 (s, 3H), 3.27 (m, 2H), 1.84 – 1.60 (m, 2H), 1.56 – 1.25 (m, 4H) ppm. <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  170.7, 167.9, 159.6, 155.9, 136.2, 134.3, 128.50, 128.46, 128.1, 119.2, 116.8, 110.0, 69.9, 67.9, 67.0, 53.6, 52.3, 38.8, 31.9, 28.9, 22.3 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for  $C_{25}H_{30}N_3O_7^{+}$ , 484.2084; obsd, 484.2107.



(*S*)-Benzyl 2-(benzyloxycarbonylamino)-6-((*S*)-2-phenyl-4,5-dihydrooxazole-4-carboxamido)hexanoate (4.24). Compound 4.24 was prepared following the same procedure used for 4.22. The use of acid 3.11 (89 mg, 0.47 mmol), amine 4.18 (204 mg, 0.550 mmol), and EDC•HCl (112 mg, 0.583 mmol) provided a light orange oil (156 mg). Chromatography through 16 g of silica using a solvent gradient from 3:1 to 3:2 CH<sub>2</sub>Cl<sub>2</sub>/EtOAc yielded 4.24 as a white solid (89 mg, 35%). mp = 125-128 °C. Rf = 0.167 (2:3 hexanes/EtOAc). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.97 (d, *J* = 7.2 Hz, 2H), 7.52 (t, *J* = 7.3 Hz, 1H), 7.42 (t, *J* = 7.5 Hz, 1H), 7.34 (br-s, 10H), 6.72 (br-t, *J* = 5.1 Hz, 1H), 5.48 (d, *J* = 6.9 Hz, 1H, NHCbz), 5.15 (d, *J* = 2.7 Hz, 2H), 5.09 (s, 2H), 4.814 (dd, *J* = 10.8, 9.0 Hz, 1H), 4.68 – 4.55 (m, 2H), 4.42 – 4.35 (m, 1H), 3.28 – 3.15 (m, 2H), 1.83 (m, 1H), 1.71 (m, 1H), 1.49 (m, 2H), 1.33 (m, 2H) ppm. HRMS (FAB) *m*/z [M+H]<sup>+</sup> calcd for C<sub>31</sub>H<sub>34</sub>N<sub>3</sub>O<sub>6</sub><sup>+</sup>, 544.2448; obsd, 544.2452.



(*S*)-Benzyl 2-(benzyloxycarbonylamino)-6-((*S*)-2-(2-hydroxyphenyl)-4,5dihydrooxazole-4-carboxamido)hexanoate (4.25). Compound 4.25 was prepared following the same procedure used for 4.22. The use of acid 2.62 (109 mg, 0.524 mmol), amine 4.18 (234 mg, 0.631 mmol), and EDC•HCl (127 mg, (0.663 mmol) provided a light orange oil (148 mg). Chromatography through 15 g of silica using 1:1 hexanes/EtOAc yielded 4.25 as a white solid (106 mg, 36%). mp = 149-151 °C. Rf = 0.270 (2:3 hexanes/EtOAc). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.68 (dd, *J* = 7.8, 1.3 Hz, 1H), 7.42 (m, 1H), 7.34 (m, 10H), 7.02 (d, *J* = 8.4 Hz, 1H), 6.91 (t, *J* = 7.8 Hz, 1H), 6.37 (br-t, 1H), 5.41 (d, *J* = 8.1 Hz, 1H, NHCbz), 5.16 (d, *J* = 3.3 Hz, 2H), 5.10 (s, 2H), 4.89 (t, *J* = 9.6 Hz, 1H), 4.67 – 4.60 (m, 2H), 4.42 – 4.35 (m, 1H), 3.26 – 3.18 (m, 2H), 1.83 (m, 1H), 1.70 (m, 1H), 1.51 (m, 2H), 1.31 (m, 2H) ppm. HRMS (FAB) *m*/z [M+H]<sup>+</sup> calcd for C<sub>31</sub>H<sub>34</sub>N<sub>3</sub>O<sub>7</sub><sup>+</sup>, 560.2397; obsd, 560.2406.



(*S*)-Methyl 2-(benzyloxycarbonylamino)-6-((*S*)-3-hydroxy-2-(2hydroxybenzamido)propanamido)hexanoate (4.26). Compound 4.26 was prepared following the same procedure used for 4.22. The use of acid 2.62 (90 mg, 0.399 mmol), amine **4.15** (123 mg, 0.418 mmol), and EDC•HCl (95 mg, 0.498 mmol) provided a pale yellow oil (217 mg). Chromatography through 25 g of silica using 1:4 hexanes/EtOAc yielded **4.26** as a white glassy solid (158 mg, 79%). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.99 (m, 1H), 7.56 (d, *J* = 7.5 Hz, 1H), 7.32 – 7.26 (m, 5H), 7.09 (t, *J* = 5.1 Hz, 1H), 6.89 (d, *J* = 8.1 Hz, 1H), 6.76 (t, *J* = 7.5 Hz, 1H), 5.77 (d, *J* = 7.8 Hz, 1H), 5.02 (s, 2H), 4.61 (m, 1H), 4.27 (m, 1H), 4.03 (m, 1H), 3.74 (q, *J* = 5.4 Hz, 1H), 3.65 (s, 3H), 3.19 (m, 2H), 1.76 – 1.30 (m, 6H) ppm. <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  172.7, 170.6, 170.2, 161.1, 156.1, 136.3, 134.4, 128.4, 128.1, 127.9, 126.6, 119.0, 118.3, 114.3, 67.1, 62.6, 54.4, 52.1, 39.1, 32.0, 28.7, 22.4 ppm.



(*S*)-Benzyl 6-(benzyloxycarbonylamino)-2-((*S*)-2-phenyl-4,5-dihydrooxazole-4-carboxamido)hexanoate (4.27). Compound 4.27 was prepared following the same procedure used for 4.22. The use of acid 3.11 (106 mg, 0.529 mmol), amine 4.21 (263 mg, 0.710 mmol), and EDC•HCl (143 mg, 0.748 mmol) provided a light orange oil (228 mg). Chromatography through 25 g of silica using a solvent gradient from 85:15 to 4:1 CH<sub>2</sub>Cl<sub>2</sub>/EtOAc yielded 4.27 as a white solid (195 mg, 65%). mp = 102-104 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.99 (dd, *J* = 8.4, 1.5 Hz, 2H), 7.51 (tt, *J* = 7.3, 1.5 Hz, 1H), 7.42 (t, *J* = 7.3 Hz, 1H), 7.35 (m, 10H), 7.23 (d, *J* = 7.8 Hz, 1H), 5.21 (d, *J* = 12 Hz, 1H), 5.18 (d, *J* = 12 Hz, 1H), 5.06 (s, 2H), 4.86 (m, 2H), 4.61 (m, 3H), 3.06 (m, 2H), 1.83 (m, 1H), 1.70 (m, 1H), 1.43 (m, 2H), 1.27 (m, 2H) ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for  $C_{31}H_{34}N_3O_6^+$ , 544.2448; obsd, 544.2445.



(S)-Benzyl 6-(benzyloxycarbonylamino)-2-((S)-2-(2-hydroxyphenyl)-4,5dihydrooxazole-4-carboxamido)hexanoate (4.28). Compound 4.28 was prepared following the same procedure used for 4.22. The use of acid 2.62 (56.9 mg, 0.275 mmol), amine 4.21 (95.8 mg, 0.259 mmol), and EDC•HCl (62 mg, 0.32 mmol) provided a cloudy oil (150 mg). Chromatography through 15 g of silica using 90% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc yielded **4.28** as an oil (111.5 mg, 77.1%). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  11.44 (br, 1H), 7.68 (d, J = 7.5 Hz, 1H), 7.41 – 7.29 (m, 10H), 7.02 (d, J = 8.5 Hz, 1H), 6.97 (d, J = 7.5Hz, 1H), 6.90 (t, J = 8 Hz, 1H), 5.22 (d, J = 12.5 Hz, 1H), 5.16 (d, J = 12.5 Hz, 1H), 5.06 (s, 2H), 4.92 (m, 2H), 4.62 (m, 3H), 3.08 (q, J = 6.5 Hz, 2H), 1.86 – 1.83 (m, 1H), 1.75 – 1.67 (m, 1H), 1.48 - 1.39 (m, 2H), 1.27 - 1.21 (m, 2H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) § 171.5, 170.3, 167.7, 159.6, 156.3, 136.5, 135.1, 134.2, 128.6, 128.5, 128.43, 128.36, 128.26, 128.0, 127.9, 119.0, 116.9, 109.9, 69.3, 67.9, 67.2, 66.4, 51.9, 40.4, 31.7, 29.6, 29.1, 22.2 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>31</sub>H<sub>34</sub>N<sub>3</sub>O<sub>7</sub><sup>+</sup>, 560.2397; obsd, 560.2375.



(*S*)-Benzyl 6-(benzyloxycarbonylamino)-2-((*R*)-2-phenyl-4,5-dihydrothiazole-4-carboxamido)hexanoate (4.31). Compound 4.31 was prepared following the same procedure used for 4.22. The use of thiazoline 4.29 (53.3 mg, 0.257 mmol), amine 4.21 (96.5 mg, 0.261 mmol), and EDC+HCl (60.6 mg, 0.316 mmol) provided a yellow oil (159 mg). Chromatography through 15 g of silica using 90% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc yielded 4.31 as a yellow oil (109 mg, 75%). Rf = 0.24 (1:1 hexanes/EtOAc – UV lamp). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.88 (d, *J* = 7.5 Hz, 2H), 7.50 (t, *J* = 7.0 Hz, 1H), 7.44 – 7.33 (m, 12H), 5.21 – 5.16 (m, 4H), 5.10 – 5.05 (m, 2H), 4.84 (m, 1H), 4.69 (td, *J* = 7.5, 5.5 Hz, 1H), 3.66 (m, 2H), 3.06 (m, 2H), 1.91 – 1.83 (m, 1H), 1.71 – 1.66 (m, 1H), 1.50 – 1.41 (m, 2H), 1.30 – 1.25 (m, 2H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  171.6, 171.3, 171.0, 156.2, 136.5, 135.1, 132.3, 131.9, 128.6, 128.5, 128.44, 128.41, 128.39, 128.2, 128.1, 128.03, 127.97, 78.8, 67.1, 66.4, 51.8, 40.5, 35.5, 32.1, 28.9, 22.2 ppm. HRMS (FAB) *m/z* [M+H]<sup>+</sup> calcd for C<sub>31</sub>H<sub>34</sub>N<sub>3</sub>O<sub>5</sub>S<sup>+</sup>, 560.2219; obsd, 560.2228.



(S)-Benzyl 6-(benzyloxycarbonylamino)-2-((R)-2-(2-hydroxyphenyl)-4,5dihydrothiazole-4-carboxamido)hexanoate (4.32). Compound 4.32 was prepared following the same procedure used for **4.22**. The use of thiazoline **4.30** (57.5 mg, 0.258 mmol), amine **4.21** (86.7 mg, 0.234 mmol), and EDC•HCl (56.5 mg, 0.295 mmol) provided a yellow oil. Chromatography through 10 g of silica using 90% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc yielded **4.32** as an off-white solid (86.2 mg, 64%). Minor epimerization was observed from analysis of <sup>1</sup>H and <sup>13</sup>C spectra. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  11.95 (br, 1H), 7.45 (d, *J* = 7.5 Hz, 1H), 7.35 – 7.29 (m, 10H), 7.03 (d, *J* = 8.5 Hz, 1H), 6.89 (t, *J* = 7.5 Hz, 2H), 5.28 – 5.06 (m, 6H), 4.88 (br-m, 1H), 4.66 (td, *J* = 8.0, 5.5 Hz, 1H), 3.69 (dd, *J* = 11, 8.0 Hz, 1H), 3.60 (m, 1H), 3.07 (m, 2H), 1.87 (m, 1H), 1.70 (m, 1H), 1.44 (m, 2H), 1.27 (m, 2H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  175.8, 171.4, 169.8, 158.8, 156.3, 136.6, 135.1, 133.8, 130.8, 128.6, 128.51, 128.49, 128.4, 128.3, 128.2, 128.03, 127.98, 119.3, 117.3, 115.8, 77.9, 67.3, 66.5, 52.1, 40.5, 33.9, 31.8, 29.0, 22.2 ppm. HRMS (FAB) *m*/*z* [M+H]<sup>+</sup> calcd for C<sub>31</sub>H<sub>34</sub>N<sub>3</sub>O<sub>6</sub>S<sup>+</sup>, 576.2168; obsd, 576.2143.



3-Oxa-2-aza-bicyclo[2.2.1]hept-5-en-2-yl((S)-2-(2-hydroxyphenyl)-4,5dihydrooxazol-4-yl)methanone (mixture of diastereomers 4.36 and 4.36'). Hydroxamic acid 3.32 (0.304 g, 1.37 mmol) was dissolved in 14 mL of MeOH in a 100mL round-bottomed flask with a stir bar. The flask and stir bar were previously washed with 6M HCl, H<sub>2</sub>O, and acetone to remove any trace metals in the glassware. The methanolic solution was cooled in an ice/H<sub>2</sub>O bath to 4 °C (internal temp.). Freshly cracked cyclopentadiene (0.60 mL, 7.2 mmol) was added to the reaction followed immediately by the dropwise addition of a solution of NaIO<sub>4</sub> (0.315 g, 1.47 mmol) in 4

mL of H<sub>2</sub>O (0.36M). After 1 min, a white solid was observed in the reaction. The addition of the NaIO<sub>4</sub> solution was complete after 8 min. The reaction was stirred in the ice/H<sub>2</sub>O bath for an additional 40 min, after which time the reaction was complete as observed by TLC (1:1 hexanes/EtOAc and 9:1 CH<sub>2</sub>Cl<sub>2</sub>/MeOH - UV lamp). The white solid was removed by vacuum filtration and washed with EtOAc (50 mL). Brine (50 mL) was added to the filtrate, and solid immediately crashed out of solution.  $H_2O$  was added until all of the solid dissolved (30 mL). The layers were separated and the aqueous layer was extracted with EtOAc (2 x 50 mL). The combined EtOAc layers were washed with brine (50 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated by rotary evaporation (bath temp  $<40^{\circ}$ C) to yield a yellow oil (497 mg). The oil was chromatographed through a Biotage 40S column using a solvent gradient from 2:1 hexanes/EtOAc to 3:2 hexanes/EtOAc to yield the two isomers **4.36** (164.5 mg, 42%) and **4.36**' (114.8 mg, 29.3%), as well as mixed 4.36 and 4.36' (71.4 mg, 18.2%), all as colorless oils. Total combined yield of 4.36 and 4.36' was 350.7 mg (90%). Absolute configuration of the stereochemistry was not assigned. **4.36:** Rf = 0.21 (1:1 hexanes/EtOAc – UV lamp, CAM stain). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  11.76 (br, 1H), 7.59 (d, J = 7.5 Hz, 1H), 7.33 (t, J = 7.5 Hz, 1H), 6.94 (d, J = 8.5 Hz, 1H), 6.81 (m, 1H), 6.56 (m, 1H), 6.41 (ddd, J = 5.5, 2.5, 1.5 Hz, 1H), 5.32 (br-m, 2H), 5.00 (m, 1H), 4.68 (t, J = 8.0 Hz, 1H), 4.46 (dd, J = 10.0, 9.0 Hz, 1H), 1.97 (d, J = 8.0 Hz, 1H), 1.84 (d, J = 8.5 Hz, 1H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 171.5, 166.9, 159.7, 136.5, 133.5, 132.5, 118.4, 116.5, 110.1, 84.8, 68.2, 66.5, 62.3, 48.1 ppm. **4.36':** Isolated with impurity (imp) as described herein. Rf = 0.14 (1:1 hexanes/EtOAc – UV lamp, CAM stain). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  11.78 (br, 1H), 7.59 (d, *J* = 7.5 Hz, 1H), 7.33 (t, *J* = 7.5 Hz, 1H), 6.96 (dd, *J* = 8.0, 0.5 Hz, 1H),

6.82 (m, 1H), 6.59 (td, J = 5.5, 2.0 Hz, 1H), 6.36 (m, 1H), 5.34 (m, 2H), 5.07 (dd, J = 9.5, 8.5 Hz, 1H), 4.42 (m, 2H), 2.02 (d, J = 8.5 Hz, 1H), 1.85 (d, J = 9.0 Hz, 1H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  171.1, 167.1, 164.8 (imp), 159.7, 159.4 (imp), 136.8, 134.0 (imp), 133.9 (imp), 133.5, 133.1, 132.7 (imp), 128.4 (imp), 128.0, 118.9 (imp), 118.4, 116.6, 110.1, 84.7, 83.8 (imp), 68.1 (imp), 67.4, 66.5, 66.4 (imp), 64.6 (imp), 62.0, 48.4, 48.2 (imp).



3-Oxa-2-aza-bicyclo[2.2.1]hept-5-en-2-yl((S)-2-(2-(benzyloxy)phenyl)-4,5dihydrooxazol-4-yl)methanone (mixture of diastereomers 4.37 and 4.37'). The mixture of diastereomers was prepared following the same procedure for 4.36 and 4.36'. The use of hydroxamate 3.21 (52 mg, 0.166 mmol), cyclopentadiene (0.100 mL, 1.25 mmol), and NaIO<sub>4</sub> (60 mg, 0.282 mmol) provided a yellow oil. Chromatography through silica gel using 4:1 EtOAc/hexanes yielded a colorless oil composed of both diastereomers 4.36 and 4.36' as well as a mixture of other impurities (37 mg).



( $\pm$ )-(3*R*,5*S*)-3,5-Bis(methoxycarbonyl)isoxazolidin-2-ium chloride (4.38). Compound 3.44 (0.153 mg , 0.529 mmol) was dissolved in Et<sub>2</sub>O (25 mL) and HCl gas, produced from the addition of concentrated H<sub>2</sub>SO<sub>4</sub> to a mixture of concentrated HCl and

NaCl, was bubbled through the solution for 1.5 h. White solid was observed and the reaction was monitored by TLC. The solvent was evaporated and the residue was used immediately without purification.



(±)-(3*R*,6*S*)-Dimethyl morpholine-3,6-dicarboxylate (4.41). TFA (0.50 mL, 6.5 mmol) was added slowly to a solution containing 3.45 (0.246 g, 0.811 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) at 4 °C (ice/H<sub>2</sub>O bath). The mixture was stirred for 10 min, then warmed to RT and stirred for an additional 3 g. 3.45 was still present by TLC, so an additional 4 eq. of TFA was added to the reaction. After stirring for 1.5 h at RT, 25 mL of 10 wt% aqueous Na<sub>2</sub>CO<sub>3</sub> was added to the solution slowly and the layers were separated. The aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 x 20 mL) and the combined organic layers were dried over MgSO<sub>4</sub>, filtered, and concentrated to yield 4.41 as a yellow oil (0.141 g, 86%). The oil was used immediately without purification. Rf = 0.16 (1:1 hexanes/EtOAc – CAM stain). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  4.41 (dd, *J* = 7.8, 4.2 Hz, 1H), 3.76 (s, 3H), 3.75 (s, 3H), 3.71 (dd, *J* = 9.0, 4.2 Hz, 1H), 2.21 – 1.82 (m, 4H) ppm.



(±)-(3*R*,6*S*)-Dimethyl morpholine-3,6-dicarboxylate (4.42). Compound 4.42 was prepared following the same procedure for 4.41. Compound 3.48 (0.623 g, 1.37

mmol) provided **4.42** as a yellow oil (0.443 g, 91%). The oil was used directly without purification. Rf = 0.38 (1:1 hexanes/EtOAc – CAM stain). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.35 (m, 10H), 6.50 (d, *J* = 2.7 Hz, 1H), 5.30 – 5.14 (m, 4H), 4.45 (dd, *J* = 6.9, 4.8 Hz, 1H), 3.75 (dt, *J* = 5.4, 4.2 Hz, 1H), 2.26 – 1.90 (m, 4H) ppm.



(S)-4-((S)-1-(Benzyloxy)-2-oxoazepan-3-ylamino)-4-oxobutan-2-yl 2-(*tert*butoxycarbonylamino)-6-(2,2-dimethyl-1,3-dioxolan-4-yl)hexanoate (mixture of diastereomers 4.47 and 4.48). Preparation using Yamaguchi esterification conditions. Acid (±)-3.101 (52.9 mg, 0.160 mmol) and 3.125 (52.4 mg, 0.164 mmol) were dissolved in 3 mL of anhydrous CH<sub>2</sub>Cl<sub>2</sub> in a 25-mL round-bottomed flask under Ar. DMAP (5.8 mg, 0.047 mmol) was added, followed by triethylamine (0.045 mL, 0.32 mmol) and 2,4,6-trichlorobenzoyl chloride (0.050 mL, 0.32 mmol). The solution was stirred at RT under Ar. After about 1 day, the reaction was progressing by TLC (100% EtOAc - UV lamp, CAM stain), but also incomplete. The reaction was allowed to stir at RT under Ar. After 3 days, an additional 3 mL of CH<sub>2</sub>Cl<sub>2</sub> was added to replace the amount that had evaporated. After 5 days, not much had changed by TLC and the reaction was diluted with EtOAc (15 mL) and  $H_2O$  (10 mL) was added. The layers were separated and the aqueous layer was extracted with EtOAc (2 x 15 mL). The combined organic layers were washed with saturated NaHCO<sub>3</sub> (20 mL) and brine (20 mL), dried

over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to yield a colorless residue. The residue was chromatographed through 12 g of silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 100% EtOAc to afford the mixture of diastereomers **4.47** and **4.48** as a colorless residue (54.7 mg, 54.1% yield) and unreacted cobactin **3.125** as a white solid (4.1 mg, 7.8% recovery). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.44 – 7.41 (m, 2H), 7.40 – 7.37 (m, 3H), 7.08 (m, 1H), 5.36 – 5.19 (m, 2H), 5.00 (dd, *J* = 10.5, 3.0 Hz, 1H), 4.90 (dd, *J* = 10.5, 3.0 Hz, 1H), 4.48 (m, 1H), 4.25 (m, 1H), 4.07 – 3.89 (m, 2H), 3.68 – 3.61 (m, 1H), 3.53 – 3.46 (m, 2H), 2.60 – 2.55 (m, 1H), 2.50 – 2.44 (m, 1H), 2.03 – 1.96 (m, 1H), 1.94 – 1.88 (m, 1H), 1.86 – 1.78 (m, 1H), 1.76 – 1.58 (m, 4H), 1.50 – 1.30 (m, 19H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  171.9, 171.8, 170.0, 169.9, 168.2, 168.1, 155.4, 155.3, 134.90, 134.87, 129.5, 128.9, 128.8, 128.52, 128.50, 108.5, 79.6, 79.5, 76.8, 75.79, 75.77, 69.3, 68.9, 53.5, 52.64, 52.58, 51.73, 51.65, 42.6, 42.5, 33.29, 33.27, 33.2, 32.3, 32.2, 31.4, 28.3, 28.2, 27.5, 26.8, 26.1, 25.6, 25.42, 25.37, 25.3, 25.24, 25.22, 19.7, 19.6 ppm. HRMS (FAB) *m*/z [M]<sup>+</sup> calcd for C<sub>33</sub>H<sub>51</sub>N<sub>3</sub>O<sub>9</sub><sup>+</sup>, 633.3625; obsd, 633.3602.

Preparation of the mixture of diastereomers 4.47 and 4.48 using EDC and DMAP (Table 4.2, entry 5). Acid 3.101 (0.472 g, 1.43 mmol) and 3.125 (0.457 g, 1.43 mmol) were dissolved in 10 mL of anhydrous  $CH_2Cl_2$  in a 100-mL round-bottomed flask under Ar. DMAP (180 mg, 1.48 mmol) was added, followed by EDC•HCl (1.28 g, 6.68 mmol). The mixture was stirred at RT under Ar and monitored by TLC (100% EtOAc - UV lamp, CAM stain). The volume of the reaction was reduced to ~5 mL and the orange mixture was partitioned between EtOAc (50 mL) and H<sub>2</sub>O (50 mL). The layers were separated and the aqueous layer was extracted with EtOAc (3 x 30 mL). The combined

organic layers were washed with  $H_2O$  (25 mL), 10 wt% citric acid (2 x 25 mL),  $H_2O$  (25 mL), saturated NaHCO<sub>3</sub> (2 x 25 mL), and brine (25 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a yellow oil. The oil was chromatographed through 100 g of silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 1:3 CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to yield a mixture of **4.47** and **4.48** as a light yellow oil. The oil was dried under vacuum (2 mm Hg) for 1 h to yield a white foam (0.626 g, 69.3% yield).

Preparation of the mixture of diastereomers 4.47 and 4.48 using EDC, DMAP, and 4-pyrrolidinopyridine (Table 4.2, entry 6). Acid 3.101 (0.805 mmol) and 3.125 (0.265 g, 0.826 mmol) were dissolved in 5 mL of anhydrous CH<sub>2</sub>Cl<sub>2</sub> in a 25-mL round-bottomed flask under Ar. DMAP (25 mg, 0.205 mmol) and 4-pyrrolidinopyridine (25 mg, 0.17 mmol) were added, followed by EDC•HCl (0.660 g, 3.44 mmol). The mixture was stirred at RT under Ar and monitored by TLC (100% EtOAc - UV lamp, CAM stain). The reaction was diluted with EtOAc (20 mL) and H<sub>2</sub>O (20 mL) and the layers were separated. The aqueous layer was extracted with EtOAc ( $2 \times 20 \text{ mL}$ ), and the combined organic layers were washed with H<sub>2</sub>O (20 mL) and brine (20 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to a yellow oil. The oil was chromatographed through 50 g of silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 25% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc and yielded a mixture of **4.47** and **4.48** as a yellow oil (0.354 g, 69% yield).

Attempted deprotection of 4.47 and 4.48 using TMSOTf and 2,6-lutidine. A mixture of 4.47 and 4.48 (75.6 mg, 0.119 mmol) was dissolved in 2 mL of anhydrous  $CH_2Cl_2$  in a flame-dried 10-mL round-bottomed flask under Ar. 2,6-Lutidine (0.030 mL,

0.26 mmol) was added, followed by trimethylsilyl trifluoromethanesulfonate (0.035 mL, 0.18 mmol) dropwise. The solution was stirred at RT under Ar and monitored by TLC (100% EtOAc - CAM stain, ninhydrin stain). After 15 min, the reaction was not clean by TLC, and starting material remained. After stirring for 3 days, with no change observed by TLC, additional 2,6-lutidine (0.030 mL, 0.26 mmol) and TMSOTf (0.035 mL, 0.18 mmol) was added. The reaction was stirred at RT under Ar for 20 min, then diluted with CH<sub>2</sub>Cl<sub>2</sub> (5 mL) and saturated NaHCO<sub>3</sub> (10 mL) was added. The layers were separated and the aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (2 x 7 mL). The combined organic layers were dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a yellow oil (103.6 mg). The mixture was analyzed by LC-MS and was found to contain large amounts of 2,6-lutidine (rt = 3.80 min), and starting material 4.47 and 4.48 (rt = 9.02 min). Approximately equal amounts of 4.51 (rt = 6.10 min, MW = 535) 4.52 (rt = 6.60 min, MW = 595) were observed in the LC trace as well as 4.53.



(S)-4-((S)-1-(Benzyloxy)-2-oxoazepan-3-ylamino)-4-oxobutan-2-yl 2-amino-

**7,8-dihydroxyoctanoate (4.53).** A mixture of **4.47** and **4.48** (0.185 g, 0.291 mmol) was dissolved in 4 mL of 90% TFA/H<sub>2</sub>O in a 25-mL round-bottomed flask. The brown solution was stirred at RT for 20 min, then added dropwise to a beaker containing saturated NaHCO<sub>3</sub> (50 mL). Solid sodium carbonate was added to the mixture until the

aqueous layer was a pH of 8-9 (pH paper).  $CH_2Cl_2$  (15 mL) was added and the layers were separated. The cloudy aqueous layer was extracted with  $CH_2Cl_2$  (3 x 10 mL). The combined organic layers were dried over MgSO<sub>4</sub>, filtered, concentrated to yield crude **4.53** as a yellow oil (92.8 mg, 64.6% yield). The material was used without purification. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.40 – 7.36 (m, 5H), 4.98 – 4.85 (m, 2H), 4.46 (m, 1H), 3.66 – 3.35 (m, 5H), 2.98 (br, 3H), 2.61 – 2.42 (m, 2H), 2.00 – 1.30 (m, 17H) ppm.



(S)-4-((S)-1-(Benzyloxy)-2-oxoazepan-3-ylamino)-4-oxobutan-2-yl 7,8dihydroxy-2-((S)-2-(2-hydroxyphenyl)-4,5-dihydrooxazole-4-carboxamido)octanoate (4.54). Amine 4.53 (92.8 mg, 0.188 mmol), 2.62 (40.7 mg, 0.196 mmol), and HOBt (29.6 mg, 0.219 mmol) were dissolved in 5 mL of anhydrous  $CH_2Cl_2$  in a 25-mL roundbottomed flask under Ar. EDC•HCl (51.7 mg, 0.270 mmol) was added in one portion and the mixture was stirred at RT under Ar. After 17 h, product formation was observed by TLC (9:1 EtOAc/MeOH - UV lamp, CAM stain). The reaction was diluted with EtOAc (25 mL) and  $H_2O$  (25 mL) and the layers were separated. The cloudy aqueous layer was extracted with EtOAc (3 x 20 mL) and the combined organic layers were washed with saturated NaHCO<sub>3</sub> (2 x 20 mL),  $H_2O$  (2 x 20 mL), and brine (20 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a yellow oil. The oil was chromatographed through 10 g of silica using a solvent gradient from 100% EtOAc to 90% EtOAc/MeOH and yielded 4.54 as a colorless residue (83.1 mg, 64.7% yield). Rf = 0.28 (9:1 EtOAc/MeOH – UV lamp, CAM stain). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.77 – 7.67 (m, 1H), 7.42 - 7.35 (m, 5H), 7.21 - 7.16 (m, 0.5H), 7.09 - 7.01 (m, 2H), 7.94 -7.89 (m, 1H), 5.36 - 5.33 (m, 0.5H), 5.26 (m, 0.5H), 5.02 - 4.95 (m, 2H), 4.89 (t, J =10.5 Hz, 1H), 4.82 – 4.78 (m, 0.5H), 4.68 – 4.59 (m, 2H), 4.54 – 4.43 (m, 2H), 3.71 – 3.57 (m, 2H), 3.55 - 3.47 (m, 2H), 3.39 - 3.33 (m, 1H), 2.58 (dd, J = 10.5, 8.0 Hz, 0.5H),2.50 (dd, J = 14.5, 4.5 Hz, 0.5H), 2.41 (d, J = 7.0 Hz, 1H), 2.00 – 1.85 (m, 3H), 1.80 – 1.60 (m, 3H), 1.50 – 1.20 (m, 11H) ppm.  $^{13}$ C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  170.99, 170.97, 170.61, 170.57, 170.4, 170.23, 170.20, 170.17, 170.1, 170.0, 168.9, 168.8, 168.4, 168.3, 167.6, 167.22, 167.18, 159.7, 159.4, 135.02, 134.98, 134.82, 134.81, 134.2, 134.1, 129.5, 129.4, 128.94, 128.92, 128.58, 128.56, 128.5, 119.2, 119.1, 116.9, 116.7, 110.2, 110.0, 76.9, 71.9, 71.55, 71.45, 70.9, 69.74, 69.69, 69.5, 69.4, 69.21, 69.18, 67.9, 67.84, 67.81, 66.82, 66.76, 66.71, 66.69, 52.70, 52.67, 52.6, 52.3, 52.20, 52.15, 51.9, 51.7, 51.64, 51.60, 51.58 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>35</sub>H<sub>47</sub>N<sub>4</sub>O<sub>10</sub><sup>+</sup>, 683.2392; obsd, 683.3283.



(S)-4-Oxo-4-((S)-2-oxoazepan-3-ylamino)butan-2-yl 7,8-dihydroxy-2-((S)-2-(2-hydroxyphenyl)-4,5-dihydrooxazole-4-carboxamido)octanoate (4.55). All glassware used was washed with 6M HCl (2x), then DI water to neutral pH (pH paper),

then acetone and was flame-dried before use to remove trace amounts of Fe and other metals. Anhydrous MeOH (2 mL) was added to **4.54** (11.6 mg, 0.0170 mmol) in a 10-mL round-bottomed flask. The mixture was purged with Ar and 10 wt% Pd/C (2 mg) was added. The black mixture was purged with Ar, then with H<sub>2</sub> for 30 seconds, then was stirred under H<sub>2</sub> (balloon). After 55 min, the mixture was analyzed by TLC (9:1 EtOAc/MeOH - UV lamp, CAM stain), but only starting material was observed. Glacial acetic acid (0.001 mL, 0.017 mmol, ~1 equiv) was added to the mixture. After 30 min, no change was observed. Additional 10 wt% Pd/C was added (about 3 mg) and the mixture was stirred under H<sub>2</sub>. After 1 h, no starting material was observed by TLC, and a new, FeCl<sub>3</sub>-positive spot was observed just above the baseline. The reaction was filtered through glass filter paper. The colorless filtrate was concentrated to yield a mixture of **4.55** and other impurities as a residue (6.8 mg, 67.5% crude yield). LC/MS (5 – 80% CH<sub>3</sub>CN/10 mM NH<sub>4</sub>OAc) R<sub>t</sub> = 5.43 and 5.57 min; *m*/z [M+H]<sup>+</sup> 577.5.



(S)-4-((S)-1-Hydroxy-2-oxoazepan-3-ylamino)-4-oxobutan-2-yl 7,8-

**dihydroxy-2-((S)-2-(2-hydroxyphenyl)-4,5-dihydrooxazole-4-carboxamido)octanoate** (4.56). All glassware used was washed with 6M HCl (2x), then H<sub>2</sub>O to neutral pH (pH paper), then acetone and was oven-dried before use to remove trace amounts of metals. Anhydrous MeOH (10 mL) was added to 4.54 (70.6 mg, 0.103 mmol) in a 25-mL round-

bottomed flask. The mixture was purged with Ar (3 min) and 10 wt% Pd/C (31.5 mg, 45 wt% of 4.54) was added. The black mixture was purged with Ar (2 min), then with  $H_2$  (2 min), then was stirred under  $H_2$  (balloon). After 1 h, no starting material remained by TLC analysis of the reaction mixture (9:1 EtOAc/MeOH - UV lamp, FeCl<sub>3</sub> stain), and a FeCl<sub>3</sub>-positive spot was observed just above the baseline. The reaction was purged with Ar (3 min) and filtered through a pad of celite over glass filter paper. The filtrate was still contaminated with fine particles of Pd/C and was filtered through a 4mm acrodisc syringe filter and concentrated to yield a light orange oil. The oil was dried under vaccuum (2 mm Hg) and solidified (51.2 mg, 83.6% crude yield). The crude material was dissolved in MeOH and analyzed by LC-MS. The desired product eluted at 5.32 min. Other peaks were observed at 6.62 min, 7.15 min, and a significant impurity at 7.72 min. The crude mixture was purified by trituration with acetonitrile and provided **4.56** as a white solid (26.2 mg, 42.8% yield). LC/MS (5%-80% CH<sub>3</sub>CN/10mM NH<sub>4</sub>OAc)  $R_t =$ 5.32 min. <sup>1</sup>H NMR (500 MHz, d<sup>6</sup>-DMSO, 30 °C)  $\delta$  8.64 (m, 1H), 8.04 (dd, J = 14, 7.0 Hz, 1H), 7.62 (d, J = 7.5 Hz, 1H), 7.44 (m, 1H), 7.97 – 7.87 (m, 2H), 5.12 (m, 1H), 4.99 (m, 1H), 4.60 (q, J = 9.5 Hz, 1H), 4.50 (m, 1H), 4.44 (m, 1H), 4.21 (m, 0.5H), 4.15 (m, 0.5H), 3.86 (d, J = 11.5 H, 0.5H), 3.82 (d, J = 11.5 Hz, 0.5H), 3.51 – 3.45 (m, 1H), 3.38 – 3.21 (m, 5H), 2.54 – 2.39 (m, 1H), 1.84 – 1.58 (m, 6H), 1.47 – 1.21 (m, 8H), 1.19 (d, J = 6.5 Hz, 1.5H), 1.14 (d, J = 6.0 Hz, 1.5H) ppm. <sup>13</sup>C NMR (125 MHz, d<sup>6</sup>-DMSO, 30 °C)  $\delta$ 171.0, 170.8, 169.7, 169.6, 168.54, 168.45, 167.87, 167.78, 165.7, 133.77, 133.73, 128.0, 118.4, 116.8, 116.7, 109.98, 109.87, 70.9, 70.8, 69.0, 68.7, 68.5, 67.04, 67.00, 65.8, 52.4, 52.3, 52.2, 50.6, 41.2, 33.01, 32.98, 30.7, 30.59, 30.55, 30.4, 26.89, 26.86, 25.4, 25.33,

25.28, 24.7, 19.4, 19.2 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>29</sub>H<sub>41</sub>N<sub>4</sub>O<sub>10<sup>+</sup></sub>, 593.2823; obsd, 593.2801.



(R)-4-((S)-1-(Benzyloxy)-2-oxoazepan-3-ylamino)-4-oxobutan-2-yl 2-(*tert*butoxycarbonylamino)-6-((S)-2,2-dimethyl-1,3-dioxolan-4-yl)hexanoate (4.57). Compound 4.57 was prepared following the procedure used for 4.47 and 4.48 using EDC, DMAP, and 4-pyrrolidinopyridine. The use of acid (2S,7R)-3.101 (0.246 g, 0.742 mmol), 3.124 (0.238 g, 0.742 mmol), EDC•HCl (0.650 g, 3.39 mmol), DMAP (33.0 mg, 0.270 mmol), and 4-pyrrolidinopyridine (40.0 mg, 0.270 mmol) provided a yellow oil. Chromatography through 50 g of silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 33% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc yielded 4.58 as a light white foam (0.268 g, 57% yield). The product consisted of approximately a 2:1 mixture of epimers from <sup>1</sup>H NMR data. Rf =0.37 (100% EtOAc - CAM stain). LC/MS (5 - 80% CH<sub>3</sub>CN/10 mM NH<sub>4</sub>OAc) R<sub>t</sub> = 8.95 min. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.43 – 7.36 (m, 5H), 7.05 (d, J = 6.5 Hz, 1H), 7.02 (d, J = 6.5 Hz, 1H epimer), 5.35 - 5.19 (m, 2H), 4.98 (d, J = 10.5 Hz, 1H), 4.99 (d, J =10.0 Hz, 1H epimer), 4.90 (d, J = 10.5 Hz, 1H), 4.46 (ddd, J = 11.0, 6.5, 1.0 Hz, 1H), 4.23 (m, 1H), 4.07 - 3.98 (m, 2H), 3.64 (d, J = 11.5 Hz, 1H epimer), 3.60 (d, J = 11.5 Hz,1H), 3.53 – 3.45 (m, 2H), 2.59 – 2.45 (m, 2H), 1.98 – 1.88 (m, 2H), 1.81 – 1.61 (m, 5H), 1.51 – 1.33 (m, 19H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 171.9, 170.00 (epimer), 169.97, 168.2, 155.4, 134.93 (epimer), 134.90, 129.58 (epimer), 129.57, 128.89, 128.87 (epimer), 128.5, 108.58 (epimer), 108.56, 79.6, 76.8, 75.8, 69.3, 68.94, 68.88 (epimer), 53.6, 53.5 (epimer), 52.7, 51.7, 42.6 (epimer), 42.5, 33.3 (epimer), 33.2, 32.5, 32.4 (epimer), 31.5, 31.4, 28.3, 27.52 (epimer), 27.48, 26.9, 26.1, 25.7, 25.42, 25.40, 25.35, 25.42, 19.62 (epimer), 19.56 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>33</sub>H<sub>52</sub>N<sub>3</sub>O<sub>9</sub><sup>+</sup>, 634.3704; obsd, 634.3709.



(*R*)-4-((*S*)-1-(Benzyloxy)-2-oxoazepan-3-ylamino)-4-oxobutan-2-yl 2-(*tert*butoxycarbonylamino)-6-((*R*)-2,2-dimethyl-1,3-dioxolan-4-yl)hexanoate (4.58). Compound 4.58 was prepared following the procedure used for 4.47 and 4.48 using EDC, DMAP, and 4-pyrrolidinopyridine. The use of acid (2*S*,7*R*)-3.101 (0.276 g, 0.832 mmol), 3.124 (0.268 g, 0.837 mmol), EDC•HCl (0.730 g, 3.81 mmol), DMAP (36.5 mg, 0.299 mmol), and 4-pyrrolidinopyridine (44 mg, 0.30 mmol) provided a yellow oil. Chromatography through 50 g of silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 33% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc yielded 4.58 as a light yellow oil (0.292 g, 55% yield). The product consisted of approximately a 2:1 mixture of epimers from <sup>1</sup>H NMR data. Rf = 0.37 (100% EtOAc – CAM stain). LC/MS (5 – 80% CH<sub>3</sub>CN/10 mM NH<sub>4</sub>OAc) R<sub>t</sub> = 8.97 min. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.44 – 7.36 (m, 5H), 7.03 (m, 1H epimer), 7.00 (d, *J* = 6.0 Hz, 1H), 5.35 – 5.18 (m, 2H), 4.99 (m, 1H), 4.90 (dd, *J* = 10.5, 2.0 Hz, 1H), 4.46 (dd, *J* = 10.5, 6.5 Hz, 1H), 4.25 (dd, J = 13.5, 8.0 Hz, 1H), 4.08 – 3.99 (m, 2H), 3.62 (m, 1H), 3.53 – 3.45 (m, 2H), 2.59 – 2.45 (m, 2H), 1.99 – 1.96 (m, 1H), 1.92 – 1.88 (m, 1H), 1.82 (m, 1H), 1.75 – 1.60 (m, 4H), 1.48 – 1.31 (m, 19H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$ 171.91, 171.89, 170.0, 168.2, 155.4, 135.0, 129.6, 128.94, 128.91, 128.6, 108.6, 79.6, 76.9, 75.8, 69.4, 68.9, 53.5, 52.8, 51.7, 42.64, 42.59 (epimer), 33.32 (epimer), 33.27, 32.6 (epimer), 32.4, 31.6 (epimer), 31.5, 28.3, 27.6, 26.9, 26.2, 25.7, 25.5, 25.42, 25.38, 25.2, 19.7, 19.6 (epimer) ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>33</sub>H<sub>52</sub>N<sub>3</sub>O<sub>9</sub><sup>+</sup>, 634.3704; obsd, 634.3706.



(*S*)-4-((*S*)-1-(Benzyloxy)-2-oxoazepan-3-ylamino)-4-oxobutan-2-yl 2-(*tert*butoxycarbonylamino)-6-((*S*)-2,2-dimethyl-1,3-dioxolan-4-yl)hexanoate (4.59). Compound 4.59 was prepared following the procedure used for 4.47 and 4.48 using EDC, DMAP, and 4-pyrrolidinopyridine. The use of (2R,7S)-3.101 (0.141 g, 0.425 mmol), 4pyrrolidinopyridine (23.0 mg, 0.155 mmol), EDC+HCl (0.356 g, 1.86 mmol), DMAP (22.1 mg, 0.181 mmol), and 3.125 (0.136 g, 0.425 mmol) provided a colorless oil. Chromatography through 30 g of silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 33% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc yielded 4.59 as a yellow oil (0.177 g, 66% yield). Isolated as a mixture of epimers. Rf = 0.36 (100% EtOAc – CAM stain). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.44 – 7.32 (m, 5H), 7.12 (m, 1H), 5.40 (d, *J* = 8.5 Hz, 1H), 5.30 (m, 2H), 5.00 (dd, *J* = 10.0, 2.5 Hz, 1H), 4.89 (dd, *J* = 10.5, 3.0 Hz, 1H), 4.50 – 4.43 (m, 1H), 4.25 (m, 1H), 4.06 – 3.98 (m, 2H), 3.68 – 3.60 (m, 1H), 3.53 – 3.44 (m, 2H), 2.58 (dd, *J* = 15.0, 7.0 Hz, 1H), 2.51 – 2.45 (m, 1H), 2.04 – 1.97 (m, 1H), 1.94 – 1.88 (m, 1H), 1.86 – 1.80 (m, 1H), 1.76 – 1.58 (m, 4H), 1.50 – 1.25 (m, 19H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) & 171.75 (epimer), 171.70, 169.9, 169.8 (epimer), 168.1, 168.0 (epimer), 155.34, 155.25 (epimer), 134.80, 134.77 (epimer), 129.4, 128.73 (epimer), 128.69, 128.4, 108.4, 79.4 (epimer), 79.3, 76.6, 75.7 (epimer), 75.6, 69.22 (epimer), 69.20, 68.8, 53.4, 53.3 (epimer), 52.5, 52.4, 51.6 (epimer), 51.5, 42.5, 42.3 (epimer), 33.2 (epimer), 33.1, 32.2 (epimer), 32.0, 31.25, 31.22, 28.2, 28.1 (epimer), 27.3, 26.7, 26.0, 25.5, 25.3, 25.2, 25.1, 19.5 ppm.



(S)-4-((S)-1-(Benzyloxy)-2-oxoazepan-3-ylamino)-4-oxobutan-2-yl 2-(*tert*-

butoxycarbonylamino)-6-((R)-2,2-dimethyl-1,3-dioxolan-4-yl)hexanoate (4.60). Compound 4.60 was prepared following the procedure used for 4.47 and 4.48 using EDC, DMAP, and 4-pyrrolidinopyridine. The use of (2*S*,7*R*)-3.101 (33.8 mg, 0.102 mmol), 4pyrrolidinopyridine (6.8 mg, 0.046 mmol), EDC•HCl (92.2 mg, 0.481 mmol), DMAP (8.0 mg, 0.65 mmol), and 3.125 (33.2 mg, 0.104 mmol) provided a yellow residue. Chromatography through 10 g of silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 50% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc yielded 4.60 as a yellow foam (40.0 mg, 62% yield). Isolated as a mixture of epimers. Rf = 0.36 (100% EtOAc – CAM stain). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>. )  $\delta$  7.44 – 7.37 (m, 5H), 7.07 (d, *J* = 6.0 Hz, 1H), 5.35 – 5.18 (m, 2H), 5.00 (m, 1H), 4.90 (m, 1H), 4.49 – 4.42 (m, 1H), 4.28 – 4.22 (m, 1H), 4.08 – 3.99 (m, 2H), 3.68 – 3.60 (m, 1H), 3.54 – 3.45 (m, 2H), 2.59 (dd, *J* = 15.0, 6.0 Hz, 1H), 2.46 (dd, *J* = 15.0, 6.0 Hz, 1H), 2.05 – 1.98 (m, 1H), 1.94 – 1.87 (m, 1H), 1.85 – 1.57 (m, 5H), 1.53 – 1.25 (m, 19H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  171.9, 170.02 (epimer), 169.97, 168.2 (epimer), 168.1, 155.4, 134.94 (epimer), 134.90, 129.6, 128.9, 128.6, 108.6, 79.64, 79.55 (epimer), 76.9, 75.8, 69.4, 68.9, 53.5, 52.7, 52.6 (epimer), 51.8, 51.7 (epimer), 42.7 (epimer), 42.5, 33.31 (epimer), 33.25, 32.4, 31.4, 28.31 (epimer), 28.27, 27.5, 26.9, 26.1, 25.7, 25.4, 25.3, 19.7 (epimer), 19.6 ppm.



(*S*)-((*R*)-4-((*S*)-1-(Benzyloxy)-2-oxoazepan-3-ylamino)-4-oxobutan-2-yl) 2amino-7,8-dihydroxyoctanoate (4.61). Compound 4.61 was prepared following the same procedure used for 4.53. Compound 4.57 (0.175 g, 0.276 mmol) provided crude 4.61 as an off-white foam (98.9 mg, 73% yield). Isolated as a mixture of epimers. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.40 – 7.14 (m, 6H), 5.32 – 5.23 (m, 1H), 4.95 (d, *J* = 10.2 Hz, 1H), 4.85 (d, *J* = 10.2 Hz, 1H), 4.56 – 4.41 (m, 1H), 3.66 – 3.33 (m, 9H), 2.58 – 2.43 (m, 2H), 1.93 – 1.22 (m, 17H) ppm. <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  174.0, 170.1, 169.9 (epimer), 168.6, 168.3 (epimer), 134.8, 129.5, 128.8, 128.5, 76.8, 71.6 (epimer), 71.4, 68.8, 66.5, 54.0, 52.6, 51.6, 42.3, 33.7, 32.5, 31.3, 27.4 (epimer), 27.3, 26.0, 24.9, 24.7, 19.7 ppm.



(S)-((R)-4-((S)-1-(Benzyloxy)-2-oxoazepan-3-ylamino)-4-oxobutan-2-yl) 7.8dihydroxy-2-((S)-2-(2-hydroxyphenyl)-4,5-dihydrooxazole-4-carboxamido)octanoate (4.62). Compound 4.62 was prepared following the same procedure used for 4.54. Oxazoline 2.62 (44.4 mg, 0.214 mmol), EDC•HCl (53 mg, 0.276 mmol), HOBt (31.9 mg, 0.236 mmol), and 4.61 (98.9 mg, 0.200 mmol) provided an oil. Chromatography through 15 g of silica using a solvent gradient from 100% EtOAc to 95% EtOAc/MeOH yielded **4.62** as a white solid (73 mg, 53% yield). Isolated as a mixture of epimers. Rf = 0.27(9:1 EtOAc/MeOH – CAM stain, UV lamp). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 11.45 (br, 1H), 6.68 (dt, J = 8.0, 1.5 Hz, 1H), 7.43 – 7.35 (m, 5H), 7.17 – 7.01 (m, 3H), 6.90 (t, J =7.5 Hz, 1H), 5.37 - 5.23 (m, 1H), 4.99 - 4.93 (m, 2H), 4.89 - 4.86 (m, 1H), 4.68 - 4.62 (m, 2H), 4.58 – 4.54 (m, 1H), 4.47 – 4.42 (m, 1H), 3.72 – 3.34 (m, 5H), 2.62 – 2.43 (m, 2H), 1.96 - 1.84 (m, 3H), 1.78 - 1.64 (m, 3H), 1.50 - 1.26 (m, 11H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 170.9 (epimer), 170.8, 170.4 (epimer), 170.3, 170.1 (epimer), 170.0, 168.4, 168.3 (epimer), 167.9, 167.6 (epimer), 159.61, 159.59 (epimer), 134.9 (epimer), 134.8, 134.2, 129.53, 129.49 (epimer), 128.91, 128.85 (epimer), 128.6, 128.5 (epimer), 119.1, 116.89 (epimer), 116.87, 110.0, 76.8, 71.5 (epimer), 71.2, 69.45 (epimer), 69.41, 67.94 (epimer), 67.90, 66.8, 66.7 (epimer), 52.64 (epimer), 52.58, 52.2, 52.0 (epimer), 51.7 (epimer), 51.6, 42.4 (epimer), 42.2, 32.5, 31.9 (epimer), 31.8, 31.5, 31.3 (epimer), 27.4 (epimer), 27.3, 26.1 (epimer), 26.0, 24.72, 24.69, 24.5, 24.4, 19.69, 19.66 ppm.



(*R*)-((*R*)-4-((*S*)-1-(Benzyloxy)-2-oxoazepan-3-ylamino)-4-oxobutan-2-yl) 2amino-7,8-dihydroxyoctanoate (4.63). Compound 4.63 was prepared following the same procedure used for 4.53. Compound 4.58 (0.254 g, 0.401 mmol) provided 4.63 as an off-white foam (125 mg, 63% yield). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.40 – 7.33 (m, 5H), 7.15 (d, *J* = 6.6 Hz, 1H), 5.27 (m, 1H), 4.95 (d, *J* = 10.2 Hz, 1H), 4.85 (d, *J* = 10.2 Hz, 1H), 4.43 (dd, *J* = 10.2, 6.3 Hz, 1H), 3.64 – 3.33 (m, 6H), 2.89 (br, 3H), 2.58 – 2.43 (m, 2H), 1.93 – 1.83 (m, 2H), 1.74 – 1.21 (m, 15H) ppm. <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$ 174.9, 170.07, 170.00 (epimer), 168.5, 134.8, 129.5, 128.8, 128.5, 76.7, 71.8, 71.7 (epimer), 68.64 (epimer), 68.56, 66.5, 54.1 (epimer), 54.0, 52.6, 51.6, 42.5, 34.1, 32.7, 31.3, 27.4, 26.0, 25.3, 25.0, 19.8 ppm.



(R)-((R)-4-((S)-1-(Benzyloxy)-2-oxoazepan-3-ylamino)-4-oxobutan-2-yl) 7.8dihydroxy-2-((S)-2-(2-hydroxyphenyl)-4,5-dihydrooxazole-4-carboxamido)octanoate (4.64). Compound 4.64 was prepared following the same procedure used for 4.54. Oxazoline 2.62 (55.8 mg, 0.269 mmol), EDC•HCl (67.0 mg, 0.350 mmol), HOBt (41.2 mg, 0.305 mmol), and 4.63 (0.125 g, 0.253 mmol) provided an oil. Chromatography through 20 g of silica using a solvent gradient from 100% EtOAc to 95% EtOAc/MeOH yielded 4.64 as a white solid (94.4 mg, 55% yield). Rf = 0.25 (9:1 EtOAc/MeOH – UV lamp, CAM stain). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.65 (dd, J = 7.5, 1.0 Hz, 1H), 7.40 – 7.32 (m, 6H), 7.13 (m, 2H), 6.99 (d, J = 8.5 Hz, 1H), 6.87 (t, J = 7.5 Hz, 1H), 5.31 (m, 1H), 5.25 (m, 1H epimer), 4.96 - 4.90 (m, 2H), 4.86 - 4.83 (m, 1H), 4.63 - 4.58 (m, 2H), 4.55 - 4.51 (m, 1H), 4.47 - 4.40 (m, 1H), 3.63 - 3.57 (m, 2H), 3.52 (dd, J = 11.0, 3.0 Hz, 1H), 3.47 – 3.34 (m, 1H), 3.40 – 3.36 (m, 1H epimer), 3.33 (dd, J = 11, 7.5 Hz, 1H), 2.57 (dd, J = 15.0, 7.5 Hz, 1H), 2.49 (dd, J = 15.0, 5.0 Hz, 1H), 2.40 (dd, J = 15.0, 5.0 Hz, 1H)epimer), 1.93 – 1.80 (m, 3H), 1.73 – 1.61 (m, 3H), 1.47 – 1.26 (m, 11H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 170.9, 170.7 (epimer), 170.5 (epimer), 170.2, 170.02, 170.0 (epimer), 168.4, 168.3 (epimer), 167.8 (epimer), 167.5, 159.54 (epimer), 159.50, 134.8, 134.7 (epimer), 134.2 (epimer), 134.1, 129.5 (epimer), 129.4, 128.9 (epimer), 128.8, 128.5, 128.4 (epimer), 119.0, 116.8, 109.9, 76.9, 71.7, 71.6 (epimer), 69.4, 69.3, 67.9,

67.8 (epimer), 66.65 (epimer), 66.61, 52.6, 52.1, 51.6, 42.3, 32.5 (epimer), 32.4, 31.7, 31.3, 27.4, 27.3 (epimer), 26.03, 26.00 (epimer), 25.0, 24.7, 19.6 ppm.



(*S*)-((*R*)-4-((*S*)-1-Hydroxy-2-oxoazepan-3-ylamino)-4-oxobutan-2-yl) 7,8dihydroxy-2-((*S*)-2-(2-hydroxyphenyl)-4,5-dihydrooxazole-4-carboxamido)octanoate (4.65). Compound 4.65 was prepared following the same procedure for 4.56. Compound 4.62 (12.0 mg, 0.018 mmol) was subjected to hydrogenolysis using 10 wt% Pd/C (6 mg) for 20 min. Trituration with acetonitrile yielded 4.65 as a white solid (3 mg, 30%). <sup>1</sup>H NMR (500 MHz, d<sup>6</sup>-DMSO, 30 °C)  $\delta$  8.62 (m, 1H), 8.02 (d, *J* = 7.0 Hz, 1H), 7.60 (d, *J* = 8.0 Hz, 1H), 7.38 (m, 1H), 6.91 – 6.78 (m, 2H), 5.10 (sextet, *J* = 6.5 Hz, 1H), 4.96 (m, 1H), 4.58 (m, 1H), 4.43 (dd, *J* = 9.5, 7.5 Hz, 2H), 4.19 (m, 1H), 3.84 (dd, *J* = 16.0, 11.5 Hz, 1H), 3.48 (m, 1H), 3.38 (m, 1H), 3.30 – 3.20 (m, 3H), 2.51 (m, 1H), 2.39 (dd, *J* = 14.5, 6.0 Hz, 1H), 1.82 (m, 1H), 1.72 – 1.58 (m, 3H), 1.46 – 1.34 (m, 3H), 1.46 – 1.18 (m, 10H) ppm. HRMS (FAB) *m*/*z* [M+H]<sup>+</sup> calcd for C<sub>28</sub>H<sub>41</sub>N<sub>4</sub>O<sub>10</sub><sup>+</sup>, 593.2823; obsd, 593.2801.



(R)-((R)-4-((S)-1-Hydroxy-2-oxoazepan-3-ylamino)-4-oxobutan-2-yl) 7,8dihydroxy-2-((S)-2-(2-hydroxyphenyl)-4,5-dihydrooxazole-4-carboxamido)octanoate (4.66). Compound 4.66 was prepared following the same procedure for 4.56. Compound **4.64** (47.0 mg, 0.069 mmol) was subjected to hydrogenolysis using 10 wt% Pd/C (18 mg) for 20 min. Trituration with acetonitrile yielded 4.66 as a white solid (13 mg, 32% yield). <sup>1</sup>H NMR (500 MHz, d<sup>6</sup>-DMSO, 30 °C)  $\delta$  8.67 (br, 1H), 8.02 (d, J = 7.5 Hz, 1H epimer), 7.98 (d, J = 7.0 Hz, 1H), 7.59 (d, J = 8.0 Hz, 1H), 7.40 (m, 1H), 6.92 (m, 1H), 6.83 (m, 1H), 5.11 (q, J = 6.5 Hz, 1H), 4.98 (dd, J = 9.5, 7.5 Hz, 1H), 4.60 (t, J = 9 Hz, 1H), 4.49 -4.41 (m, 2H), 4.20 (m, 1H), 3.84 (dd, J = 16.0, 11.5 Hz, 1H), 3.46 (dd, J = 15.5, 5.0 Hz, 1H), 3.28 - 3.19 (m, 3H), 2.51 (m, 1H), 2.40 (dd, J = 14.5, 6.5 Hz, 1H), 1.83 - 1.58 (m, 6H), 1.46 – 1.16 (m, 11H) ppm. <sup>13</sup>C NMR (125 MHz, d<sup>6</sup>-DMSO, 30 °C) δ 170.9, 170.8 (epimer), 169.8, 168.6, 168.5 (epimer), 167.9, 167.9 (epimer), 165.75, 165.69 (epimer), 133.7, 128.0, 118.2 (br), 116.8 (br), 109.9 (br), 70.9, 68.9, 68.7, 67.1, 65.9, 52.3, 52.2 (epimer), 50.7, 41.3, 41.2 (epimer), 33.0, 30.7, 30.3 (epimer), 30.1, 26.8, 25.4 (epimer), 25.3, 24.72 (epimer), 24.66, 19.3, 19.2 (epimer) ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>28</sub>H<sub>41</sub>N<sub>4</sub>O<sub>10</sub><sup>+</sup>, 593.2823; obsd, 593.2825.

8.4 Experimental procedures for chapter 5



(*S*)-Methyl 2-benzamido-3-hydroxypropanoate (5.20). Compound 5.20 was prepared following the same procedure for compound 3.5 using EDC-mediated amide formation. Benzoic acid (3.55 g, 29.0 mmol), **3.18** (4.03 g, 25.9 mmol), EDC•HCl (5.57 g, 29.1 mmol), and triethylamine (3.85 mL, 27.4 mmol) provided 5.20 as an oil that solidified upon storage (3.64 g, 63% yield). mp = 84-86 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.78 (d, *J* = 6.9 Hz, 2H), 7.50 – 7.28 (m, 3H), 4.81 (dt, *J* = 7.5, 3.9 Hz, 1H), 4.06 – 3.94 (m, 2H), 3.75 (s, 3H), 3.53 (br, 1H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  171.0, 167.8, 133.3, 131.9, 128.5, 127.1, 63.0, 55.1, 52.7 ppm.



(*S*)-Methyl 3-hydroxy-2-(4-methoxybenzamido)propanoate (5.21). Compound 5.21 was prepared following the same procedure for compound 3.5 using EDC-mediated amide formation. 4-Anisic acid (0.558 g, 3.67 mmol), 3.18 (0.632 g, 4.06 mmol), EDC•HCl (0.780 g, 4.07 mmol), and triethylamine (0.60 mL, 4.3 mmol) provided 5.21 as a colorless oil (0.512 g, 55% yield). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.79 (m, 2H), 7.08 (d, *J* = 6.9 Hz, 1H), 6.90 (m, 2H), 4.84 (td, *J* = 7.2, 3.6 Hz, 1H), 4.08 – 3.99 (m, 2H), 3.84 (s, 3H), 3.81 (s, 3H) ppm.



(*S*)-Methyl 3-hydroxy-2-(4-nitrobenzamido)propanoate (5.22). Compound 5.22 was prepared following the same procedure for compound 3.5 using EDC-mediated amide formation. 4-Nitrobenzoic acid (0.519 g, 3.11 mmol), 3.18 (0.413 g, 3.44 mmol), EDC•HCl (0.660 g, 3.44 mmol), and triethylamine (0.50 mL, 3.6 mmol) provided 5.22 as a light yellow solid (0.304 g, 36% yield). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  8.29 (m, 2H), 7.99 (m, 2H), 7.22 (d, *J* = 7.2 Hz, 1H), 4.88 (td, *J* = 7.0, 3.4 Hz, 1H), 4.14 (dd, *J* = 11.4, 3.6 Hz, 1H), 4.05 (dd, *J* = 11.4, 3.6 Hz, 1H), 3.84 (s, 3H), 2.40 (br, 1H) ppm.



Methyl 2-benzamidoacrylate (5.23). Compound 5.20 (1.206 g, 5.403 mmol) was dissolved in 50 mL of dry  $CH_2Cl_2$  in a flame-dried 100-mL round-bottomed flask under Ar. EDC•HCl (1.238 g, 6.458 mmol) was added to the solution in one portion followed by CuCl (160.7 mg, 1.623 mmol). The reaction was wrapped in foil and stirred under Ar at rt. After 2 h, TLC of the reaction (1:1 hexanes/EtOAc - UV lamp) indicated 5.20 had been completely consumed and a new, less polar compound was observed. After 3 h, H<sub>2</sub>O (100 mL) was added to the reaction and the layers were separated. The aqueous layer was extracted with  $CH_2Cl_2$  (2 x 20 mL) and the combined  $CH_2Cl_2$  layers were washed with H<sub>2</sub>O (100 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a yellow oil. The oil was purified through a plug (30 g) of silica using 100%  $CH_2Cl_2$  and yielded 5.23 as a colorless, cloudy liquid (1.06 g, 96% yield). <sup>1</sup>H NMR (500 MHz,

CDCl<sub>3</sub>)  $\delta$  8.49 (br-s, 1H), 7.75 (dd, J = 5.1, 0.9 Hz, 2H), 7.43 (m, 1H), 7.37 (m, 2H), 6.71 (s, 1H), 5.90 (d, J = 0.6 Hz, 1H), 3.76 (s, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  165.3, 164.3, 133.9, 131.7, 130.8, 128.4, 126.6, 108.5, 52.7 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>11</sub>H<sub>12</sub>NO<sub>3</sub><sup>+</sup>, 206.0817; obsd, 206.0818.



Benzyl 2-benzamidoacrylate (5.24). Compound 5.24 was prepared following the same procedure for 5.23. Compound 3.9 (1.51 g, 5.03 mmol), EDC•HCl (1.17 g, 6.12 mmol), and CuCl (0.150 g, 1.51 mmol) provided an oil. Chromatography through silica using 100% CH<sub>2</sub>Cl<sub>2</sub> yielded 5.24 as a white solid (1.45 g, 99% yield). mp = 51-52 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 8.56 (br-s, 1H), 7.84 (dd, J = 7.0, 1.5 Hz, 2H), 7.55 (m, 1H), 7.47 (m, 2H), 7.41 – 7.36 (m, 5H), 6.82 (s, 1H), 6.06 (m, 1H), 5.32 (s, 2H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 165.7, 164.2, 135.0, 134.2, 132.0, 131.0, 128.8, 128.7, 128.6, 128.2, 126.9, 109.1, 67.9 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>17</sub>H<sub>16</sub>NO<sub>3</sub><sup>+</sup>, 282.1130; obsd, 282.1140.



Methyl 2-(4-methoxybenzamido)acrylate (5.25). Compound 5.25 was prepared following the same procedure for 5.23. Compound 5.21 (0.495 g, 1.95 mmol), EDC•HCl (0.448 g, 2.33 mmol), and CuCl (59.7 mg, 0.603 mmol) provided an oil. Chromatography through silica using 100%  $CH_2Cl_2$  yielded 5.25 as a white solid (0.426

g, 93% yield). mp = 47.0-47.5 °C. Rf = 0.47 (1:1 hexanes/EtOAc – UV lamp). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  8.44 (br, 1H), 7.77 (m, 2H), 6.93 (m, 2H), 6.73 (s, 1H), 5.93 (d, *J* = 1.2 Hz, 1H), 3.86 (s, 3H), 3.83 (s, 3H) ppm. <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  165.2, 164.8, 162.6, 131.0, 128.8, 126.4, 113.9, 108.3, 55.4, 43.0 ppm. HRMS (FAB) *m/z* [M+H]<sup>+</sup> calcd for C<sub>12</sub>H<sub>14</sub>NO<sub>4</sub><sup>+</sup>, 236.0923; obsd, 236.0922.



Methyl 2-(4-nitrobenzamido)acrylate (5.26). Compound 5.26 was prepared following the same procedure for 5.23. Compound 5.22 (0.289 g, 1.08 mmol), EDC•HCl (0.249 g, 1.30 mmol), CuCl (36.0 mg, 0.363 mmol) provided an oil. Chromatography through silica using 100% CH<sub>2</sub>Cl<sub>2</sub> yielded 5.26 as a white solid (0.244 g, 90% yield). mp = 158-159 °C. Rf = 0.45 (1:1 hexanes/EtOAc – UV lamp). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 8.57 (br, 1H), 8.33 (m, 2H), 8.00 (m, 2H), 6.81 (s, 1H), 6.06 (m, 1H), 3.91 (s, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 164.5, 163.6, 149.8, 139.6, 130.6, 128.2, 124.0, 110.0, 53.3 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>11</sub>H<sub>11</sub>N<sub>2</sub>O<sub>5</sub><sup>+</sup>, 251.0668; obsd, 251.0670.



(S)-Benzyl 3-hydroxy-2-(4-nitrobenzamido)propanoate (5.27). Compound 5.27 was prepared following the same procedure for 3.5 using acid chloride-mediated amide formation. *p*-nitrobenzoic acid (1.01 g, 6.06 mmol), oxalyl chloride (1.10 mL,

12.6 mmol), and catalytic DMF provided the acid chloride intermediate as a yellow semisolid. **3.4** (1.42 g, 6.12 mmol) and diisopropylethylamine (2.70 mL, 15.5 mmol) yielded a peach-colored solid. Chromatography through 150 g of silica using a solvent gradient from 90% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to 60% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc yielded **5.27** as a light yellow solid (1.40 g, 67% yield). mp = 129.5-130.0 °C. Rf = 0.15 (1:1 hexanes/EtOAc - UV lamp). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  8.29 (d, *J* = 8.5 Hz, 2H), 7.99 (d, *J* = 9.0 Hz, 2H), 7.37 (m, 5H), 7.16 (d, *J* = 6.5 Hz, 1H), 5.29 (d, *J* = 12.0 Hz, 1H), 5.25 (d, *J* = 12.0 Hz, 1H), 4.92 (dt, *J* = 7.5, 3.5 Hz, 1H), 4.16 (dd, *J* = 11.5, 3.5 Hz, 1H), 2.18 (br, 1H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  170.1, 165.5, 139.1, 134.9, 128.7, 128.4, 128.3, 123.9, 67.9, 63.2, 55.3 ppm. HRMS (FAB) *m*/z [M+H]<sup>+</sup> calcd for C<sub>17</sub>H<sub>17</sub>N<sub>2</sub>O<sub>6</sub><sup>+</sup>, 345.1087; obsd, 345.1111.



**Benzyl 2-(4-nitrobenzamido)acrylate (5.28).** Compound **5.28** was prepared following the same procedure for **5.23**. Compound **5.27** (0.945 g, 2.75 mmol), EDC•HCl (0.585 g, 3.05 mmol), and CuCl (83.9 mg, 0.847 mmol) provided an oil. Chromatography through silica using 100% CH<sub>2</sub>Cl<sub>2</sub> yielded **5.28** as a white solid (0.869 g, 97% yield). mp = 144-145 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  8.59 (br-s, 1H), 8.31 (dt, *J* = 9.0, 2.1 Hz, 2H), 7.98 (m, 2H), 7.40 (m, 5H), 6.83 (s, 1H), 6.12 (d, *J* = 1.2 Hz, 1H), 5.32 (s, 2H) ppm. <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  163.9, 163.5, 149.8, 139.6, 134.8, 130.7, 128.7, 128.2, 128.1, 124.0, 110.2, 68.1 ppm. HRMS (FAB) *m/z* [M+H]<sup>+</sup> calcd for C<sub>17</sub>H<sub>15</sub>N<sub>2</sub>O<sub>5</sub><sup>+</sup>, 327.0981; obsd, 327.0984.

General procedure for one-pot preparation of 2-amidoacrylates from benzoic acid and serine esters. Acrylate 5.24. Anhydrous CH<sub>2</sub>Cl<sub>2</sub> (50 mL) was added to 3.4 (1.01 g, 4.37 mmol) and benzoic acid (0.542 g, 4.44 mmol) in a flame-dried singlenecked 200-mL round-bottomed flask under Ar. Triethylamine (0.65 mL, 4.6 mmol) was added, followed by EDC•HCl (1.83 g, 9.55 mmol). The mixture was stirred at rt under Ar. After 1 h, the reaction was progressing by TLC (1:1 hexanes/EtOAc - UV lamp, ninhydrin stain). After 2.5 h, CuCl (0.137 g, 1.38 mmol) was added and the color of the solution turned light blue/green in color. After 4 h, there was no change in the reaction by TLC and more EDC•HCl (0.9 g, 4.7 mmol, ~1 equiv) was added. After about 10 h, the reaction color had changed to dark greenish/yellow, but was still incomplete by TLC. More CuCl (about 0.1 g, ~1 mmol, 22 mol%) was added and the solution was stirred at rt. The color of the reaction changed to orange and the reaction was complete. The reaction was diluted with CH<sub>2</sub>Cl<sub>2</sub> (50 mL) and H<sub>2</sub>O (100 mL) was added. The mixture was diluted with brine (100 mL) and EtOAc (~400 mL). The layers were separated and the organic layer was washed with brine (2 x 50 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to yield a yellow oil. The oil was chromatographed through an Analogix SF 25-40g silica column using 100%  $CH_2Cl_2$  and yielded **5.24** as a white solid (0.945 g, 77%) yield).



(2*S*,3*R*)-Methyl 2-benzamido-3-hydroxybutanoate (5.30). Compound 5.30 was prepared following the same procedure for compound 3.5 using EDC-mediated amide
formation. Benzoic acid (2.39 g, 19.6 mmol), **5.29** (2.97 g, 17.5 mmol), EDC•HCl (4.03 g, 21.0 mmol), and triethylamine (2.65 mL, 18.9 mmol) provided a yellow oil that solidified upon standing. Chromatography through an Analogix F40M column using a solvent gradient from 90% to 60% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc yielded **5.30** as a white solid (2.61 g, 63% yield). mp = 95-96 °C. Rf = 0.5 (1:1 hexanes/EtOAc – UV lamp). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.82 (d, *J* = 7.5 Hz, 2H), 7.48 (m, 1H), 7.39 (t, *J* = 7.5 Hz, 2H), 7.13 (d, *J* = 9.0 Hz, 1H), 4.78 (dd, *J* = 9.0, 2.5 Hz, 1H), 4.42 (qd, *J* = 6.5, 2.5 Hz, 1H), 3.74 (s, 3H), 3.17 (br, 1H), 1.24 (d, *J* = 6.5 Hz, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  171.5, 168.0, 133.5, 131.9, 128.5, 127.2, 68.0, 57.7, 52.6, 20.0 ppm.



Methyl 2-benzamidobut-2-enoate (5.31). Compound 5.31 was prepared following the same procedure for 5.23. Compound 5.30 (1.21 g, 5.08 mmol), EDC•HCl (1.18 g, 6.13 mmol), and CuCl (0.150 g, 1.52 mmol) were stirred at rt for 2 h and concentrated. The mixture was chromatographed through 30 g of silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 80% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc and yielded 5.31 as a pale yellow oil that solidified upon storage (1.07 g, 97% yield). 5.31 was isolated as a 29:1 mixture of *Z:E* isomers based on NMR data. mp = 66-68 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  8.22 (br, 1H, *E* isomer), 7.86 (m, 2H, *Z* isomer), 7.79 (m, 2H, *E* isomer), 7.65 (br, 1H, *Z* isomer), 7.52 (t, *J* = 7.5 Hz, 1H), 7.44 (m, 2H), 7.37 (q, *J* = 8.0 Hz, 1H, *E* isomer), 6.88 (q, *J* = 7 Hz, 1H, *Z* isomer), 3.86 (s, 3H, *E* isomer), 3.76 (s, 3H, *Z* isomer), 2.12 (d, *J* = 7.5 Hz, 3H, *E* isomer), 1.83 (d, *J* = 7.0 Hz, 3H, *Z* isomer) ppm. <sup>13</sup>C NMR (125 MHz,

CDCl<sub>3</sub>)  $\delta$  165.4 (*E*), 165.1 (*Z*), 133.85 (*E*), 133.77 (*Z*), 131.9 (*Z*), 131.7 (*E*), 128.6 (*Z*), 127.4 (*Z*), 126.8 (*E*), 126.0 (*E*), 52.4, 15.0 (*Z*), 14.1 (*E*) ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>12</sub>H<sub>14</sub>NO<sub>3</sub><sup>+</sup>, 220.0974; obsd, 220.0975.



3-hydroxy-2-(4-methylphenylsulfonamido)propanoate (S)-Methyl (5.32). Compound 3.18 (2.107 g, 13.54 mmol), p-toluenesulfonyl chloride (2.87 g, 15.1 mmol), and anhydrous  $CH_2Cl_2$  (100 mL) were added to a flame-dried single-necked 250-mL round-bottomed flask under Ar. N,N-Diisopropyl-N-ethylamine (5.50 mL, 31.6 mmol) was added and the solid material dissolved. The resultant solution was stirred at rt under Ar overnight. After 22 h, the reaction was analyzed by TLC (1:1 hexanes/EtOAc and 9:1 CH2Cl2/EtOAc - UV lamp) and one new polar spot was observed (along with unreacted TsCl). The light yellow solution was concentrated by rotary evaporation and the resultant residue was partitioned between EtOAc (100 mL) and  $H_2O$  (100 mL). The aqueous layer was extracted with EtOAc (2 x 50 mL) and the combined organic layers were washed with saturated NaHCO<sub>3</sub> (2 x 50 mL), H<sub>2</sub>O (50 mL), 1M HCl (2 x 50 mL), H<sub>2</sub>O (50 mL), and brine (2 x 50 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to yield an orange oil. The oil solidified upon standing. The crude material was chromatographed through a Biotage 40S column using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 50% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc and yielded **5.32** as a white solid (3.04 g, 82% yield). mp = 89-90 °C. Rf = 0.11 (1:1) hexanes/EtOAc – UV lamp). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.75 (d, J = 8.0 Hz, 2H), 7.31 (d, J = 8.0 Hz, 2H), 5.85 (d, J = 8.0 Hz, 1H), 4.01 (dt, J = 8.0, 3.5 Hz, 1H), 3.89 (m,

2H), 3.61 (s, 3H), 2.61 (br, 1H), 2.42 (s, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  170.1, 143.9, 136.4, 129.7, 127.2, 63.6, 57.6, 52.9, 21.5 ppm. HRMS (FAB) *m*/*z* [M+H]<sup>+</sup> calcd for C<sub>11</sub>H<sub>16</sub>NO<sub>5</sub>S<sup>+</sup>, 274.0749; obsd, 274.0725.



Methyl 2-(4-methylphenylsulfonamido)acrylate (5.33). Compound 5.33 was prepared following the same procedure for 5.23. Compound 5.32 (1.01 g, 3.71 mmol), EDC•HCl (0.826 g, 4.31 mmol), and CuCl (0.120 g, 1.21 mmol) were stirred at rt for 1 h, concentrated, and chromatographed through a Biotage 40S column using 100% CH<sub>2</sub>Cl<sub>2</sub> and yielded 5.33 as a free-flowing yellow solid (0.924 g, 98% yield). mp = 99-100 °C. Rf = 0.45 (1:1 hexanes/EtOAc – UV lamp). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 7.75 (d, *J* = 8.0 Hz, 2H), 7.30 (d, *J* = 8.5 Hz, 2H), 5.67 (m, 1H), 5.64 (m, 1H), 3.76 (s, 3H), 2.42 (s, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 163.6, 144.3, 135.3, 130.8, 129.7, 127.5, 106.8, 53.2, 21.5 ppm. HRMS (FAB) *m*/*z* [M+H]<sup>+</sup> calcd for C<sub>11</sub>H<sub>15</sub>NO<sub>4</sub>S<sup>+</sup>, 256.0644; obsd, 256.0668.



Methyl 2-acetamidoacrylate (5.34). To a flame-dried 50-mL round-bottomed flask containing 3.18 (2.021 g, 12.99 mmol) under Ar was added 15 mL of anhydrous triethylamine. The mixture was stirred at RT for 10 min, then cooled in a crushed ice/H<sub>2</sub>O bath. Acetic anhydride (3.00 mL, 32.0 mmol) was added slowly the reaction

under Ar. The mixture was stirred in the ice/H<sub>2</sub>O bath for 15 min, then stirred at rt. After stirring 1 h, the reaction color changed from a white mixture to a yellow mixture. After 4 days, the brown mixture was partitioned between H<sub>2</sub>O (100 mL) and EtOAc (50 mL). The layers were separated and the aqueous layer was extracted with EtOAc (2 x 50 mL). The combined organic layers were washed with 1M HCl (3 x 50 mL), H<sub>2</sub>O (50 mL), and brine (50 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield an orange oil (~0.8 g crude material). The oil solidified upon storage at 10 °C. The crude material was purified through 100 g of silica gel using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 90% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc and yielded **5.34** as a yellow oil that solidified upon standing (0.681 g, 37% yield). mp = 49-50 °C. Rf = 0.30 (1:1 hexanes/EtOAc – UV lamp). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.76 (br, 1H), 6.54 (s, 1H), 5.82 (d, *J* = 1.5 Hz, 1H), 3.80 (s, 3H), 2.09 (s, 3H) ppm. <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  168.8, 164.4, 130.8, 108.6, 52.8, 24.5 ppm. HRMS (FAB) *m*/z [M+H]<sup>+</sup> calcd for C<sub>6</sub>H<sub>10</sub>NO<sub>3</sub><sup>+</sup>, 144.0661; obsd, 144.0650.



**Methyl 2-formamidoacrylate (5.35).** Methyl formate (30 mL) was added to **3.18** (1.516 g, 9.744 mmol) and potassium carbonate (5.403 g, 39.10 mmol) in a single-necked 100-mL round-bottomed flask. One drop of triethylamine was added and the mixture was stirred at rt overnight. After 28 h, the solid material was removed by filtration and washed with small portions of methyl formate (20 mL). The colorless filtrate was concentrated to yield a light yellow oil. The oil was chromatographed through a short column of silica (~35 g) using 50% hexanes/EtOAc and yielded **5.35** as a

colorless oil that solidified upon standing to a white solid (0.88 g, 70% yield). Isolated as a mixture of rotamers (88:12 *trans:cis*). mp = 55-57 °C (lit. 56.9-57.2 °C). Rf = 0.29 (1:1 hexanes/EtOAc – UV lamp). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  8.53 (m, 1H, *cis*), 8.39 (s, 1H, *trans*), 8.06 (br, 1H, *trans*), 7.76 (br, 1H, *cis*), 6.60 (s, 1H, *trans*), 5.92 (s, 1H, *trans*), 5.67 (s, 1H, *cis*), 5.42 (s, 1H, *cis*), 3.81 (s, 3H, *trans/cis*) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  164.1, 161.1, 159.6, 131.8, 130.1, 110.3, 104.8, 95.8, 52.9 ppm. HRMS (FAB) *m/z* [M+H]<sup>+</sup> calcd for C<sub>5</sub>H<sub>8</sub>NO<sub>3</sub><sup>+</sup>, 130.0504; obsd, 130.0527.



General procedure for the preparation of alkyl azides. Benzyl azide (5.36a). CAUTION: Azides may be explosive and shock sensitive and should be handled with care. A blast shield was used for every reaction that involved the use or preparation of azides. 75 mL of DMSO was added to sodium azide (2.45 g, 37.7 mmol) in a 250-mL single-necked round-bottomed flask under Ar. The mixture was stirred at rt until the sodium azide dissolved. Benzyl bromide (4.00 mL, 34.6 mmol) was added and the solution was stirred at rt under Ar for 4 h. 100 mL of H<sub>2</sub>O was added slowly to quench the reaction (exothermic), the resultant cloudy mixture was allowed to warm to rt (30 min), then poured into 100 mL of H<sub>2</sub>O. The mixture was extracted with Et<sub>2</sub>O (3 x 100 mL), and the combined Et<sub>2</sub>O layers were washed with brine, dried over MgSO<sub>4</sub>, filtered, and concentrated by rotary evaporation (30 °C, 21 mm Hg) and yielded **5.36a** as a colorless oil (4.604 g, 99.9% yield). Stored at 4 °C until ready to use. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.46 – 7.36 (m, 5H), 4.37 (s, 2H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  135.2, 128.7, 128.13, 128.06, 54.6 ppm. FT-IR (film) 3066, 3033, 2931, 2097 (N<sub>3</sub>), 1496, 1456, 1256, 1203, 1078, 1029 cm<sup>-1</sup>



**1-Azidooctane (5.36b).** Compound **5.36b** was prepared according to the general procedure for alkyl azides. 1-bromooctane (0.32 mL, 1.8 mmol) and sodium azide (0.112 g, 2.0 mmol) were stirred in DMSO at rt for 4 h and yielded **5.36b** as a pale yellow oil (0.251 g, 88% yield). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  3.25 (t, *J* = 6.9 Hz, 2H), 1.60 (m, 2H), 1.32 – 1.27 (m, 10H), 0.88 (t, *J* = 6.9 Hz, 3H) ppm. FT-IR (film) 2929 (CH), 2858 (CH), 2096 (N<sub>3</sub>), 1467 (CH bend), 1378 (CH bend), 1349, 1260 (N<sub>3</sub>) cm<sup>-1</sup>.



Azidocyclopentane (5.36c). Compound 5.36c was prepared according to the general procedure for alkyl azides. Bromocyclopentane (1.00 mL, 9.14 mmol) and sodium azide (0.658 g, 10.1 mmol) were stirred in DMSO for 25 h and yielded 5.36c as a pale yellow liquid (0.738 g, 73% yield). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  3.92 (m, 1H), 1.83 – 1.71 (m, 2H), 1.71 – 1.69 (m, 4H), 1.61 – 1.59 (m, 2H) ppm.



**4-Methylbenzenesulfonyl azide (5.37).** To a 500-mL round-bottomed flask cooled in an ice/ $H_2O$  bath was added *p*-toluenesulfonyl chloride (8.00 g, 42.0 mmol) and

240 mL of 1:1 acetone/H<sub>2</sub>O. The resultant suspension was stirred vigorously as sodium azide (2.767 g, 42.6 mmol) was added in one portion to the reaction. The cloudy suspension was stirred vigorously for 6 h, allowing the reaction to slowly reach rt as the ice/H<sub>2</sub>O bath melted. Most of the acetone was removed by rotary evaporation (~40°C, 21 mm Hg) to yield a light yellow oil beneath the aqueous layer. Et<sub>2</sub>O (~100 mL) was added and the layers were separated. The aqueous layer was extracted with Et<sub>2</sub>O (2 x 100 mL), and the combined ether layers were dried over MgSO<sub>4</sub>, filtered, and concentrated by rotary evaporation (~30°C, 21 mm Hg) and yielded **5.37** as a colorless liquid that solidified upon storage at -10 °C (7.59 g, 92% yield). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.84 (d, *J* = 8.4 Hz, 2H), 7.40 (d, *J* = 8.1 Hz, 2H), 2.48 (s, 3H) ppm. <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  146.2, 135.5, 130.3, 127.5, 21.8 ppm. FT-IR (film) 2360, 2128 (N<sub>3</sub>), 1596, 1372, 1121, 1168, 1086, 814, 748, 665 cm<sup>-1</sup>.

N<sub>3</sub>

**Phenyl azide (5.38a). Preparation using NaNO<sub>2</sub>/HCl.** 75 mL of H<sub>2</sub>O and concentrated HCl (14.0 mL, 169 mmol) were added to a 500-mL 3-necked round-bottomed flask equipped with a thermometer and a 10-mL addition funnel. The flask was cooled in a ice/NaCl/H<sub>2</sub>O bath to an internal temperature of about -5 °C. Phenylhydrazine (7.6 mL, 77.2 mmol) was added to the solution dropwise over 5 min. Phenylhydrazine-HCl precipitated out of the mixture as a fine white solid. 25 mL of Et<sub>2</sub>O was added and the mixture was stirred vigorously as a solution of NaNO<sub>2</sub> (6.254 g, 90.64 mmol) in 7.5 mL of H<sub>2</sub>O was added to the solution dropwise over about 25 min.

The white flakes of Phenylhydrazine-HCl had dissolved in the reaction and the mixture had turned a bright yellow color. After stirring for 10 min, the mixture was subjected to steam distillation until about 100 mL of distillate were collected. During the distillation, the mixture turned from yellow to brick red in color. An appreciable amount of tar was also formed. The orange ether layer of the distillate was removed and the aqueous layer was extracted with Et<sub>2</sub>O (2 x 20 mL). The combined Et<sub>2</sub>O layers were dried over MgSO<sub>4</sub> and filtered into a 100-mL round-bottomed flask. The Et<sub>2</sub>O was removed by rotary evaporation (RT, 21 torr) to yield an orange oil. The oil was purified by vacuum distillation and yielded **5.38a** as a yellow oil (1.58 g, 17% yield). bp = 26-27 °C (1.5 mm Hg). Isolated with impurities.

**Preparation of phenyl azide from aniline.** Aniline (0.50 mL, 5.48 mmol) was suspended in 4 mL of H<sub>2</sub>O and cooled in a crushed ice/H<sub>2</sub>O bath. Conc. H<sub>2</sub>SO<sub>4</sub> (1.20 mL, 21.7 mmol) was added dropwise and aniline hydrochloride precipitated out of the solution. To this mixture was added a solution of NaNO<sub>2</sub> (432 mg, 6.26 mmol) in 2.5 mL of H<sub>2</sub>O dropwise. A yellow color was observed in the mixture. Hexanes (8 mL) was added, followed by a solution of NaN<sub>3</sub> (383 mg, 5.89 mmol) in 2.5 mL of H<sub>2</sub>O dropwise to the biphasic mixture. A lot of bubbling was observed. The mixture was stirred vigorously in the ice/H<sub>2</sub>O bath over 3 h, allowing the ice/H<sub>2</sub>O bath to melt and reach rt. The layers were separated and the organic layer was dried over MgSO<sub>4</sub>, filtered, and concentrated by rotary evaporation (30 °C, 21 mm Hg) and yielded **5.38a** as a yellow oil (465 mg, 71% yield). A blast sheild was used when concentrating the azide as a precautionary measure. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 7.36 (m, 2H), 7.14 (m, 1H), 7.03

(m, 2H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 140.0, 129.7, 124.9, 119.0 ppm. FT-IR (film) 2130, 2096, 1594, 1493, 1296, 749 cm<sup>-1</sup>.



1-Azido-4-methoxybenzene (5.38b). Compound 5.38b was prepared following the same procedure for 5.38a from aniline. *p*-anisidine (0.662 g, 5.38 mmol), concentrated H<sub>2</sub>SO<sub>4</sub> (1.20 mL, 21.7 mmol), NaNO<sub>2</sub> (0.434 g, 6.29 mmol), and sodium azide (0.375 g, 5.77 mmol) yielded 5.38b as a brown oil (0.343 g, 44% yield). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 6.96 (m, 2H), 6.89 (m, 2H), 3.80 (s, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 156.9, 132.3, 119.9, 115.1, 55.5 ppm. FT-IR (film) 2955, 2837, 2105, 1505, 1465, 1286, 1246, 1182, 1035, 825, 626 cm<sup>-1</sup>.



General procedure for the preparation of imidazoles from 2-amidoacrylates and azides. Methyl 1-benzyl-2-phenyl-1H-imidazole-4-carboxylate (5.40a). Compound 5.23 (116.2 mg, 0.5663 mmol) and benzyl azide (0.11 mL, 0.87 mmol) were dissolved in 5 mL of toluene in a sealed tube fitted with a stir bar. The tube was heated in an oil bath maintained at 190 °C overnight. After 1 h, the colorless solution turned yellow in color and gradually the color became brown. After 24 h, the reaction was cooled to rt and concentrated. The crude brown oil was purified through 25 g of silica gel using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 80% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc and yielded pure **5.41a** (see below) (26.5 mg, 15.1% yield), pure **5.40a** as an amber oil (84.1 mg, 50.8% yield), and a mixture of **5.41a** and **5.40a** (26.1 mg). Total yield of **5.40a** isolated as an amber oil was 90.1 mg (54% yield). **5.40a** was recrystallized from EtOAc/hexanes to produce light yellow crystals suitable for x-ray diffraction. Rf = 0.14 (1:1 hexanes/EtOAc – UV lamp). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.69 (s, 1H), 7.60 (m, 2H), 7.44 – 7.40 (m, 3H), 7.37 – 7.32 (m, 3H), 7.09 (m, 2H), 5.23 (s, 2H), 3.90 (s, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  163.0, 148.8, 135.4, 132.5, 129.2, 129.0, 128.8, 128.7, 128.2, 127.9, 126.9, 126.5, 51.3, 50.5 ppm. HRMS (FAB) *m/z* [M+H]<sup>+</sup> calcd for C<sub>18</sub>H<sub>17</sub>N<sub>2</sub>O<sub>2</sub><sup>+</sup>, 293.1290; obsd, 293.1293.



(Z)-Methyl 2-benzamido-3-(benzylamino)acrylate (5.41a). Compound 5.41a was isolated as an amber oil. Total yield isolated was 46.6 mg (27% yield). 5.41a was recrystallized from 70% EtOH/H<sub>2</sub>O to yield colorless crystals suitable for x-ray diffraction. Rf = 0.26 (1:1 hexanes/EtOAc – UV lamp). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  8.00 (br, 1H), 7.84 (m, 2H), 7.51 (m, 1H), 7.44 (m, 2H), 7.37 – 7.26 (m, 6H), 6.50 (br, 1H), 4.42 (d, *J* = 6.0 Hz, 2H), 3.73 (s, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  166.5, 164.8, 140.1, 138.5, 134.0, 131.6, 131.3, 128.73, 128.69, 128.6, 128.51, 128.46, 127.2, 127.1, 127.0, 126.9, 98.6, 52.5, 51.6 ppm. HRMS (FAB) *m/z* [M]<sup>+</sup> calcd for C<sub>18</sub>H<sub>18</sub>N<sub>2</sub>O<sub>3</sub><sup>+</sup>, 310.1317; obsd, 310.1297.



Benzyl 1-benzyl-2-phenyl-1H-imidazole-4-carboxylate (5.40b). Compound 5.40b was prepared following the general procedure for the synthesis of imidazoles from acrylates and azides. Acrylate 5.24 (0.144 g, 0.513 mmol) and benzyl azide (0.10 mL, 0.79 mmol) were heated in toluene (5 mL) in a sealed tube for 24 h in a 190 °C oil bath. Chromatography through 25 g of silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 90% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc yielded pure 5.41b (see below), 5.40b, and a mixture of 5.40b and 5.41b as amber oils. Total yield of 5.40b was 120 mg (63% yield). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 7.65 (s, 1H), 7.56 (m, 2H), 7.44 (m, 2H), 7.42 – 7.29 (m, 9H), 7.05 (d, *J* = 7.0 Hz, 2H), 5.36 (s, 2H), 5.18 (s, 2H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 162.6, 149.2, 136.1, 135.6, 132.8, 129.4, 129.3, 129.1, 129.0, 128.44, 128.36, 128.3, 128.2, 128.0, 127.2, 126.7, 66.0, 50.7 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>24</sub>H<sub>21</sub>N<sub>2</sub>O<sub>2</sub><sup>+</sup>, 369.1603; obsd, 369.1602.



(Z)-Benzyl 2-benzamido-3-(benzylamino)acrylate (5.41b). Compound 5.41b was isolated as an amber oil (35 mg, 18% yield). 5.41b was isolated as an 8:1 mixture of isomers as evidenced by <sup>1</sup>H NMR data. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  9.04 (br, 1H), 7.84 (d, J = 7.5 Hz, 2H), 7.79 (d, J = 7.5 Hz, 2H, isomer), 7.51 (m, 1H), 7.45 – 7.26 (m, 13H), 6.59 (br, 1H), 5.21 (s, 2H), 4.44 (d, J = 6.0 Hz, 2H) ppm. <sup>13</sup>C NMR (125 MHz,

CDCl<sub>3</sub>)  $\delta$  164.9, 138.4, 136.5, 134.0, 131.6, 128.8, 128.7, 128.6, 128.53, 128.47, 128.02, 127.98, 127.5, 127.2, 127.1, 126.9, 66.1, 52.6 ppm. HRMS (FAB) m/z [M•]<sup>+</sup> calcd for C<sub>24</sub>H<sub>22</sub>N<sub>2</sub>O<sub>3</sub><sup>+</sup>, 386.1630; obsd, 386.1628.



**Benzyl 1-benzyl-2-(4-nitrophenyl)-1H-imidazole-4-carboxylate** (5.69b). Compound **5.69b** was prepared following the general procedure for the synthesis of imidazoles from acrylates and azides. Acrylate **5.28** (0.166 g, 0.510 mmol) and benzyl azide (0.117 g, 0.879 mmol) were heated in toluene. Chromatography through 20 g of silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 85% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc provided **5.70b** (see below), **5.69b** (123 mg), and a mixture of **5.69b** and **5.70b** as amber oils. Total yield of **5.69b** was 135 mg (64% yield). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  8.20 (m, 2H), 7.75 (m, 2H), 7.72 (s, 1H), 7.43 (m, 2H), 7.37 – 7.26 (m, 6H), 7.04 (m, 2H), 5.35 (s, 2H), 5.25 (s, 2H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  162.1, 147.9, 146.5, 135.8, 135.2, 134.8, 133.4, 129.6, 129.1, 128.44, 128.40, 128.3, 128.0, 126.3, 123.6, 66.2, 51.0 ppm. HRMS (FAB) *m*/*z* [M+H]<sup>+</sup> calcd for C<sub>24</sub>H<sub>20</sub>N<sub>3</sub>O<sub>4</sub><sup>+</sup>, 414.1454; obsd, 414.1460.



## (Z)-Benzyl 3-(benzylamino)-2-(4-nitrobenzamido)acrylate (5.70b).

Compound **5.70b** was isolated as an amber oil with impurities (18.1 mg, 8% yield). <sup>1</sup>H

NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  8.29 (m, 2H), 8.12 (br, 1H), 7.99 (m, 2H), 7.44 – 7.27 (m, 10H), 6.51 (br, 1H), 5.21 (s, 2H), 4.46 (d, *J* = 6.0 Hz, 2H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  162.5, 149.6, 139.5, 138.1, 136.3, 128.8, 128.6, 128.4, 128.2, 128.1, 127.7, 127.1, 123.9, 66.4, 52.8 ppm. HRMS (FAB) *m*/*z* [M•]<sup>+</sup> calcd for C<sub>24</sub>H<sub>21</sub>N<sub>3</sub>O<sub>5</sub><sup>+</sup>, 431.1481; obsd, 431.1465.



Methyl 1-benzyl-2-methyl-1H-imidazole-4-carboxylate (5.73). Compound 5.73 was prepared following the general procedure for the synthesis of imidazoles from acrylates and azides. Acrylate 5.34 (0.100 g, 0.701 mmol) and benzyl azide (0.122 g, 0.916 mmol) were heated in toluene (6 mL) in a sealed tube for 15 h in a 190 °C oil bath. Chromatography through 25 g of silica using a solvent gradient from 25% hexanes/EtOAc to 100% EtOAc provided 5.73 and 5.74. Byproduct 5.74 was isolated with a large amount of impurities. Imidazole 5.73 was isolated as an amber oil (120 mg, 74% yield). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.50 (s, 1H), 7.31 – 7.25 (m, 3H), 7.03 (d, *J* = 6.5 Hz, 2H), 5.02 (s, 2H), 3.80 (s, 3H), 2.31 (s, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  163.2, 146.1, 134.9, 131.4, 129.0, 128.2, 126.7, 126.1, 51.4, 50.1, 13.0 ppm. HRMS (FAB) *m*/z [M+H]<sup>+</sup> calcd for C<sub>13</sub>H<sub>15</sub>N<sub>2</sub>O<sub>2</sub><sup>+</sup>, 231.1134; obsd, 231.1139.



Methyl 1-benzyl-1H-imidazole-4-carboxylate (5.75). Compound 5.75 was prepared following the general procedure for the synthesis of imidazoles from acrylates and azides. Acrylate 5.35 (76.6 mg, 0.593 mmol) and benzyl azide (0.101 g, 0.756 mmol) were heated in toluene (6 mL) in a sealed tube in a 190 °C oil bath for 15 h. Chromatography through 25 g of silica using a solvent gradient from 50% hexanes/EtOAc to 33% hexanes/EtOAc provided 5.75 and 5.76. Byproduct 5.76 was isolated with a large amount of impurities. Imidazole 5.75 was isolated as an amber oil (90.9 mg, 71% yield). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.53 (d, *J* = 1.0 Hz, 1H), 7.50 (d, *J* = 1.0 Hz, 1H), 7.33 – 7.28 (m, 3H), 7.15 (m, 2H), 5.08 (s, 2H), 3.79 (s, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  163.0, 137.9, 134.8, 133.7, 129.0, 128.5, 127.4, 125.2, 51.4, 51.1 ppm. HRMS (FAB) *m*/*z* [M+H]<sup>+</sup> calcd for C<sub>12</sub>H<sub>13</sub>N<sub>2</sub>O<sub>2</sub><sup>+</sup>, 277.0977; obsd, 277.0983.



**3-Butyl-1-methyl-1H-imidazol-3-ium chloride, BMIM**<sup>+</sup>Cl<sup>-</sup> (5.67). Freshly distilled 1-methylimidazole (18.4 mL, 0.231 mol) was added to a 3-necked 250-mL round-bottomed flask fitted with a internal thermometer adapter, a condenser, under Ar. Dry acetonitrile (12.5 mL) was added, followed by 1-chlorobutane (31.5 mL, 0.299 mol). The colorless solution was heated in an oil bath (oil temp = 85-88 °C) to reflux (internal temperature = 75-80 °C). After 48 h, the yellow solution was cooled to rt, transferred to a single-necked 200-mL round-bottomed flask, and the volume was reduced by rotary

evaporation (55-60 °C, 21 mm Hg). The resultant thick yellow oil was stored under Ar at rt, then dried under vacuum (~1 mm Hg) for 5 h. The oil was dissolved in 32 mL of dry acetonitrile and added dropwise via cannula to a single-necked 500-mL round-bottomed flask containing EtOAc (125 mL) and one seed crystal of BMIM chloride (obtained from a previous preparation). The mixture is stirred vigorously during the addition and solid BMIM chloride crystallized immediately upon the addition of the acetonitrile solution. Once the addition was complete, the mixture was stirred vigorously under Ar and cooled in a -30 °C bath (dry ice/acetone) for 2 h. The mixture was quickly filtered while still cold and the white solid was washed with a minimal amount of EtOAc. (NOTE: The solid was very hygroscopic and started to turn into an oil very quickly). The solid was dried with P<sub>2</sub>O<sub>5</sub> under vacuum (1 mm Hg) for 24 h. Yielded 5.67 as a white solid material (35.5 g, 88% yield). lit. 89% yield. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  10.31 (s, 1H), 7.54 (t, J = 2 Hz, 1H), 7.37 (t, J = 2 Hz, 1H), 4.07 (s, 2H), 3.85 (s, 3H), 1.63 (m, 2H), 1.09 (septet, J = 7.5 Hz, 2H), 0.67 (t, J = 7.5 Hz, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 137.0, 123.3, 121.6, 49.1, 35.9, 31.6, 18.8, 12.8 ppm.



## 3-Butyl-1-methyl-1H-imidazol-3-ium tetrafluoroborate, BMIM<sup>+</sup>BF<sub>4</sub><sup>-</sup> (5.68). Compound 5.67 (30.46 g, 174.4 mmol) was dissolved in 35 mL of H<sub>2</sub>O in a 125-mL erlenmayer flask with a stir bar and thermometer. NaBF<sub>4</sub> (20.16 g, 183.6 mmol) was added to the solution in portions with stirring over 15 min. The NaBF<sub>4</sub> dissolved and the solution cooled to ~15 °C during the addition. The solution was stirred until reaching ambient temperature (~21 °C), then CH<sub>2</sub>Cl<sub>2</sub> (30 mL) was added. The layers were

separated and the aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (20 mL). The combined CH<sub>2</sub>Cl<sub>2</sub> layers were washed with a solution of NaBF<sub>4</sub> (10 g in 20 mL of H<sub>2</sub>O), dried over a mixture of Na<sub>2</sub>SO<sub>4</sub> (1 g) and MgSO<sub>4</sub> (3 g), filtered, and concentrated by rotary evaporation (45 °C, 20 mm Hg). The resultant oil was dried under vacuum (~1 mm Hg) at ambient temperature for 24 h to yield **5.68** as a thick, colorless/light yellow oil (36.2 g, 92% yield). lit. 89% yield. <sup>1</sup>H NMR (300 MHz, CD<sub>2</sub>Cl<sub>2</sub>)  $\delta$  8.70 (s, 1H), 7.28 (dt, *J* = 6.0, 1.8 Hz, 2H), 4.13 (t, *J* = 7.2 Hz, 2H), 3.89 (s, 3H), 1.81 (m, 2H), 1.31 (sextet, *J* = 7.5 Hz, 2H), 0.91 (t, *J* = 7.5 Hz, 3H) ppm.



Benzyl 1-(4-methoxyphenyl)-2-phenyl-1H-imidazole-4-carboxylate (5.81).

Acrylate **5.24** (162.1 mg, 0.576 mmol) and azide **5.38b** (169.3 mg, 1.14 mmol) were dissolved in 5 mL of nitromethane in a single-necked 25-mL round-bottomed flask fitted with a stir bar and condenser. The solution was heated to reflux in an oil bath (oil temperature = 115 °C). The reaction was monitored by TLC (1:1 hexanes/EtOAc and 3:2 hexanes/EtOAC - UV lamp). The reaction color changed to deep brown overnight, but was incomplete. The reaction was still incomplete after 2 and 3 days at reflux. After heating for 4 days, no starting material was observed by TLC and the solution was cooled to rt and concentrated. The brown oil was chromatographed through 25 g of silica using a gradual solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 85% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc and yielded the major product **5.81** as an orange oil (46.3 mg, 21% yield). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$ 

7.76 (s, 1H), 7.45 – 7.39 (m, 4H), 7.36 – 7.21 (m, 6H), 7.10 (m, 2H), 6.88 (m, 2H), 5.39 (s, 2H), 3.81 (s, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  162.8, 159.6, 147.9, 136.2, 132.8, 130.4, 129.2, 128.9, 128.61, 128.57, 128.44, 128.38, 128.1, 126.9, 114.7, 66.2, 55.5 ppm. HRMS (FAB) *m*/*z* [M+H]<sup>+</sup> calcd for C<sub>24</sub>H<sub>21</sub>N<sub>2</sub>O<sub>3</sub><sup>+</sup>, 385.1552; obsd, 385.1532.



*N*-Tosyliminoisodosobenzene (5.82). *p*-Toluenesulfonamide (3.405 g, 19.89 mmol) and KOH (3.30 g, 58.8 mmol) were dissolved in 30 mL of MeOH (exothermic). The solution was cooled in a crushed ice/H<sub>2</sub>O bath and became a white slurry that was hard to stir. About 10 mL of MeOH were added to the suspension. Iodobenzene diacetate (7.505 g, 23.30 mmol) was added to the vigorously stirred mixture in portions over about 5 min, maintaining an internal reaction temperature of less than 10 °C. The mixture turned yellow and most of the solid material dissolved. After about 5 minutes, white crystals were observed in the solution. The mixture was removed from the ice/H<sub>2</sub>O bath and stored in the refridgerator (about 5 °C) overnight. No solid was observed (except for the bit that did not dissolve) in the morning. The yellow solution was stirred at rt. After 30 min, yellow solid precipitated out of solution. The mixture was stirred for an additional 3 h at rt, then stored at 5 °C for 4 h. The solid was collected by filtration and **5.82** was obtained as an off-white solid (3.36 g, 45% yield). mp = 103-105 °C (dec.

violently). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.70 (m, 2H), 7.46 (m, 3H), 7.30 (m, 2H), 7.06 (m, 2H), 2.27 (s, 3H) ppm.

Attempted aziridination of acrylate 5.35 using iminoiodinane 5.82. Anhydrous CH<sub>3</sub>CN (10 mL) was added to a flame-dried single-necked 50-mL roundbottomed flask containing 5.35 (0.134 g, 1.04 mmol) and 5.82 (0.452 g, 1.21 mmol), under Ar. Cu(CH<sub>3</sub>CN)<sub>4</sub>PF<sub>6</sub> (21.9 mg, 0.0588 mmol) was added to the light yellow suspension. The color of the mixture changed immediately to a light lime-green color. After 2 min, most of the solid had dissolved and the mixture was a mint green color. The reaction flask was wrapped in foil and stirred at rt under Ar. After 2 h, the TLC of the reaction (1:1 hexanes/EtOAc - UV lamp) still showed mostly 5.35 and a two additional (faint) more nonpolar spots. The mixture was placed in an oil bath (oil temperature 55-60 °C) and stirred under Ar. After stirring overnight, no change was observed and the oil bath temperature was raised to ~90 °C. The reaction was heated to reflux overnight. No change by TLC. After heating for 4 days at reflux, no change was observed by TLC except for decomposition. Minor products were observed, but mostly starting material was left in the reaction.

Synthesis of imidazoles from oxazolines. Imidazole 5.40b from oxazoline 3.10. Compound 3.10 (143.2 mg, 0.509 mmol) and benzyl azide (118.4 mg, 0.889 mmol) were dissolved in 5 mL of toluene in a 25-mL round-bottomed flask. pTsOH (24.0 mg, 0.139 mmol) was added and the solution was heated in an oil bath to reflux (oil temp ~123 °C). The pTsOH did not seem to dissolve completely in the reaction. TLC analysis

of the reaction mixture (1:1 hexanes/EtOAc - UV lamp) indicated the reaction was complete after 4 days of reflux. The reaction was filtered, concentrated by rotary evaporation (40°C, 21 torr) and purified through 15 g of silica using a solvent gradient from 100%  $CH_2Cl_2$  to 95%  $CH_2Cl_2$ /EtOAc and yielded **5.40b** as a brown/orange oil (92 mg, 49% yield).

8.5 Experimental procedures for chapter 6

**General Procedure for the Synthesis of Triazolines Using Method A.** The alkene (1 mmol) and azide (1.5 mmol) were combined and stirred *neat* at 25 °C in a single-necked round-bottomed flask. The progress of the reaction was monitored by TLC or <sup>1</sup>H NMR for the disappearance of the alkene, and the crude material was purified through silica.

**General Procedure for the Synthesis of Triazolines Using Method B.** The alkene (1 mmol) and azide (1.5 mmol) were dissolved in 10 mL of CHCl<sub>3</sub> in a single-necked round-bottomed flask and stirred at 25 °C. The progress of the reaction was monitored by TLC or <sup>1</sup>H NMR for the disappearance of the alkene (typically 4 weeks), and the crude material was purified through silica.

General Procedure for the Synthesis of Triazolines Using Method C. The alkene (1 mmol) and azide (1.5 mmol) were dissolved in 10 mL of CHCl<sub>3</sub> in a single-

necked round-bottomed flask fitted with a condensor and heated to reflux in an oil bath (oil temp. = 80 °C). The progress of the reaction was monitored by TLC or <sup>1</sup>H NMR for the disappearance of the alkene (typically 3 days), and the crude material was purified through silica.

General Procedure for the Synthesis of Triazolines Using Method D. The alkene (1 mmol) and azide (1.5 mmol) were dissolved in 10 mL of PhCH<sub>3</sub> in a single-necked round-bottomed flask fitted with a condensor and heated to reflux in an oil bath (oil temp. =  $125 \,^{\circ}$ C). The progress of the reaction was monitored by TLC or <sup>1</sup>H NMR for the disappearance of the alkene (typically 3-4 h), and the crude material was purified through silica.



tert-Butyl  $(3a\alpha,4\beta,7\beta,7a\alpha)$ -1-benzyl-3a,4,7,7a-tetrahydro-4,7methano[1,2,3]triazole[4,5-d][1,2]oxazine-6(1*H*)-carboxylate (6.7a) and tert-butyl  $(3a\alpha,4\beta,7\beta,7a\alpha)$ -3-benzyl-3a,4,7,7a-tetrahydro-4,7-methano[1,2,3]triazole[4,5-

d][1,2]oxazine-6(3*H*)-carboxylate (6.8a). The title compounds were prepared following the general procedure for the synthesis of triazolines using Method A. Cycloadduct 3.38 (198 mg, 1.01 mmol) and benzyl azide (211 mg, 1.58 mmol) were reacted for 48 h. The brown crude material was purified through 15 g of silica using a solvent gradient of 100% CH<sub>2</sub>Cl<sub>2</sub> to 98% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to afford triazoline 6.8a (159 mg, 48% yield) as a white solid, then 85% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to afford triazoline **6.7a** (170 mg, 51% yield) as a white solid (99% total combined yield). Analytical and x-ray crystallographic samples of **6.7a** and **6.8a** were prepared by recrystallization from EtOAc/hexanes. **6.7a**: mp = 104-105 °C.  $\lambda$ max = 255 nm. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.37–7.27 (m, 5H), 4.92 (s, 1H), 4.90 (d, *J* = 14.5 Hz, 1H), 4.82 (d, *J* = 9.8 Hz, 1H), 4.72 (d, *J* = 14.5 Hz, 1H), 4.09 (s, 1H), 3.55 (d, *J* = 9.8 Hz, 1H), 1.70 (dt, *J* = 11.4, 1.5 Hz, 1H), 1.45 (d, *J* = 11.5 Hz, 1H), 1.39 (s, 9H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  156.4, 135.6, 128.8, 128.24, 128.17, 84.6, 82.6, 79.9, 61.2, 59.3, 53.4, 32.4, 27.9 ppm. HRMS (FAB) *m/z* [M+H]<sup>+</sup> calcd for C<sub>17</sub>H<sub>23</sub>N<sub>4</sub>O<sub>3</sub><sup>+</sup>, 331.1770; obsd, 331.1745. **6.8a**: mp = 100-101 °C.  $\lambda$ max = 255 nm. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.35-7.24 (m, 5H), 4.92 (d, *J* = 14.7 Hz, 1H), 4.85 (s, 1H), 4.84 (d, *J* = 9.9 Hz, 1H), 1.45-1.43 (m, 10H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  156.4, 135.6, 128.8, 128.3, 128.2, 83.7, 82.7, 79.6, 61.5, 59.9, 53.6, 32.3, 27.9 ppm. HRMS (FAB) *m/z* [M+H]<sup>+</sup> calcd for C<sub>17</sub>H<sub>23</sub>N<sub>4</sub>O<sub>3</sub><sup>+</sup>, 331.1770; obsd, 331.1770; obsd, 331.1770; obsd, 331.1770; obsd, 331.350 (d, *J* = 9.9 Hz, 1H), 1.68 (d, *J* = 11.5 Hz, 1H), 1.45-1.43 (m, 10H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  156.4, 135.6, 128.8, 128.3, 128.2, 83.7, 82.7, 79.6, 61.5, 59.9, 53.6, 32.3, 27.9 ppm. HRMS (FAB) *m/z* [M+H]<sup>+</sup> calcd for C<sub>17</sub>H<sub>23</sub>N<sub>4</sub>O<sub>3</sub><sup>+</sup>, 331.1770; obsd, 331.1770; obsd, 331.1753.



*tert*-Butyl  $(3a\alpha,4\beta,7\beta,7a\alpha)$ -1-(1-adamantyl)-3a,4,7,7a-tetrahydro-4,7methano[1,2,3]triazole[4,5-d][1,2]oxazine-6(1*H*)-carboxylate (6.7b) and *tert*-butyl  $(3a\alpha,4\beta,7\beta,7a\alpha)$ -3-(1-adamantyl)-3a,4,7,7a-tetrahydro-4,7-

methano[1,2,3]triazole[4,5-d][1,2]oxazine-6(3H)-carboxylate (6.8b). The title

compounds were prepared following the general procedure for the synthesis of triazolines using Method B. Cycloadduct 3.38 (104 mg, 0.527 mmol) and 1-azidoadamantane (140 mg, 0.79 mmol) were reacted for 4 weeks. The crude material was purified through 20 g of silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 97% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to afford 6.8b (86 mg, 43% yield) as a white solid, then 85% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to afford 6.7b (102 mg, 52% yield) as a white solid (95% total combined yield). **6.7b:** mp > 144  $^{\circ}$ C (dec.). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  4.93 (s, 1H), 4.77 (dt, J = 9.5, 1.5 Hz, 1H), 4.42 (s, 1H), 3.85 (d, J = 9.5 Hz, 1H), 2.14-2.07 (m, 6H), 1.84-1.81 (m, 4H), 1.72-1.65 (m, 6H), 1.47(m, 10H) ppm.  ${}^{13}$ C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  156.5, 83.2, 82.7, 80.4, 63.1, 57.2, 41.7, 36.0, 32.4, 29.2, 28.1 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>20</sub>H<sub>31</sub>N<sub>4</sub>O<sub>3</sub><sup>+</sup>, 375.2396; obsd, 375.2375. **6.8b:** mp > 166 °C (dec.). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  4.85 (m, 1H), 4.76 (dt, J = 10.0, 1.4 Hz, 1H), 4.49 (t, J = 1.4 Hz, 1H), 3.80 (dt, J = 10.0, 1.2 Hz, 1H), 2.12 (m, 2H), 2.03 (ddd, J = 11.6, 4.7, 3.0 Hz, 4H), 1.77 (m, 4H), 1.70-1.62 (m, 6H), 1.47 (m, 10H) ppm.  $^{13}$ C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  156.5, 82.782.3, 81.5, 61., 57.0, 56.7, 41.7, 36.0, 32.3, 29.2, 28.0 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>20</sub>H<sub>31</sub>N<sub>4</sub>O<sub>3</sub><sup>+</sup>, 375.2396; obsd, 375.2372.



*tert*-Butyl (3aα,4β,7β,7aα)-1-(*n*-octyl)-3a,4,7,7a-tetrahydro-4,7-



## $(3a\alpha, 4\beta, 7\beta, 7a\alpha)$ -3-(n-octyl)-3a, 4, 7, 7a-tetrahydro-4, 7-methano[1, 2, 3]triazole[4, 5-

d][1,2]oxazine-6(3H)-carboxylate (6.8c). The title compounds were prepared following the general procedure for the synthesis of triazolines using Method C. Cycloadduct 3.38 (199 mg, 1.01 mmol) and n-octyl azide (237 mg, 1.53 mmol) were reacted for 2 days. The crude material was purified through 40 g of silica using 80% hexanes/EtOAc to afford 6.8c (124 mg, 35% yield), mixed 6.7c/6.8c (192 mg, 54%), and 6.7c (38 mg, 11% yield) as off-white semi-solids (99% total combined yield). 6.7c: mp = 58-59 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  4.97 (s, 1H), 4.87 (d, J = 9.5 Hz, 1H), 4.55 (s, 1H), 3.71-3.66 (m, 2H), 3.58-3.52 (m, 1H), 1.80 (dt, J = 11.5, 1.5 Hz, 1H), 1.70-1.64 (m, 2H), 1.50 (m, 10H), 1.33-1.24 (m, 12H), 0.88 (t, J = 7.0 Hz, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$ 156.9, 84.0, 83.0, 80.2, 61.6, 60.1, 49.3, 32.4, 31.7, 29.7, 29.1, 28.8, 28.1, 26.7, 22.6, 14.1 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>18</sub>H<sub>33</sub>N<sub>4</sub>O<sub>3</sub><sup>+</sup>, 353.2553; obsd, 353.2525. **6.8c:** mp = 49-50 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  4.87 (m, 1H), 4.58 (s, 1H), 3.66-3.61 (m, 2H), 3.52-3.47 (m, 1H), 1.80 (dt, J = 11.5, 1.5 Hz, 1H), 1.61-1.58 (m, 2H), 1.48 (m, 10H), 1.30-1.24 (m, 12H), 0.86 (t, J = 7.0 Hz, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl-3) δ 156.6, 83.0, 82.9, 80.0, 61.7, 60.1, 49.3, 32.4, 31.7, 29.0, 28.8, 28.1, 26.6, 22.6, 14.0 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>18</sub>H<sub>33</sub>N<sub>4</sub>O<sub>3</sub><sup>+</sup>, 353.2553; obsd, 353.2525.



*tert*-Butyl (3aα,4β,7β,7aα)-1-cyclopentyl-3a,4,7,7a-tetrahydro-4,7methano[1,2,3]triazole[4,5-d][1,2]oxazine-6(1H)-carboxylate (6.7d) and tert-butyl (3aα,4β,7β,7aα)-3-cyclopentyl-3a,4,7,7a-tetrahydro-4,7-methano[1,2,3]triazole[4,5**d**][1,2]oxazine-6(3H)-carboxylate (6.8d). The title compounds were prepared following the general procedure for the synthesis of triazolines using Method C. Cycloadduct 3.38 (201 mg, 1.02 mmol) and 1-azidocyclopentane (155 mg, 1.39 mmol) were reacted for 2 days. The crude material was purified through 30 g of silica using 80% hexanes/EtOAc to afford 6.8d (88 mg, 28% yield) as a white solid, mixed 6.7d/6.8d (194 mg, 62%) as an off-white solid, and 6.7d (25 mg, 8% yield) as a colorless oil (97% total combined yield). **6.7d:** mp >70 °C (dec.). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  4.96 (s, 1H), 4.85 (d, J = 9.5 Hz, 1H), 4.52 (s, 1H), 4.05 (p, J = 7.0 Hz, 1H), 3.70 (d, J = 9.5 Hz, 1H), 2.08-1.92 (m, 2H), 1.83-1.62 (m, 8H), 1.49 (m, 10H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 156.7, 83.7, 82.9, 80.3, 62.1, 60.9, 59.4, 32.5, 31.6, 31.1, 28.1, 23.6, 23.4 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for  $C_{15}H_{25}N_4O_3^+$ , 309.1927; obsd, 309.1913. **6.8d:** mp > 120 °C (dec.). <sup>1</sup>H NMR  $(500 \text{ MHz}, \text{CDCl}_3) \delta 4.84 \text{ (s, 1H)}, 4.81 \text{ (d, } J = 9.5 \text{ Hz}, 1\text{H}), 4.55 \text{ (s, 1H)}, 3.97 \text{ (p, } J = 7.0 \text{ (s, 1H)})$ Hz, 1H), 3.61 (d, J = 9.5 Hz, 1H), 2.02-1.95 (m, 1H), 1.87-1.83 (m, 1H), 1.78-1.57 (m, 7H), 1.45-1.42 (m, 10H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 156.5, 82.8, 82.7, 80.4, 61.6, 60.8, 59.8, 32.3, 31.5, 31.1, 28.0, 23.5, 23.3 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>15</sub>H<sub>25</sub>N<sub>4</sub>O<sub>3</sub><sup>+</sup>, 309.1927; obsd, 309.1913.



tert-Butyl  $(3a\alpha,4\beta,7\beta,7a\alpha)$ -1-phenyl-3a,4,7,7a-tetrahydro-4,7methano[1,2,3]triazole[4,5-d][1,2]oxazine-6(1*H*)-carboxylate (6.7e) and tert-butyl  $(3a\alpha,4\beta,7\beta,7a\alpha)$ -3-phenyl-3a,4,7,7a-tetrahydro-4,7-methano[1,2,3]triazole[4,5-

d][1,2]oxazine-6(3*H*)-carboxylate (6.8e). The title compounds were prepared following the general procedure for the synthesis of triazolines using Method D. Cycloadduct **3.38** (202 mg, 1.02 mmol) and phenyl azide (165 mg, 1.39 mmol) were reacted for 3 h. The crude material was purified through 35 g of silica using 100% CH<sub>2</sub>Cl<sub>2</sub> to afford **6.8e** (175 mg, 54% yield) as a light yellow solid, then 80% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to afford **6.7e** (144 mg, 45% yield) as a light yellow oil that solidified upon standing (99% total combined yield). **6.7e:** mp = 117-120 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.41-7.35 (m, 4H), 7.10 (t, *J* = 7.0 Hz, 1H), 5.09 (d, *J* = 10.0 Hz, 1H), 5.085 (s, 1H), 4.86 (s, 1H), 4.46 (d, *J* = 10.0 Hz, 1H), 1.91 (dt, *J* = 12.0, 2.0 Hz, 1H), 1.54 (m, 10H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  156.4, 139.3, 129.6, 123.0, 113.8, 83.9, 83.2, 79.7, 60.2, 57.1, 32.2, 28.0 ppm. HRMS (FAB) *m*/*z* [M+H]<sup>+</sup> calcd for C<sub>16</sub>H<sub>21</sub>N<sub>4</sub>O<sub>3</sub><sup>+</sup>, 317.1614; obsd, 317.1622. **6.8e:** mp > 158 °C (dec.). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.37 (t, *J* = 8.0 Hz, 3H), 7.30 (d, *J* = 8.5 Hz, 2H), 7.08 (t, *J* = 7.5 Hz, 1H), 5.10 (d, *J* = 9.5 Hz, 1H), 5.02 (s, 1H), 4.92 (s, 1H), 4.24 (d, *J* = 9.5 Hz, 1H), 1.86 (dt, *J* = 12.0, 2.0 Hz, 1H), 1.54 (m, 10H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  156.6, 139.4, 129.7, 123.1, 113.9, 83.2, 78.7, 61.4, 58.0, 32.3, 28.1 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>16</sub>H<sub>21</sub>N<sub>4</sub>O<sub>3</sub><sup>+</sup>, 317.1614; obsd, 317.1605.

Attempted formation of aziridine 6.9 directly from cycloadduct 3.38. Compound 3.38 (197.0 mg, 0.9988 mmol) was dissolved in 10 mL of toluene in a 50-mL round-bottomed flask. Azidotrimethylsilane (0.20 mL, 1.5 mmol) was added and the reaction was heated to reflux (oil bath temp ~125°C) for 4 h. The reaction was concentrated by rotary evaporation (~20 torr, ~40°C) to yield an amber oil. No triazoline or aziridine products were observed by TLC or NMR.

Second reaction attempt without heating: A 1-dram screw-cap vial was charged with **3.38** (21 mg, 0.10 mmol) and dissolved in 1 mL of CDCl<sub>3</sub>. Azidotrimethylsilane (21 uL, 0.16 mmol) was added and the solution was stirred at rt. The reaction was monitored by <sup>1</sup>H NMR after 3, 7, 14, 22, and 28 days. No reaction was observed.



(±)-tert-Butyl (1 $\alpha$ ,2 $\beta$ ,4 $\beta$ ,5 $\alpha$ )-3-[(4-methylphenyl)sulfonyl]-6-oxa-3,7diazatricyclo[3.2.1.0<sup>2,4</sup>]octane-7-carboxylate (6.10). Cycloadduct 3.38 (205 mg, 1.04 mmol) and tosyl azide (268 mg, 1.36 mmol) were dissolved in 10 mL of PhCH<sub>3</sub> in a 25mL single-necked round-bottomed flask fitted with a condenser and heated to reflux in an oil bath (oil temperature = 125 °C). The reaction was monitored by TLC (1:1 hexanes/EtOAc; UV lamp) for the disappearance of 3.38. After 9 h, the reaction was complete and the solution was concentrated to yield a brown oil. The oil was purified through 35 g of silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 90% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to yield **6.10** as a tan solid (263 mg, 69% yield). mp = 111-115 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.74 (d, *J* = 8.0 Hz, 2H), 7.31 (d, *J* = 8.5 Hz, 2H), 4.78 (s, 1H), 4.62 (s, 1H), 3.26 (d, *J* = 6.0 Hz, 1H), 3.22 (d, *J* = 6.0 Hz, 1H), 2.40 (s, 3H), 2.01 (d, *J* = 11.0 Hz. 1H), 1.44 (s, 10H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  156.6, 145.0, 133.9, 129.7, 127.8, 82.9, 78.1, 59.4, 36.5, 36.2, 29.0, 27.9, 21.5 ppm. MS (FAB) *m*/*z* [M+H]<sup>+</sup> at 367, 311, 267 (100%). HRMS (FAB) *m*/*z* [M+H]<sup>+</sup> calcd for C<sub>17</sub>H<sub>23</sub>N<sub>2</sub>O<sub>5</sub>S<sup>+</sup>, 367.1328; obsd, 367.1314.



Reaction of 3.39 With Benzyl Azide: Formation of triazolines 6.11, 6.12, 6.13,

and 6.14. Compound 3.39 (421 mg, 1.99 mmol) and benzyl azide (1.33 g, 9.97 mmol) were dissolved in 10 mL of toluene in a 25-mL single-necked round-bottomed flask fitted with a condenser and heated to reflux in an oil bath (oil temp. = 125 °C). The reaction was monitored by TLC (1:1 hexanes/EtOAc; UV lamp) for the disappearance of 3.39. After 28 h, the deep brown solution was concentrated to yield a brown oil. The oil was chromatographed through silica using a solvent gradient from 100% hexanes to 50% hexanes/EtOAc to afford 6.12 (136 mg, 20% yield), 6.11 (128 mg, 19% yield), and an inseparable mixture of 6.13 and 6.14 (163 mg, 24% yield), all as brown solids (62% total combined yield). 6.11: <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.38-7.30 (m, 5H), 4.99 (ddd, *J* =

13.0, 4.5, 1.5 Hz, 1H), 4.85 (d, J = 14.5 Hz, 1H), 4.77 (d, J = 14.5 Hz, 1H), 4.54 (t, J = 4.0 Hz, 3.83 (br-s, 1H), 3.63 (dd, J = 13.0, 4.0 Hz, 1H), 1.98-1.90 (m, 1H), 1.76-1.70 (m, 1H), 1.63 (tdd, J = 13.8, 4.2, 2.5 Hz, 1H), 1.52-1.46 (m, 1H), 1.43 (m, 10H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  155.4, 132.9, 128.8, 128.5, 128.2, 81.9, 78.1, 69.5, 53.5, 47.2, 28.1, 19.6, 18.5 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>18</sub>H<sub>25</sub>N<sub>4</sub>O<sub>3</sub><sup>+</sup>, 345.1927; obsd, 345.1931. **6.12:** <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.32-7.21 (m, 5H), 4.90 (ddd, J = 12.5, 4.0, 1.0 Hz, 1H), 4.75 (d, J = 14.5 Hz, 1H), 4.66 (d, J = 14.5 Hz, 1H), 4.45 (m, 1H), 3.71 (m, 1H), 3.60 (dd, J = 12.5, 4.5 Hz, 1H), 1.86-1.80 (m, 1H), 1.77-1.70 (m, 1H), 1.55-1.49 (m, 1H), 1.47-1.43 (m, 1H), 1.42 (m, 10H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  155.7, 134.9, 128.9, 128.7, 128.4, 82.1, 78.4, 70.3, 54.2, 54.1, 47.6, 47.6, 28.2, 20.2, 17.9 ppm; HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>18</sub>H<sub>25</sub>N<sub>4</sub>O<sub>3</sub><sup>+</sup>, 345.1931. **6.13** and **6.14** were obtained as ~1:1 mixture as evidenced by <sup>1</sup>H NMR. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>18</sub>H<sub>25</sub>N<sub>4</sub>O<sub>3</sub><sup>+</sup>, 345.1931.

General procedure for the addition of benzyl azide to other alkenes (3.73, 3.74, 3.77, 3.40, and 6.15). Benzyl azide (1.5 equivalents or greater) was heated with the alkene (1 equivalent) in toluene or neat. Reactions were monitored by <sup>1</sup>H NMR and TLC for consumption of the alkene starting material. When alkene 3.73 was heated with 1.5 eq. of BnN<sub>3</sub> in toluene at reflux for 2 days, a complex mixture of triazoline products were observed by <sup>1</sup>H NMR; however, the reaction did not progress to completion. All other alkenes examined resulted in no reaction.

General procedure for investigation of the effect of Ru(II) catalyst 6.16 on azide [3+2] cycloaddition reactions. Cycloadduct 3.38 or 3.40 (1 mmol) and benzyl azide (1.2 – 1.5 mmol) were dissolved in 10 mL of benzene in a 50-mL round-bottomed flask equipped with a stir bar and condenser. Catalyst 6.16 (0.05 mmol) was added to the reaction and the mixture was stirred at rt. Control reactions were performed as described above without the addition of catalyst 6.16. The reaction was monitored by TLC for the disappearance of cycloadduct 3.38 or 3.40. After 1 h, no reaction was observed and the reaction was stirred in a 55 °C oil bath for 3 days. The reaction was incomplete at this time. The mixture was heated to reflux (oil bath temperature = 85 °C) for 6 h and the mixture was concentrated. The residue was analyzed by TLC and <sup>1</sup>H NMR. % conversion was determined by <sup>1</sup>H NMR.



General Procedure for Photolysis of Triazolines. ( $\pm$ )-*tert*-Butyl (1 $\alpha$ ,2 $\beta$ ,4 $\beta$ ,5 $\alpha$ )-3-benzyl-6-oxa-3,7-diazatricyclo[3.2.1.0<sup>2,4</sup>]octane-7-carboxylate (6.17). Compound 6.7a (333 mg, 1.01 mmol) was dissolved in 300 mL of degassed CH<sub>3</sub>CN and transferred to a 450-mL photochemical reaction vessel. The solution was irradiated in an immersion-well reactor under a stream of Ar with a Hanovia 450W mercury lamp equipped with a Vycor filter sleeve. Reaction progress was monitored by TLC and <sup>1</sup>H NMR for the disappearance of 6.7a. After 3 h, the reaction was concentrated and the crude material was purified through 30 g of silica using a gradient consisting of 100% CH<sub>2</sub>Cl<sub>2</sub> to 95% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc to afford 6.17 as a colorless oil (205 mg, 67% yield). <sup>1</sup>H

NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.35-7.27 (m, 5H), 4.79 (m, 1H), 4.62 (m, 1H), 3.40 (d, J = 13.5 Hz, 1H), 3.35 (d, J = 14.0 Hz, 1H), 2.29-2.22 (m, 3H), 1.50 (s, 9H), 1.39 (d, J = 10.5 Hz, 1H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  157.5, 138.6, 128.4, 127.7, 127.3, 82.3, 79.9, 60.7, 59.1, 37.1, 36.7, 29.4, 28.2 ppm. MS (FAB) m/z [M+H]<sup>+</sup> at 303, 247, 203, 171 (100%). HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>17</sub>H<sub>23</sub>N<sub>2</sub>O<sub>3</sub><sup>+</sup>, 303.1709; obsd, 303.1712.

Attempted photolytic conversion of triazoline 6.7a to aziridine 6.17 using TPP as a sensitizer. Compound 6.7a (100.6 mg, 0.305 mmol) and tetraphenylporphyrin (TPP, 1.9 mg, 0.0031 mmol) were dissolved in 10 mL of anhydrous  $CH_3CN$  and 2 mL of  $CH_2Cl_2$  (to help solubilize the TPP). The solution was degassed with Ar for 5 min, then irradiated using a 250-W sunlamp while bubbling the solution with Ar. After 30 min, the heat of the sunlamp had melted the ice bath, part of the outlet needle, and the upper portion of the rubber septum (it made a big mess...). Only the triazoline 6.7a and TPP were observed when the reaction mixtre (now a darker purple color) was analyzed by TLC (1:1 hexanes/EtOAc - UV lamp, CAM stain). None of the desired product was observed, and no obvious decomposition of starting materials was observed either.

Attempted conversion of triazoline 6.7a to aziridine 6.17 with pyridine. Compound 6.7a (101 mg, 0.306 mmol) was dissolved in 3 mL of toluene in a 25-mL round-bottomed flask. Pyridine (0.030 mL, 0.37 mmol) was added and the solution was heated to reflux (oil bath temp ~125 °C). After ~5 h, TLC of the reaction did not indicate the presence of aziridine 6.17, and only showed starting material 6.7a (1:1 hexanes/EtOAc - UV lamp). The reaction was heated at reflux overnight. Even when heated neat overnight, no reaction was observed.

Attempted conversion of triazoline 6.7a to aziridine 6.17 using TMSOTf. Compound 6.7a (80.2 mg, 0.243 mmol) was dissolved in 2 mL of  $CHCl_3$  in a 20-mL scintillation vial under Ar. TMSOTf (0.010 mL, 0.052 mmol) was added and the solution was stirred under Ar at RT. After 1 h, the initially yellow solution appeared slightly darker, however TLC of the reaction (1:1 hexanes/EtOAc - UV lamp) only showed triazoline 6.7a plus a minor amount of decomposition.



General procedure for N-O bond reduction of triazolines using  $Mo(CO)_6/NaBH_4$ . (±)-*tert*-Butyl ( $3a\alpha,4\alpha,6\alpha,6a\alpha$ )-3-benzyl-6-hydroxy-3,3a,4,5,6,6a-hexahydrocyclopenta[d][1,2,3]triazol-4-ylcarbamate (6.19). Triazoline 6.7a (207 mg, 0.628 mmol) was dissolved in 5 mL of 4:1 CH<sub>3</sub>CN/H<sub>2</sub>O in a 25-mL single-necked round-bottomed flask and heated in a 50 °C oil bath. Molybdenum hexacarbonyl (72 mg, 0.27 mmol) was added to the solution in one portion, followed by sodium borohydride (80 mg, 2.1 mmol) added in portions. Bubbling was observed and the color of the reaction changed from light yellow to a deep, murky brown. After the bubbling subsided, the reaction was heated to reflux (oil temperature = 70 °C) and monitored by TLC (1:1 hexanes/EtOAc; UV lamp) for the disappearance of 6.7a. After 18 h, the reaction was cooled in a crushed ice/H<sub>2</sub>O bath and the solid material was removed by filtration through

a pad of celite. The celite was washed with EtOAc (20 mL) and the filtrate was concentrated to yield an off-white solid, which was purified through 20 g of silica using 50% hexanes/EtOAc to afford **6.19** as a white solid (123 mg, 59% yield). mp = 153-155 °C (dec.). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.34-7.28 (m, 5H), 5.60 (d, *J* = 9.0 Hz, 1H), 5.15 (d, *J* = 15.0 Hz, 1H), 4.78 (d, *J* = 10.8 Hz, 1H), 4.63 (d, *J* = 15.0 Hz, 1H), 4.51 (d, *J* = 4.2 Hz, 1H), 4.11 (m, 1H), 3.66 (d, *J* = 10.8 Hz, 1H), 1.76 (d, *J* = 14.4 Hz, 1H), 1.55-1.49 (m, 1H), 1.43 (s, 9H) ppm. HRMS (FAB) *m*/*z* [M+H]<sup>+</sup> calcd for C<sub>17</sub>H<sub>25</sub>N<sub>4</sub>O<sub>3</sub><sup>+</sup>, 333.1927; obsd, 333.1906.





hexahydrocyclopenta[d][1,2,3]triazol-4-ylcarbamate (6.20). Compound 6.20 was prepared following the same procedure for 6.19. 6.8a (506 mg, 1.53 mmol), molybdenum hexacarbonyl (181 mg, 0.686 mmol) and NaBH<sub>4</sub> (220 mg, 5.8 mmol) provided an amber foam (540 mg). Chromatography through 10 g of silica using 83% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc yielded 6.20 as a light yellow solid (295 mg, 58% yield). mp > 110 °C (dec.). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.30-7.20 (m, 5H), 5.63 (br-m, 1H), 4.91-4.85 (m, 2H), 4.63 (d, *J* = 15.0 Hz, 1H), 4.21 (t, *J* = 8.0 Hz, 1H), 4.07 (d, *J* = 4.0 Hz, 1H), 3.64 (d, *J* = 10.5 Hz, 1H), 1.69 (d, *J* = 14.0 Hz, 1H), 1.53 (br-s, 1H), 1.39 (s, 10H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  155.1, 136.0, 128.7, 128.2, 127.9, 89.4, 79.6, 76.0, 66.9, 56.7, 52.9, 37.8, 28.4 ppm. HRMS (FAB) *m*/*z* [M+H]<sup>+</sup> calcd for C<sub>17</sub>H<sub>25</sub>N<sub>4</sub>O<sub>3</sub><sup>+</sup>, 333.1927; obsd, 333.1930.



(±)-*tert*-**Butyl** (1α,2α,4α,5α)-6-benzyl-4-hydroxy-6-aza-bicyclo[3.1.0]hexan-2ylcarbamate (6.21). Triazoline 6.19 (108 mg, 0.323 mmol) was dissolved in 300 mL of degassed CH<sub>3</sub>CN in a 450-mL photochemical reaction reaction vessel. The solution was irradiated in an immersion-well reactor under a stream of Ar with a Hanovia 450W mercury lamp fitted with a Vycor filter sleeve. The progress of the reaction was monitored by <sup>1</sup>H NMR. After 5 h, the solution was concentrated (40 °C, 21 torr) to yield an amber oil (110 mg). The oil was chromatographed through a pad of silica using a solvent gradient from 100% CH<sub>2</sub>Cl<sub>2</sub> to 40% CH<sub>2</sub>Cl<sub>2</sub>/EtOAc and yielded **6.21** as a yellow oil (89.9 mg, 91% yield). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 7.32-7.23 (m, 5H), 5.16 (m, 1H), 4.26 (d, *J* = 5.0 Hz, 1H), 4.08 (t, *J* = 7.5 Hz, 1H), 3.52 (d, *J* = 14 Hz, 1H), 3.32 (d, *J* = 14 Hz, 1H), 2.30 (m, 2H), 2.05 (m, 1H), 1.47 (d, *J* = 15 Hz, 1H), 1.43 (s, 9H), 1.25 (m, 1H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 155.4, 138.7, 128.3, 127.3, 126.9, 71.8, 60.7, 51.0, 48.4, 47.6, 39.8, 29.6, 28.3 ppm. HRMS (FAB) *m*/*z* [M]<sup>+</sup> calcd for C<sub>17</sub>H<sub>24</sub>N<sub>2</sub>O<sub>3</sub><sup>+</sup>, 304.1787; obsd, 304.1785. 8.6 Experimental procedures for chapter 7

$$\overset{N^{-}NH_{2}}{\overset{||}{\overset{||}{\overset{||}{\overset{}}}}} Ph$$

(**Diphenylmethylene**)**hydrazine** (**7.12**). Benzophenone (10.099 g, 55.323 mmol) was dissolved in 60 mL of absolute EtOH in a 250-mL round-bottomed flask. The solution was heated in a 50 °C oil bath and hydrazine monohydrate (4.10 mL, 82.8 mmol) was added. A condenser was attached to the flask and the colorless solution was heated to reflux (oil bath temp = 100 °C) overnight. After 16 h, the reaction was cooled to rt and the colorless solution was concentrated to yield a white solid. The solid was recrystallized from absolute EtOH to yield **7.12** as white needles (6.17 g, 57%). mp = 96-98 °C (lit.<sup>2</sup> 97-98 °C).



**Diphenyldiazomethane (7.13). 7.12** (2.011 g, 10.25 mmol) was dissolved in 20 mL of anhydrous  $CH_2Cl_2$  in a 100-mL round-bottomed flask and cooled in an ice/H<sub>2</sub>O bath under Ar. Anhydrous MgSO<sub>4</sub> (3.21 g, 26.7 mmol) was added and the mixture was stirred vigorously. A blast shield was placed in front of the reaction as a precautionary measure as activated MnO<sub>2</sub> (88%, 1.187 g, 12.02 mmol) was added in one portion to the reaction. The reddish-purple suspension was stirred in the ice/H<sub>2</sub>O bath for 2 h, then warmed to rt and stirred an additional 1 h. The solid material was removed by vacuum

filtration and washed with  $CH_2Cl_2$ . The filtrate was concentrated by rotary evaporation (~20 torr, ~30°C) to yield crude **7.13** as a deep purple-colored solid (2.02 g). The solid was stored in the freezer (-10°C) under Ar. The crude mixture of **7.13** was used without purification in further reactions.

**Treatment of cycloadduct 3.38 with diphenyldiazomethane.** Compound **3.38** (207.8 mg, 1.054 mmol) and crude **7.13** (388 mg, 2.00 mmol) were dissolved in 5 mL of benzene in a 25-mL round-bottomed flask under Ar. The purple solution was purged with Ar and stirred at rt in the dark (covered in foil). The reaction was monitored by TLC (1:1 and 2:1 hexanes/EtOAc - UV lamp), and after 3 days the reaction appeared to be progressing; however, there was still a lot of starting material (**3.38**) in the reaction. After 10 days, the reaction was concentrated and a crude <sup>1</sup>H NMR of the residue indicated that the reaction was progressing. After stirring for another week, **3.38** was completely consumed.



(±)-(3aR,6aS)-3-Hydroxy-3a,4-dihydro-3*H*-cyclopenta[*d*]oxazol-2(6a*H*)-one (7.34). Initial preparation using un-purified dimethyl sulfate. Compound 3.38 (100.2 mg, 0.5080 mmol) was dissolved in 5 mL of anhydrous  $CH_2Cl_2$  in a flame-dried singlenecked 25-mL round-bottomed flask under Ar. Dimethyl sulfate (0.10 mL, 1.1 mmol, not distilled prior to use) was added and bubbling was observed. After 5 min, the bubbling had ceased and the reaction was analyzed by TLC (1:1 hexanes/EtOAc - UV

lamp, CAM stain). After 4 h, no cycloadduct was observed and anhydrous methanol (0.20 mL, 4.9 mmol) was added to the reaction. After 1 h, there was no change by TLC and the reaction was stirred overnight at rt under Ar. The reaction was concentrated and the residue was purified through 15 g of silica using a solvent gradient from 100%  $CH_2Cl_2$  to 50%  $CH_2Cl_2/EtOAc$  and yielded **7.34** as a white solid (37.6 mg, 52% yield). Hydroxamate **7.34** was insoluble in  $CH_2Cl_2$  and  $CHCl_3$ , sparingly soluble in MeOH, and soluble in DMSO and  $H_2O$ . See below for full characterization data.

Optimized synthesis of hydroxamate ( $\pm$ )-7.34 using catalytic triflic acid (TfOH). Cycloadduct 3.38 (1.07 g, 5.40 mmol) was dissolved in 50 mL of anhydrous THF in a 200-mL round-bottomed flask that was rinsed with concentrated HCl (2x), washed with H<sub>2</sub>O, and acetone, then flame-dried under Ar. The solution was cooled in a crushed ice/H<sub>2</sub>O bath under Ar and trifluromethanesulfonic acid (0.010 mL, 0.11 mmol) was added. The solution was stirred in the ice/H<sub>2</sub>O bath under Ar and monitored by TLC (1:1 hexanes/EtOAc - UV lamp, CAM stain). After 1 h, the reaction was complete and the solution was warmed to rt and concentrated by rotary evaporation to a yellow oil. The oil was triturated with Et<sub>2</sub>O (50 mL) to yield pure **7.34** as a white powdery solid (0.562 g, 74% yield). <sup>1</sup>H NMR (500 MHz, d<sup>6</sup>-DMSO, 40 °C)  $\delta$  9.72 (s, 1H), 6.12 (d, *J* = 5.0 Hz, 1H), 5.85 (d, *J* = 5.0 Hz, 1H), 5.41 (d, *J* = 7.0 Hz, 1H), 4.34 (t, *J* = 6.0 Hz, 1H), 2.54 (m, 2H) ppm. <sup>13</sup>C NMR (125 MHz, d<sup>6</sup>-DMSO, 40 °C)  $\delta$  156.1, 135.9, 128.2, 81.4, 59.3, 36.1 ppm. HRMS (FAB) *m*/z [M+H]<sup>+</sup> calcd for C<sub>6</sub>H<sub>8</sub>NO<sub>3</sub><sup>+</sup>, 142.0504; obsd, 142.0521.


General procedure for the investigation of nitrone formation under acid (±)-(3aR,6aS)-2-Phenyl-4,6a-dihydro-3aH-cyclopenta[d]oxazole 3-oxide catalysis. (7.50). Cycloadduct 7.49 (0.104 g, 0.517 mmol) was dissolved in 10 mL of anhydrous CH<sub>2</sub>Cl<sub>2</sub> under Ar in a flame-dried single-necked 50-mL round-bottomed flask. Triflic acid (0.050 mL, 0.56 mmol) was added and the solution was stirred at rt and monitored by TLC. After 15 min, no starting material was observed by TLC (1:1 hexanes/EtOAc -UV lamp) and a new, more nonpolar spot was observed. Saturated NaHCO<sub>3</sub> (5 mL) was added and the layers were separated. The aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (2 x 5 mL) and the combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to yield an off-white residue. The residue was chromatographed through 10 g of silica using 100% CH<sub>2</sub>Cl<sub>2</sub> and yielded **7.50** as a yellow residue (19.5 mg, 20% yield). Rf = 0.48 (1:1 hexanes/EtOAc – UV lamp). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.85 (dd, J = 8.1, 1.3 Hz, 2H), 7.41 (m, 3H), 6.09 (dd, J = 5.7, 1.8 Hz, 1H), 5.97 (dd, J = 6.0, 1.8 Hz, 1H), 5.37 (m, 1H), 4.39 (q, J = 3.9 Hz, 1H), 2.71 (m, 2H) ppm. <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 172.6, 133.9, 130.7, 129.6, 128.2, 125.9, 81.8, 72.0, 37.4 ppm. HRMS (FAB) m/z [M+H]<sup>+</sup> calcd for C<sub>12</sub>H<sub>12</sub>NO<sub>2</sub><sup>+</sup>, 202.0868; obsd, 202.0869.

Attempted homo-Diels-Alder reaction with norbornadiene by direct oxidation of hydroxamate using NaIO<sub>4</sub>. A solution of 3.37 (47.5 mg, 0.357 mmol) in 2.8 mL of MeOH/H<sub>2</sub>O (4:1) was cooled to 0 °C (ice/H<sub>2</sub>O bath). Norbornadiene (0.045 mL, 0.442 mmol) was added (did not dissolve), followed by the dropwise addition of a

solution of sodium periodate (84.5 mg, 0.395 mmol) in 1.5 mL of  $H_2O$  over 5 min using a syringe. White solid was observed in the reaction and the mixture was stirred in the ice/ $H_2O$  bath for 1.5 h. The mixture was diluted with  $H_2O$  (20 mL), saturated with NaCl, and extracted with EtOAc (5 x 10 mL). The combined organic layers were washed with brine (2 x 10 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a light yellow oil (19 mg). NMR analysis did not indicate the formation of the desired adduct and was inconclusive.

Attempted homo-Diels-Alder reaction with norbornadiene by direct oxidation of hydroxamate using Swern oxidation. Compound 3.37 (44.3 mg, 0.333 mmol) was dissolved in 5.1:1 CH<sub>2</sub>Cl<sub>2</sub>/DMSO under Ar and cooled to -78 °C in a dry ice/acetone bath. In a separate flask, 1.6 mL of CH<sub>2</sub>Cl<sub>2</sub> was cooled to -78 °C and oxalyl chloride (0.12 mL, 1.40 mmol) was added followed by DMSO (0.15 mL, 2.11 mmol). This mixture was stirred for 5 min, then transferred to the flask containing 3.37 via cannula over 3-4 min. A yellow color was observed and the resultant solution was stirred at -78 °C for 20 min under Ar. Triethylamine (0.48 mL, 3.45 mmol) was added slowly over 5 min and the mixture was stirred at -78 °C for 5 min before slowly warming to rt. Et<sub>2</sub>O (10 mL), EtOAc (7 mL), and 1M HCl (7 mL) were added to the mixture and the layers were separated. The aqueous layer was extracted with EtOAc (2 x 10 mL) and the combined organic layers were washed with saturated NaHCO<sub>3</sub> (2 x 15 mL) and brine (2 x 15 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a yellow oil (25 mg). NMR analysis did not indicate the formation of the desired adduct and was inconclusive.

Attempted homo-Diels-Alder reaction with norbornadiene by direct oxidation of hydroxamate using Dess-Martin periodinane. Compound 7.65 (66.0 mg, 0.437 mmol) was dissolved in THF (4.5 mL) under Ar. Norbornadiene (0.050 mL, 0.492 mmol) was added and the solution was cooled to -78 °C in a dry ice/acetone bath. Dess-Martin periodinane (0.48M in CH<sub>2</sub>Cl<sub>2</sub>, 1.05 mL, 0.504 mmol) was added dropwise over 3 min and the mixture was stirred at -78 °C for 35 min under Ar, then warmed to rt and stirred for an additional 4 h. Saturated NaHCO<sub>3</sub> (5 mL) and saturated Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> (5 mL) were added and the mixture was stirred for 10 min until all of the solid material dissolved completely. The layers were separated and the aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (2 x 15 mL). The combined organic layers were washed with saturated NaHCO<sub>3</sub> (2 x 40 mL) and brine (2 x 40 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a white solid (36 mg). NMR analysis did not indicate the formation of the desired adduct and was inconclusive.



General procedure for cycloaddition reactions with 9,10-Dimethylanthracene using Bu<sub>4</sub>NIO<sub>4</sub>. Cycloadduct 7.67. Tetrabutylammonium periodate (0.204 g, 0.471 mmol) and 9,10-dimethylanthracene (80.8 mg, 0.392 mmol) were dissolved in 1 mL of CHCl<sub>3</sub> and cooled in an ice/H<sub>2</sub>O bath. A solution of **3.37** (62.4 mg, 0.469 mmol) in 0.25 mL of DMF was added dropwise over 3 min. The resultant mixture was stirred in the ice/H<sub>2</sub>O bath for 25 min, then at rt for an additional 1.5 h. The reaction was poured into

10 mL of EtOAc and washed with saturated Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> (2 x 10 mL), H<sub>2</sub>O (10 mL), and brine (2 x 10 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to yield a yellow solid (204 mg). Chromatography through 20 g of silica using 3:2 CH<sub>2</sub>Cl<sub>2</sub>/hexanes yielded **7.67** as a yellow solid (80.3 mg, 61% yield). Product obtained is 90% pure by NMR and contains 10% 9,10-dimethylanthracene. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.45 (dd, *J* = 5.9, 2.9 Hz, 2H), 7.37 (dd, *J* = 5.1, 3.4 Hz, 2H), 7.25 (dd, *J* = 5.6, 3.4 Hz, 4H), 2.57 (s, 3H), 2.23 (s, 3H), 1.22 (s, 9H) ppm.



**9,10-Dimethylanthracene cycloadduct 7.68.** Comopund **7.68** was prepared following the same procedure for **7.67**. Hydroxamic acid **7.65** (71.1 mg, 0.470 mmol), 9,10-dimethylanthracene (80.7 mg, 0.391 mmol), and tetrabutylammonium periodate (0.203 g, 0.469 mmol) provided a yellow oil. Chromatography through 30 g of silica using a solvent gradient from 2:1 to 3:1 CH<sub>2</sub>Cl<sub>2</sub>/hexanes yielded **7.68** as a yellow oil (62.6 mg, 45% yield). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.49 (d, *J* = 6.9 Hz, 2H), 7.31 – 7.12 (m, 6H), 7.12 (d, *J* = 6.3 Hz, 3H), 6.82 (dd, *J* = 7.5, 1.8 Hz, 2H), 3.54 (s, 2H), 2.75 (s, 3H), 2.07 (s, 3H) ppm.



1-(2,3-Dimethyl-1H-indol-1-yl)ethanone (7.82). 2,3-Dimethylindole (5.00 g, 34.4 mmol) was dissolved in 75 mL of acetic anhydride in a single-necked 250-mL round-bottomed flask fitted with a stir bar and reflux condenser. p-Toluenesulfonic acid monohydrate (68 mg, 0.36 mmol) was added and the deep red solution was heated to reflux in an oil bath (oil temperature = 130 - 150 °C) overnight. After 15 h, the resultant black mixture was cooled to rt, then concentrated by rotary evaporation to afford a black solid. TLC of the crude mixture seems to indicate starting material remained, however the product also appears at the same Rf in the solvent systems used for visualization (3:7 ether/pet Eter and 85:15 hexanes/EtOAc). The solid was loaded onto silica and purified through a Biotage 40M column using 100% hexanes to 95% hexanes/EtOAc and yielded **7.82** as a white solid (4.61 g, 72% yield). The solid gradually became light orange after standing for 1 h to overnight, and was stored in a scintillation vial under Ar at -10 °C until use. mp = 66-68 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.99 (m, 1H), 7.45 (m, 1H), 7.27 (m, 2H), 2.72 (s, 3H), 2.57 (s, 3H), 2.21 (s, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 170.1, 135.5, 132.5, 131.2, 123.6, 122.8, 118.1, 115.4, 114.9, 27.5, 14.4, 8.7 ppm. HRMS (FAB) m/z [M]<sup>+</sup> calcd for C<sub>12</sub>H<sub>13</sub>NO<sup>+</sup>, 187.0997; obsd, 187.1003.

## APPENDIX A:

## NMR SPECTRA FOR SELECTED COMPOUNDS



 $^{1}$ H and  $^{13}$ C NMR spectra for compound **2.62** 





<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.8** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.9** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.10** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.14a** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.14b** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.17** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.20** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.27** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.28** 





<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.29** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.31** 

300 MHz, d6-DMSO



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.32** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.33** 





500 MHz, d6-DMSO



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.34** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.37** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.38** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.39** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.40** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.42** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.43** 



 $^{1}$ H and  $^{13}$ C NMR spectra for compound **3.45** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.47** 



<sup>1</sup>H NMR spectrum for compound **3.48** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.52** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.53** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.59** 



COSY NMR spectra for compound 3.59



HETCOR NMR spectra for compound 3.59



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.63**


<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.64** (includes impurities)



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.66** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.69** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.71** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.73** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.74** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.75** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.76** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.77** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.78** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.79** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.80** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.81** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.86** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.90** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.92** (includes impurities)



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.93** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.94** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.95** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.96** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.97** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.98** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.99** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.100** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.101** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.102** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.105** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.106** 





<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.108** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.110** 



COSY NMR spectra for compound 3.110



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.111** 



COSY NMR spectra for compound 3.111



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.115** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.116** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.117**


<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.118** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.119** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.120** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.121** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.122** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.124** 



COSY NMR spectra for compound 3.124



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **3.125** 



COSY NMR spectra for compound 3.125



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **4.6** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for a mixture of compounds **4.7a** and **4.7b** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **4.8** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for the purified diastereomer of compound **4.10** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **4.20** 



<sup>1</sup>H NMR spectra for compound **4.22** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **4.23** 







<sup>1</sup>H NMR spectra for compound **4.25** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **4.26** 



<sup>1</sup>H NMR spectra for compound **4.27** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **4.28** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **4.31** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **4.32** 





<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **4.36** 





<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **4.36**'







<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **4.42** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for a mixture of diastereomers **4.47** and **4.48** 





<sup>1</sup>H NMR spectra for compound **4.53** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **4.54** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **4.56** 



COSY NMR spectra for compound 4.56



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **4.57** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **4.58** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **4.59** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **4.60** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **4.61** 



 $^{1}$ H and  $^{13}$ C NMR spectra for compound **4.62**


COSY NMR spectra for compound 4.62



HETCOR NMR spectra for compound 4.62



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **4.63** 



COSY NMR spectra for compound 4.64



<sup>1</sup>H NMR spectra for compound **4.65** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **5.20** 



<sup>1</sup>H NMR spectra for compound **5.21** 



<sup>1</sup>H NMR spectra for compound **5.22** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **5.23** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **5.24** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **5.25** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **5.26** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **5.27** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **5.28** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **5.30** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **5.31** 



 $^{1}$ H and  $^{13}$ C NMR spectra for compound **5.32** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **5.33** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **5.34** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **5.35** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **5.36a** 











<sup>1</sup>H NMR spectra for compound **5.36c** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **5.37** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **5.38a** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **5.38b** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **5.40a** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **5.40b** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **5.41a** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **5.41b** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **5.67** 



<sup>1</sup>H NMR spectra for compound **5.68** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **5.69b** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **5.70b** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **5.73** 



 $^{1}$ H and  $^{13}$ C NMR spectra for compound 5.75



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **5.81** 





<sup>1</sup>H NMR spectra for compound **5.82**


<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **6.7a** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **6.7b** 



HETCOR NMR spectra for compound 6.7b



gHMBC NMR spectra for compound 6.7b



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **6.7c** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **6.7d** 



COSY NMR spectra for compound 6.7d



ROESY NMR spectra for compound 6.7d



HETCOR NMR spectra for compound 6.7d



gHMBC NMR spectra for compound 6.7d



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **6.7e** 



HETCOR NMR spectra for compound 6.7e



gHMBC NMR spectra for compound 6.7e



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **6.8a** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **6.8b** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **6.8c** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **6.8d** 



COSY NMR spectra for compound 6.8d



ROESY NMR spectra for compound 6.8d



HETCOR NMR spectra for compound 6.8d



gHMBC NMR spectra for compound 6.8d



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **6.8e** 



COSY NMR spectra for compound 6.8e



ROESY NMR spectra for compound 6.8e



gHMBC NMR spectra for compound 6.8e



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **6.10** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **6.11** 



COSY NMR spectra for compound 6.11



ROESY NMR spectra for compound 6.11



HETCOR NMR spectra for compound 6.11



gHMBC NMR spectra for compound 6.11



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **6.12** 



COSY NMR spectra for compound 6.12



ROESY NMR spectra for compound 6.12



HETCOR NMR spectra for compound 6.12



gHMBC NMR spectra for compound 6.12


<sup>1</sup>H and <sup>13</sup>C NMR spectra for a mixture of compounds **6.13** and **6.14** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **6.17** 



Fromation of compound **6.17** monitored by <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz)



<sup>1</sup>H NMR spectra for compound **6.19** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **6.20** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **6.21** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **7.34** 



COSY NMR spectra for compound 7.34





<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **7.50** 







<sup>1</sup>H NMR spectra for compound **7.68** 



<sup>1</sup>H and <sup>13</sup>C NMR spectra for compound **7.82** 

## REFERENCES

- 1. Yamamoto, H.; Kawasaki, M., Nitroso and Azo Compounds in Modern Organic Synthesis: Late Blooming but Very Rich. *Bull. Chem. Soc. Jpn.* **2007**, 80, (4), 595-607.
- 2. Yamamoto, Y.; Yamamoto, H., Recent Advances in Asymmetric Nitroso Diels-Alder Reactions. *Eur. J. Org. Chem.* **2006**, 2006, (9), 2031-2043.
- 3. Vogt, P. F.; Miller, M. J., Development and Applications of Amino Acid-Derived Chiral Acylnitroso Hetero Diels-Alder Reactions. *Tetrahedron* **1998**, 54, 1317-1348.
- 4. Streith, J.; Defoin, A., Azasugar syntheses and multi-step cascade rearrangements via hetero Diels-Alder cycloadditions with nitroso dienophiles. *Synlett* **1996**, (3), 189-200.
- 5. Streith, J.; Defoin, A., Hetero Diels-Alder Reactions with Nitroso Dienophiles: Application to the Synthesis of Natural Product Derivatives. *Synthesis* **1994**, (11), 1107-1117.
- 6. Waldmann, H., Asymmetric Hetero Diels-Alder Reactions. *Synthesis* **1994**, (6), 535-551.
- 7. Weinreb, S. M.; Staib, R. R., Synthetic aspects of diels-alder cycloadditions with heterodienophiles. *Tetrahedron* **1982**, 38, (21), 3087-3128.
- 8. Baeyer, A., Nitrosobenzol und Nitrosonaphtalin. *Ber.* **1874**, 7, (2), 1638-1640.
- 9. Ehrlich, P.; Sachs, F., Ueber Condensationen von aromatischen Nitrosoverbindungen mit Methylenderivaten. *Chem. Ber.* **1899**, 32, (2), 2341-2346.
- 10. Yamamoto, H.; Momiyama, N., Rich chemistry of nitroso compounds. *Chem. Commun.* **2005**, (28), 3514-3525.
- Momiyama, N.; Yamamoto, H., Bronsted acid catalysis of achiral enamine for regio- and enantioselective nitroso aldol synthesis. J. Am. Chem. Soc. 2005, 127, (4), 1080-1081.

- 12. Yamamoto, Y.; Momiyama, N.; Yamamoto, H., Enantioselective Tandem O-Nitroso Aldol/Michael Reaction. J. Am. Chem. Soc. **2004**, 126, (19), 5962-5963.
- 13. Momiyama, N.; Yamamoto, H., Simple Synthesis of a-Hydroxyamino Carbonyl Compounds: New Scope of the Nitroso Aldol Reaction. *Org. Lett.* **2002**, 4, (21), 3579-3582.
- 14. Adam, W.; Krebs, O., The Nitroso Ene Reaction: A Regioselective and Stereoselective Allylic Nitrogen Functionalization of Mechanistic Delight and Synthetic Potential. *Chem. Rev.* **2003**, 103, (10), 4131-4146.
- 15. Sha, X.; Isbell, T. S.; Patel, R. P.; Day, C. S.; King, S. B., Hydrolysis of Acyloxy Nitroso Compounds Yields Nitroxyl (HNO). J. Am. Chem. Soc. 2006, 128, (30), 9687-9692.
- 16. Bahr, N.; Guller, R.; Reymond, J. L.; Lerner, R. A., A Nitroxyl Synthase Catalytic Antibody. J. Am. Chem. Soc. **1996**, 118, (15), 3550-3555.
- 17. Kirby, G. W., Tilden Lecture. Electrophilic C-nitroso-compounds. *Chem. Soc. Rev.* **1977**, 6, (1), 1-24.
- 18. Labaziewicz, H.; Lindfors, K. R.; Kejonen, T. H., The synthesis of novel N-pyridyl-3,6-dihydro-1,2-oxazines. *Heterocycles* **1989**, 29, (12), 2327-2334.
- 19. Noguchi, H.; Aoyama, T.; Shioiri, T., Total synthesis and absolute configuration of radiosumin, a strong trypsin inhibitor from the blue-green alga Plectonema radiosum. *Heterocycles* **2002**, 58, 471-504.
- 20. Zhang, D.; Sueling, C.; Miller, M. J., The Hetero Diels-Alder Reactions between D-Mannose-Derived Halonitroso Compounds and Cyclopentadiene: Scope and Limitations. *J. Org. Chem.* **1998**, 63, (3), 885-888.
- Tronchet, J. M. J.; Jean, E.; Barbalat-Rey, F.; Bernardinelli, G., Hetero Diels-Alder adducts of deoxy C-nitroso sugar derivatives. J. Chem. Res., Synop. 1992, (7), 228-229.
- 22. Calvet, G.; Guillot, R.; Blanchard, N.; Kouklovsky, C., Intermolecular nitroso Diels-Alder cycloaddition of a-acetoxynitroso derivatives in aqueous medium. *Org. Biomol. Chem.* **2005**, *3*, (24), 4395-4401.
- 23. Calvet, G.; Dussaussois, M.; Blanchard, N.; Kouklovsky, C., Lewis Acid-Promoted Hetero Diels-Alder Cycloaddition of alpha-Acetoxynitroso Dienophiles. *Org. Lett.* **2004**, 6, (14), 2449-2451.
- Miller, C. A.; Batey, R. A., Hetero Diels-Alder Reactions of Nitrosoamidines: An Efficient Method for the Synthesis of Functionalized Guanidines. *Org. Lett.* 2004, 6, (5), 699-702.

- 25. Kirby, G. W.; McGuigan, H.; Mackinnon, J. W. M.; Mclean, D.; Sharma, R. P., Formation and reactions of C-nitrosoformate esters, a new class of transient dienophiles. *J. Chem. Soc.*, *Perkin Trans. 1* **1985**, 1437-1442.
- 26. Ware, R. W.; Day, C. S.; King, S. B., Diastereoselective and Intramolecular Cycloadditions of Asymmetric P-Nitroso Phosphine Oxides. *J. Org. Chem.* **2002**, 67, (17), 6174-6180.
- 27. Ware, R. W., Jr.; King, S. B., P-Nitrosophosphate Compounds: New N-O Heterodienophiles and Nitroxyl Delivery Agents. J. Org. Chem. 2000, 65, (25), 8725-8729.
- 28. Ware, R. W., Jr.; King, S. B., N-Phosphinoylnitroso Compounds: New Asymmetric N-O Heterodienophiles and Nitroxyl Delivery Agents. *J. Am. Chem. Soc.* **1999**, 121, (28), 6769-6770.
- 29. Singal, K. K.; Singh, B.; Raj, B., Studies in 1-Aza-1,3-butadienes: Diels-Alder Cycloaddition Reactions of 1,4-Diaryl-1-aza-1,3-butadienes with Aryl Sulphonyl Nitrosites Leading to the Synthesis of New Oxadiazines. *Synth. Commun.* **1993**, 23, (1), 107 114.
- 30. Gouverneur, V.; Dive, G.; Ghosez, L., Asymmetric Diels-Alder reactions of a nitroso compound derived from D-bornane-10,2-sultam. *Tetrahedron: Asymmetry* **1991**, 2, (12), 1173-1176.
- 31. Kirby, G. W.; Sweeny, J. G., Nitrosocarbonyl compounds as intermediates in the oxidative cleavage of hydroxamic acids. J. Chem. Soc., Chem. Commun. 1973, (19), 704-705.
- 32. Kirby, G. W.; Sweeny, J. G., Formation and dienophilic reactions of transient Cnitrosocarbonyl compounds. *J. Chem. Soc.*, *Perkin Trans. 1* **1981**, (12), 3250-3254.
- 33. Martin, S. F.; Hartmann, M.; Josey, J. A., Diastereoselective [4+2] Cycloadditions of Acyl Nitroso Compounds. *Tetrahedron Lett.* **1992**, 33, (25), 3583-3586.
- 34. Dao, L. H.; Dust, J. M.; Mackay, D.; Watson, K. N., The formation and interconversion of oxazines and dioxazines from the reaction of nitrosocarbonyl compounds with cyclopentadienes. *Can. J. Chem.* **1979**, 57, (13), 1712–1719.
- 35. Jenkins, N. E.; Ware, R. W., Jr.; Atkinson, R. N.; King, S. B., Generation of acyl nitroso compounds by the oxidation of N-acyl hydroxylamines with the Dess-Martin periodinane. *Synth. Commun.* **2000**, 30, (5), 947-953.
- 36. Howard, J. A. K.; Ilyashenko, G.; Sparkes, H. A.; Whiting, A., Development of new transition metal catalysts for the oxidation of a hydroxamic acid with in situ Diels-Alder trapping of the acyl nitroso derivative. *Dalton Trans.* **2007**, (21), 2108-2111.

- 37. Iwasa, S.; Fakhruddin, A.; Tsukamoto, Y.; Kameyama, M.; Nishiyama, H., Iridium(I)-catalyzed hydrogen peroxide oxidation of hydroxamic acids and hetero Diels-Alder reaction of the acyl nitroso intermediates with cyclopentadiene. *Tetrahedron Lett.* **2002**, 43, (35), 6159-6161.
- 38. Flower, K. R.; Lightfoot, A. P.; Wan, H.; Whiting, A., The development and application of ruthenium catalyzed oxidations of a hydroxamic acid and in situ Diels-Alder trapping of the acyl nitroso derivative. *J. Chem. Soc., Perkin Trans. 1* **2002**, (18), 2058-2064.
- 39. Flower, K. R.; Lightfoot, A. P.; Wan, H.; Whiting, A., Direct evidence for a ruthenium(iv) oxo complex-mediated oxidation of a hydroxamic acid in the presence of phosphine oxide donors. *Chem. Commun. (Cambridge, U. K.)* **2001**, (18), 1812-1813.
- 40. Iwasa, S.; Tajima, K.; Tsushima, S.; Nishiyama, H., A mild oxidation method of hydroxamic acids: efficient trapping of acyl nitroso intermediates. *Tetrahedron Lett.* **2001**, 42, (34), 5897.
- 41. Adamo, M. F. A.; Bruschi, S., Generation of Acylnitroso Dienophiles: A Study of Metal Catalysis. *J. Org. Chem.* **2007**, 72, (7), 2666-2669.
- 42. Quadrelli, P.; Mella, M.; Invernizzi, A. G.; Caramella, P., The mild oxidation of nitrile oxides affords a convenient entry to nitrosocarbonyl intermediates, versatile tools in organic syntheses. *Tetrahedron* **1999**, 55, (34), 10497-10510.
- 43. Corrie, J. E. T.; Kirby, G. W.; Mackinnon, J. W. M., Reactions of transient Cnitrosocarbonyl compounds with dienes, monoolefins, and nucleophiles. *J. Chem. Soc., Perkin Trans. 1* 1985, (4), 883-886.
- 44. Quadrelli, P.; Mella, M.; Caramella, P., A photochemical generation of nitrosocarbonyl intermediates. *Tetrahedron Lett.* **1999**, 40, (4), 797-800.
- 45. O'Bannon, P. E.; P., W. D., Rearrangement of nitrocarbenes to acyl nitroso compounds. *Tetrahedron Lett.* **1988**, 29, (45), 5719-5722.
- 46. O'Bannon, P. E.; Suelzle, D.; Schwarz, H., Characterization of nitrosocarbonyl compounds by neutralization-reionization mass spectrometry. *Helv. Chim. Acta* **1991**, 74, (8), 2068-2072.
- 47. Cohen, A. D.; Zeng, B. B.; King, S. B.; Toscano, J. P., Direct Observation of an Acyl Nitroso Species in Solution by Time-Resolved IR Spectrocopy. *J. Am. Chem. Soc.* **2003**, 125, (6), 1444-1445.
- 48. Gouverneur, V.; McCarthy, S. J.; Mineur, C.; Belotti, D.; Dive, G.; Ghosez, L., New acylnitroso compounds for the asymmetric oxyamination of dienes. *Tetrahedron* **1998**, 54, (35), 10537-10554.

- 49. Defoin, A.; Brouillard-Poichet, A.; Streith, J., Asymmetric Diels-Alder Cycloadditions with Chiral Carbamoyl Dienophiles. *Helv. Chim. Acta* **1992**, 75, (1), 109-123.
- 50. Wang, Y.-C.; Lu, T.-M.; Elango, S.; Lin, C.-K.; Tsai, C.-T.; Yan, T.-H., Switch in asymmetric induction sense in cycloadditions using camphor-based nitroso dienophiles. *Tetrahedron: Asymmetry* **2002**, 13, (7), 691-695.
- 51. Cardillo, B.; Galeazzi, R.; Mobbili, G.; Orena, M.; Rossetti, M., Diastereoselective hetero-diels-alder cycloaddition of a C-nitroso compound prepared starting from a homochiral imidazolidin-2-one. *Tetrahedron: Asymmetry* **1994**, 5, (8), 1535-1540.
- 52. Miller, A.; Procter, G., On the origin of the diastereoselectivity observed in cycloadditions of chiral acyl-nitroso compounds. *Tetrahedron Lett.* **1990**, 31, (7), 1041-1042.
- 53. Miller, A.; Paterson, T. M.; Procter, G., Stereoselective cycloadditions of chiral acyl-nitroso compounds. *Synlett* **1989**, (1), 32-34.
- 54. Corrie, J. E. T.; Kirby, G. W.; Sharma, R. P., Formation of isocyanates by deoxygenation of C-nitrosocarbonyl compounds. *J. Chem. Soc., Perkin Trans.* 1 **1982**, (7), 1571-1574.
- 55. Wichterle, O., Addition of nitroso compounds to a conjugated system. *Collect. Czech. Chem. Commun.* **1947,** 12, 292-304.
- 56. Arbuzov, Y. A., Reaction of diene hydrocarbons with nitroso compounds. Addition of 1,3-butadiene to nitrosobenzene. *Dokl. Akad. Nauk SSSR* **1948**, 60, 993-996.
- 57. Leach, A. G.; Houk, K. N., Diels-Alder and ene reactions of singlet oxygen, nitroso compounds and triazolinediones: transition states and mechanisms from contemporary theory. *Chem. Commun.* **2002**, (12), 1243-1255.
- 58. Leach, A. G.; Houk, K. N., Transition States and Mechanisms of the Hetero-Diels-Alder Reactions of Hyponitrous Acid, Nitroso alkanes, Nitroso arenes, and Nitroso carbonyl Compounds. *J. Org. Chem.* **2001**, 66, (15), 5192-5200.
- 59. McCarrick, M. A.; Wu, Y. D.; Houk, K. N., Hetero-Diels-Alder reaction transition structures: reactivity, stereoselectivity, catalysis, solvent effects, and the exo-lone-pair effect. *J. Org. Chem.* **1993**, 58, (12), 3330-3343.
- 60. McCarrick, M. A.; Wu, Y. D.; Houk, K. N., exo-Lone-pair effect on hetero-Diels-Alder cycloaddition stereochemistry. *J. Am. Chem. Soc.* **1992**, 114, (4), 1499-1500.

- 61. Boger, D. L.; Patel, M.; Takusagawa, F., Regioselectivity of the intermolecular Diels-Alder reaction of acyl nitroso compounds (C-nitrosocarbonyl compounds) and nitrosoformates (O-nitrosocarbonyl compounds). Preparation of functionalized cis-.DELTA.6-1-octalones. *J. Org. Chem.* **1985**, 50, (11), 1911-1916.
- Lemire, A.; Beaudoin, D.; Grenon, M.; Charette, A. B., [4+2] Cycloaddition of 2-Substituted 1,2-Dihydropyridines with Nitrosobenzene: Asymmetric Synthesis of <i>trans</i>-2-Substituted 3-Amino-1,2,3,6-tetrahydropyridines. J. Org. Chem. 2005, 70, (6), 2368-2371.
- 63. Dubey, S. K.; Knaus, E. E., Some regio- and stereochemical aspects of the Diels-Alder reaction of nitrosocarbonyl compounds with N-substituted 1,2dihydropyridines. *J. Org. Chem.* **1985**, 50, (12), 2080-2086.
- 64. Aoyagi, S.; Tanaka, R.; Naruse, M.; Kibayashi, C., Asymmetric total synthesis of (-)-epibatidine. *Tetrahedron Lett.* **1998**, 39, (25), 4513-4516.
- 65. Vogt, P. F.; Hansel, J.-G.; Miller, M. J., Asymmetric synthesis of an important precursor to 5'-nor carbocyclic nucleosides. *Tetrahedron Lett.* **1997**, 38, (16), 2803-2804.
- 66. Ritter, A. R.; Miller, M. J., Amino Acid-Derived Chiral Acyl Nitroso Compounds: Diastereoselectivity in Intermolecular Hetero Diels-Alder Reactions. *J. Org. Chem.* **1994**, 59, 4602-4611.
- 67. Pepper, A. G.; Procter, G.; Voyle, M., Stereoselective cycloadditions of chiral acyl-nitroso compounds; selective reactions of ring-cleaved cycloadducts leading to a new approach to polyoxamic acid. *Chem. Commun.* **2002**, (10), 1066-1067.
- 68. Morley, A. D.; Hollinshead, D. M.; Procter, G., Stereoselective cycloadditions of acyl-nitroso compounds; an approach to carbapenem synthesis. *Tetrahedron Lett.* **1990,** 31, (7), 1047-1050.
- 69. Miller, A.; Procter, G., Stereoselective cycloadditions of chiral acyl-nitroso compounds; synthetically useful reactions of the adducts. *Tetrahedron Lett.* **1990**, 31, (7), 1043-1046.
- 70. Kirby, G. W.; Nazeer, M., Asymmetric induction in the diels-alder reactions of alpha-hydroxyacylnitroso compounds. *Tetrahedron Lett.* **1988**, 29, (47), 6173-6174.
- 71. Behr, J.-B.; Chevrier, C.; Defoin, A.; Tarnus, C.; Streith, J., Asymmetric synthesis of potent glycosidase and very potent [alpha]-mannosidase inhibitors: 4-amino-4-deoxy--erythrose and 4-amino-4,5-dideoxy--ribose. *Tetrahedron* **2003**, 59, (4), 543-553.

- 72. Defoin, A.; Pires, J.; Streith, J., Asymmetric hetero-Diels-Alder cycloadditions using a chiral N-dienyl lactam and acylnitroso dienophiles. *Synlett* **1991**, (6), 417-419.
- 73. Hussain, A.; Wyatt, P. B., 2-(penta-1,3-dienyl)oxazolidines: synthesis of hydroxylated piperidines by a stereoselective diels-alder Reaction. *Tetrahedron* **1993**, 49, (10), 2123-2130.
- Arribas, C.; Carreno, M. C.; Garcia-Ruano, J. L.; Rodriguez, J. F.; Santos, M.; AscensionSanz-Tejedor, M., First Asymmetric Hetero Diels-Alder Reaction of 1-Sulfinyl Dienes with Nitroso Derivatives. A New Entry to the Synthesis of Optically Pure 1,4-Imino-L-ribitol Derivatives. Org. Lett. 2000, 2, (20), 3165-3168.
- 75. Carreno, M. C.; Cid, M. B.; Garcia Ruano, J.; Santos, M., Short enantioselective approach to substituted triazolopyridazines from [(S)R]-1-(1E, 3E)-1-p-tolylsulfinyl-1,3-pentadiene. *Tetrahedron Lett.* **1998**, 39, (11), 1405-1408.
- 76. Hudlicky, T.; Olivo, H. F., Stereospecific synthesis of aminocyclitols via cycloadditions of unsymmetrical, optically pure dienes: conduramine A-1 and dihydroconduramine A-1. *Tetrahedron Lett.* **1991**, 32, (43), 6077-6080.
- 77. Lightfoot, A. P.; Pritchard, R. G.; Wan, H.; Warren, J. E.; Whiting, A., A novel scandium ortho-methoxynitrosobenzene-dimer complex: mechanistic implications for the nitroso-Diels-Alder reaction. *Chem. Commun.* **2002**, (18), 2072-2073.
- 78. Yamamoto, Y.; Yamamoto, H., Catalytic, Highly Enantio, and Diastereoselective Nitroso Diels-Alder Reaction. J. Am. Chem. Soc. **2004**, 126, (13), 4128-4129.
- 79. Krchnak, V.; Moellmann, U.; Dahse, H.-M.; Miller, M. J., Solid-Supported Nitroso Hetero Diels-Alder Reactions. 1. Acylnitroso Dienophiles: Scope and Limitations. *J. Comb. Chem.* **2008**, 10, (1), 94-103.
- Krchnak, V.; Moellmann, U.; Dahse, H.-M.; Miller, M. J., Solid-Supported Nitroso Hetero-Diels-Alder Reactions. 3. Acid-Mediated Transformation of Cycloadducts by Scission of the Oxazine C−O Bonds. J. Comb. Chem. 2008, 10, (1), 112-117.
- 81. Krchnak, V.; Moellmann, U.; Dahse, H.-M.; Miller, M. J., Solid-Supported Nitroso Hetero Diels-Alder Reactions. 2. Arylnitroso Dienophiles: Scope and Limitations. *J. Comb. Chem.* **2008**, 10, (1), 104-111.
- Faita, G.; Mella, M.; Paio, A. M.; Quadrelli, P.; Seneci, P., Solid-Supported Nitrile Oxides as Stable and Valuable Reactive Intermediates. *Eur. J. Org. Chem.* 2002, 2002, (7), 1175-1183.
- 83. Quadrelli, P.; Scrocchi, R.; Piccanello, A.; Caramella, P., Photochemical Generation of Nitrosocarbonyl Intermediates on Solid Phase: Synthons toward

Hetero Diels-Alder and Ene Adducts through Photocleavage. J. Comb. Chem. 2005, 7, (6), 887-892.

- 84. Nahm, S.; Weinreb, S. M., N-methoxy-n-methylamides as effective acylating agents. *Tetrahedron Lett.* **1981**, 22, (39), 3815-3818.
- 85. Keck, G. E.; Webb, R. R.; Yates, J. B., A versatile method for carbon-nitrogen bond formation via ene reactions of acylnitroso compounds. *Tetrahedron* **1981**, 37, (23), 4007-4016.
- 86. Tsoungas, P. G., Synthesis of 1,2-oxazines and their N-oxides. *Heterocycles* **2002**, 57, (6), 1149-1178.
- 87. Tsoungas, P. G., 1,2-oxazines and their N-oxides in synthesis. *Heterocycles* **2002**, 57, (5), 915-953.
- 88. Le Flohic, A.; Meyer, C.; Cossy, J.; Desmurs, J.-R., Synthesis of unsaturated [1,2]oxazines by using sigmatropic rearrangements and the ring-closing metathesis reaction. *Tetrahedron Lett.* **2003**, 44, (47), 8577-8580.
- 89. Yang, Y. K.; Choi, J. H.; Tae, J., Synthesis of 2,3-Di- and 2,3,4-Trisubstituted Furans from 1,2-Dioxines Generated by an Enyne-RCM/Diels-Alder Reaction Sequence. *J. Org. Chem.* **2005**, 70, (17), 6995-6998.
- 90. Yang, Y.-K.; Tae, J., Ring-closing metathesis of enynes tethered by an N-O bond: Synthesis of 1,2-oxaza polycycles by diels-alder reaction of the ring-closing metathesis products. *Synlett* **2003**, (13), 2017-2020.
- 91. Pulz, R.; Cicchi, S.; Brandi, A.; Reißig, H.-U., Synthesis of New Enantiopure Bicyclic 1,2-Oxazines by Addition of Lithiated Methoxyallene to Chiral Cyclic Nitrones. *Eur. J. Org. Chem.* **2003**, 2003, (7), 1153-1156.
- 92. Sibi, M. P.; Ma, Z.; Jasperse, C. P., Enantioselective Addition of Nitrones to Activated Cyclopropanes. J. Am. Chem. Soc. 2005, 127, (16), 5764-5765.
- 93. Ganton, M. D.; Kerr, M. A., Magnesium Iodide Promoted Reactions of Nitrones with Cyclopropanes: A Synthesis of Tetrahydro-1,2-oxazines. *J. Org. Chem.* **2004**, 69, (24), 8554-8557.
- 94. Kumarn, S.; Shaw, D. M.; Longbottom, D. A.; Ley, S. V., A Highly Selective, Organocatalytic Route to Chiral Dihydro-1,2-oxazines. *Org. Lett.* **2005**, 7, (19), 4189-4191.
- 95. Gallos, J. K.; Stathakis, C. I.; Kotoulas, S. S.; Koumbis, A. E., An Improved Approach to Chiral Cyclopentenone Building Blocks. Total Synthesis of Pentenomycin I and Neplanocin A. *J. Org. Chem.* **2005**, 70, (17), 6884-6890.

- 96. Cicchi, S.; Goti, A.; Brandi, A.; Guarna, A.; De Sarlo, F., 1,3-Aminoalcohols by Reductive Cleavage of Isoxazolidines With Molybdenum Hexacarbonyl. *Tetrahedron Lett.* **1990**, 31, (23), 3351-3354.
- 97. Nitta, M.; Kobayashi, T., Metal-carbonyl-induced reaction of isoxazoles. Ring cleavage and reduction by hexacarbonylmolybdenum, pentacarbonyliron, or nonacarbonyldiiron. J. Chem. Soc., Perkin Trans. 1 1985, (7), 1401-6.
- 98. Keck, G. E.; Wager, T. T.; McHardy, S. F., Reductive cleavage of N---O bonds in hydroxylamines and hydroxamic acid derivatives using samarium diiodide. *Tetrahedron* **1999**, **55**, (40), 11755-11772.
- 99. Keck, G. E.; McHardy, S. F.; Wager, T. T., Reductive cleavage of N-O bonds in hydroxylamine and hydroxamic acid derivatives using SmI2/THF. *Tetrahedron Lett.* **1995**, 36, (41), 7419-7422.
- 100. Lee, Y. H.; Choo, D. J., Photochemistry of 3,6-dihydro-1,2-oxazines. 1. A mechanistic study for pyrrole cyclization. *Bull. Korean Chem. Soc.* **1993**, 14, (4), 423-424.
- 101. Klier, K.; Kresze, G.; Werbitzky, O.; Simon, H., The microbial reductive splitting of the N---O bond of dihydrooxazines; an alternative to the chemical reduction. *Tetrahedron Lett.* **1987**, 28, (24), 2677-2680.
- 102. Calvet, G.; Blanchard, N.; Kouklovsky, C., Synthesis of polysubstituted pyrroles from nitroso-Diels-Alder cycloadducts. *Synthesis* **2005**, (19), 3346-3354.
- 103. Mulvihill, M. J.; Gage, J. L.; Miller, M. J., Enzymatic Resolution of Aminocyclopentenols as Precursors to D- and L-Carbocyclic Nucleosides. *J. Org. Chem.* **1998**, 63, (10), 3357-3363.
- 104. Kefalas, P.; Grierson, D. S., Synthesis of pyrrolo-castanospermine analogs: A novel fluoride ion induced rearrangement of a 3-trialkylsilyl substituted dihydro-2H-1,2-oxazine to a pyrrole. *Tetrahedron Lett.* **1993**, 34, (22), 3555-3558.
- 105. Kornblum, N.; DeLaMare, H. E., The Base Catalyzed Decomposition of a Dialkyl Peroxide. *J. Am. Chem. Soc.* **1951**, 73, (2), 880-881.
- Desai, M. C.; Doty, J. L.; Stephens, L. M.; Brighty, K. E., Novel fragmentation of 3-lithio-3,6-dihydro-1,2-oxazines: Synthesis of 2,5-dihydro-2-furanylamines. *Tetrahedron Lett.* 1993, 34, (6), 961-962.
- 107. Mulvihill, M. J.; Surman, M. D.; Miller, M. J., Regio- and Stereoselective Fe(III)and Pd(0)-Mediated Ring Openings of 3-Aza-2-oxabicyclo[2.2.1]hept-5-ene Systems. J. Org. Chem. **1998**, 63, (15), 4874-4875.

- 108. Surman, M. D.; Miller, M. J., Regio- and stereochemically controlled formation of hydroxamic acid containing anti- or syn-1,4-cycloalkenols from acylnitrosoderived Diels-Alder adducts. *J. Org. Chem.* **2001**, 66, (7), 2466-2469.
- 109. Muxworthy, J. P.; Wilkinson, J. A.; Procter, G., Stereoselective cycloadditions of chiral acyl-nitroso compounds; unexpected formation of an oxazolidinone. *Tetrahedron Lett.* **1995**, 36, (41), 7539-7540.
- 110. Muxworthy, J. P.; Wilkinson, J. A.; Procter, G., Stereoselective cycloadditions of chiral acyl-nitroso compounds; hydrolytic reactions of a clyclopentadiene adduct. *Tetrahedron Lett.* **1995**, 36, (41), 7535-7538.
- 111. Surman, M. D.; Mulvihill, M. J.; Miller, M. J., Regio- and Stereoselective Ring Openings of 3-Aza-2-oxabicyclo[2.2.1]hept-5-ene Systems with Copper Catalyst-Modified Grignard Reagents: Application to the Synthesis of an Inhibitor of 5-Lipoxygenase. J. Org. Chem. 2002, 67, (12), 4115-4121.
- 112. Pineschi, M.; DelMoro, F.; Crotti, P.; Macchia, F., Catalytic Asymmetric Ring Opening of 2,3-Substituted Norbornenes with Organometallic Reagents: A New Formal Aza Functionalization of Cyclopentadiene. *Org. Lett.* **2005**, 7, (17), 3605-3607.
- 113. Pineschi, M.; Del Moro, F.; Crotti, P.; Macchia, F., Catalytic asymmetric ringopening of cyclopentadiene-heterodienophile cycloadducts with organometallic reagents. *Pure Appl. Chem.* **2006**, 78, (2), 463-467.
- 114. Chen, W.; Day, C. S.; King, S. B., Grignard Reagent-Mediated Conversion of an Acyl Nitroso-anthracene Cycloadduct to a Nitrone. *J. Org. Chem.* **2006**, 71, (24), 9221-9224.
- 115. Surman, M. D.; Mulvihill, M. J.; Miller, M. J., Novel 1,4-Benzodiazepines from Acylnitroso-Derived Hetero-Diels-Alder Cycloadducts. *Org. Lett.* **2002**, 4, (1), 139-141.
- 116. Lee, W.; Kim, K.-H.; Surman, M. D.; Miller, M. J., Stereo- and Regioselectivity of Pd0/InI-Mediated Allylic Additions to Aldehydes and Ketones. In Situ Generation of Allylindium(III) Intermediates from N-Acylnitroso Diels-Alder Cycloadducts and 1-Amino-4-acetoxycyclopentenes. J. Org. Chem. 2003, 68, (1), 139-149.
- 117. Quadrelli, P.; Mella, M.; Paganoni, P.; Caramella, P., Cycloadditions of Nitrile Oxides to the Highly Reactive N-Acyl-2-oxa-3-azanorborn-5-enes Afford Versatile Cycloadducts and a Convenient Entry to Highly Functionalized Derivatives. *Eur. J. Org. Chem.* **2000**, 14, 2613-2620.
- 118. Quadrelli, P.; Scrocchi, R.; Caramella, P.; Rescifina, A.; Piperno, A., From cyclopentadiene to isoxazoline-carbocyclic nucleosides: a rapid access to

biological molecules through nitrosocarbonyl chemistry. *Tetrahedron* **2004**, 60, (16), 3643-3651.

- 119. Ranganathan, S.; George, K. S., 2-Aza-3-oxabicyclo[2.2.1]heptene hydrochloride: An exceptionally versatile synthon for carbocyclic sugars and nucleosides. *Tetrahedron* **1997**, **5**3, (9), 3347-3362.
- 120. Lin, C. C.; Wang, Y. C.; Hsu, J. L.; Chiang, C. C.; Su, D. W.; Yan, T. H., Novel Asymmetric Route to Carbanucleoside and Prostanoid Intermediates: Efficient Preparation of Both Optical Antipodes of Chiral Cyclopentenone. *J. Org. Chem.* 1997, 62, (12), 3806-3807.
- 121. Cowart, M.; Bennett, M. J.; Kerwin, J. F., Synthesis of Novel Carbocyclic Adenosine Analogues as Inhibitors of Adenosine Kinase. *J. Org. Chem.* **1999**, 64, (7), 2240-2249.
- 122. Bigeault, J.; Giordano, L.; deRiggi, I.; Gimbert, Y.; Buono, G., Platinum(II)-Coordinated Phosphinous Acid-Catalyzed Alkylidenecyclopropanation of Bicyclic Alkenes with Terminal Alkynes. *Org. Lett.* **2007**, *9*, (18), 3567-3570.
- 123. Calvet, G.; Blanchard, N.; Kouklovsky, C., Domino Metathesis of 3,6-Dihydro-1,2-oxazine: Access to Isoxazolo[2,3-a]pyridin-7-ones. *Org. Lett.* **2007**, 9, (8), 1485-1488.
- 124. Ellis, J. M.; King, S. B., Ring-opening cross metathesis of 1,3-cyclopentadieneheterodienophile cycloadducts to produce cyclic hydrazines and hydroxylamines. *Tetrahedron Lett.* **2002**, 43, (33), 5833-5835.
- 125. Freer, A. A.; Islam, M. A.; Kirby, G. W.; Mahajan, M. P., Cycloaddition of acylnitroso compounds and nitrosobenzene with the pyrrolidine dienamine of Pummerer's ketone: x-ray crystal structure of a bridged 1,2-oxazine derivative. *J. Chem. Soc., Perkin Trans. 1* 1991, (5), 1001-1007.
- 126. Kirby, G. W.; Mackinnon, J. W. M., Cycloadducts of C-nitrosocarbonyl compounds and ergosteryl acetate; [3,3]sigmatropic rearrangements of N-aroyl-3,6-dihydro-1,2-oxazines. J. Chem. Soc., Perkin Trans. 1 1985, (4), 887-892.
- Jeong, L. S.; Lee, J. A., Recent advances in the synthesis of the carbocyclic nucleosides as potential antiviral agents. *Antiviral Chem. Chemother.* 2004, 15, (5), 235-250.
- 128. Rodriguez, J. B.; Comin, M. J., New Progresses in the Enantioselective Synthesis and Biological Properties of Carbocyclic Nucleosides. *Mini Reviews in Medicinal Chemistry* **2003**, *3*, (2), 95-114.
- 129. Roy, A.; Schneller, S. W., 4'- and 1'-Methyl-Substituted 5'-Norcarbanucleosides. *J. Org. Chem.* **2003**, 68, (24), 9269-9273.

- 130. Schneller, S. W., Carbocyclic Nucleosides (Carbanucleosides) as New Therapeutic Leads. *Current Topics in Medicinal Chemistry* **2002**, 2, (10), 1087-1092.
- 131. Hansel, J.-G.; O'Hogan, S.; Lensky, S.; Ritter, A. R.; Miller, M. J., Oxazoline formation via a palladium-catalyzed cyclization: a direct, stereoselective approach to cis-5-amino-2-cyclopenten-1-ol derivatives. *Tetrahedron Lett.* **1995**, 36, (17), 2913-2916.
- 132. De Clercq, E., John Montgomery's Legacy: Carbocyclic Adenosine Analogues as Sah Hydrolase Inhibitors with Broad-Spectrum Antiviral Activity. *Nucleosides, Nucleotides Nucleic Acids* 2005, 24, (10-12), 1395-1415.
- 133. Shireman, B. T.; Miller, M. J., Rapid syntheses of either enantiomer of important carbocyclic nucleoside precursors. *Tetrahedron Lett.* **2000**, 41, (49), 9537-9540.
- 134. Mulvihill, M. J.; Miller, M. J., Syntheses of novel hydroxylamine carbanucleosides. *Tetrahedron* **1998**, 54, (24), 6605-6626.
- 135. Zhang, D.; Ghosh, A.; Suling, C.; Miller, M. J., Efficient functionalization of acylnitroso cycloadducts: Application to the syntheses of carbocyclic nucleoside precursors. *Tetrahedron Lett.* **1996**, 37, (22), 3799-3802.
- 136. Ghosh, A.; Ritter, A. R.; Miller, M. J., Synthesis of enantiomerically pure 5'-azanoraristeromycin analogs. *J. Org. Chem.* **1995**, 60, (18), 5808-5813.
- 137. Jiang, M. X.-W.; Jin, B.; Gage, J. L.; Priour, A.; Savela, G.; Miller, M. J., Substrate-Dependent Dihydroxylation of Substituted Cyclopentenes: Toward the Syntheses of Carbocyclic Sinefungin and Noraristeromycin. J. Org. Chem. 2006, 71, (11), 4164-4169.
- Li, F.; Brogan, J. B.; Gage, J. L.; Zhang, D.; Miller, M. J., Chemoenzymatic synthesis and synthetic application of enantiopure aminocyclopentenols: total synthesis of carbocyclic (+)-uracil polyoxin C and its alpha-epimer. *J. Org. Chem.* 2004, 69, (13), 4538-4540.
- 139. Kim, K.-H.; Miller, M. J., An enantioselective synthesis of the cyclopentene fragment of nucleoside Q. *Tetrahedron Lett.* **2003**, 44, (24), 4571-4573.
- 140. Jean-Bernard Behr, A. D. N. M. J. S., (±)-4-Amino-4,5-dideoxyribose, (±)-4amino-4-deoxyerythrose, and (±)-dihydroxyproline derivatives from <I>N</I>dienyl-?-lactams. *Helv. Chim. Acta* **1995**, 78, (5), 1166-1177.
- 141. Joubert, M.; Defoin, A.; Tarnus, C.; Streith, J., Synthesis of potent a-Dmannosidase and a-L-fucosidase inhibitors: 4-amino-4-deoxy-D-erythrose and 4amino-4,5-dideoxy-L-lyxose. *Synlett* **2000**, (9), 1366-1368.

- 142. Defoin, A.; Pires, J.; Streith, J., Synthesis of aminoerythrose and of aminoerythritol derivatives via Diels-Alder cycloadditions of 1-tertbutyldimethylsiloxybutadiene with acylnitrose dienophiles. *Synlett* **1990**, (2), 111-113.
- 143. Robinson, R., Synthesis of tropinone. J. Chem. Soc. 1917, 111, 762-768.
- 144. Iida, H.; Watanabe, Y.; Kibayashi, C., A New Synthetic Route to Tropane Alkaloids Based on [4+2] Nitroso Cycloaddition to 1,3-Cycloheptadienes. J. Org. Chem. **1985**, 50, 1818-1825.
- 145. Aoyagi, S.; Tanaka, R.; Naruse, M.; Kibayashi, C., Total Synthesis of (-)-Epibatidine Using an Asymmetric Diels-Alder Reaction with a Chiral N-Acylnitroso Dienophile. J. Org. Chem. **1998**, 63, (23), 8397-8406.
- 146. Bathgate, A.; Malpass, J. R., A simple approach to nortropane and nortrop-6-ene derivatives. *Tetrahedron Lett.* **1987**, 28, (47), 5937-5940.
- 147. Smith, C. R.; Justice, D.; Malpass, J. R., Synthetic approaches to homotropanes and homotrop-7-enes using intramolecular displacement and amidomercuration strategies. *Tetrahedron* **1993**, 49, (47), 11037-11054.
- 148. Malpass, J. R.; Hemmings, D. A.; Wallis, A. L.; Fletcher, S. R.; Patel, S., Synthesis and nicotinic acetylcholine-binding properties of epibatidine homologues: homoepibatidine and dihomoepibatidine. *J. Chem. Soc., Perkin Trans. 1* 2001, (9), 1044-1050.
- 149. Malpass, J. R.; Smith, C., A simple approach to homotropanes and homotrop-7enes. *Tetrahedron Lett.* **1992**, 33, (2), 273-276.
- 150. Justice, D. E.; Malpass, J. R., Total synthesis of scopine, pseudoscopine, and norderivatives. *Tetrahedron* **1996**, 52, (36), 11977-11994.
- 151. Groetzl, B.; Handa, S.; Malpass, J. R., The first syntheses of 6,7-dihydroxylated calystegines and homocalystegines. *Tetrahedron Lett.* **2006**, 47, (51), 9147-9150.
- 152. Soulie, J.; Betzer, J.-F.; Muller, B.; Lallemand, J.-Y., General access to polyhydroxylated nortropane derivatives through hetero diels -alder cycloaddition. *Tetrahedron Lett.* **1995**, 36, (52), 9485-9488.
- 153. Martin, S. F., In *The Alkaloids*, Brossi, A. R., Ed. Academic Press: New York, 1987; Vol. 31, p 252.
- 154. Hudlicky, T.; Rinner, U.; Gonzalez, D.; Akgun, H.; Schilling, S.; Siengalewicz, P.; Martinot, T. A.; Pettit, G. R., Total Synthesis and Biological Evaluation of Amaryllidaceae Alkaloids: Narciclasine, ent-7-Deoxypancratistatin, Regioisomer of 7-Deoxypancratistatin, 10b-epi-Deoxypancratistatin, and Truncated Derivatives. J. Org. Chem. 2002, 67, (25), 8726-8743.

- 155. Gonzalez, D.; Martinot, T.; Hudlicky, T., A short chemoenzymatic synthesis of (+)-narciclasine. *Tetrahedron Lett.* **1999**, 40, (16), 3077-3080.
- 156. Hudlicky, T.; Olivo, H. F., A short synthesis of (+)-lycoricidine. J. Am. Chem. Soc. **1992**, 114, (24), 9694-9696.
- 157. Martin, S. F.; Tso, H. H., Synthetic studies on the narciclasine alkaloids. A synthesis of (+-)-lycoricidine. *Heterocycles* **1993**, 35, (1), 85-88.
- 158. Shukla, K. H.; Boehmler, D. J.; Bogacyzk, S.; Duvall, B. R.; Peterson, W. A.; McElroy, W. T.; DeShong, P., Application of Palladium-Catalyzed Allylic Arylation to the Synthesis of a (&#177)-7-Deoxypancratistatin Analogue. *Org. Lett.* **2006**, 8, (19), 4183-4186.
- 159. Olivo, H. F.; Hemenway, M. S.; Hartwig, A. C.; Chan, R., New preparation of activated 2-vinylaziridines from 1,4-amino alcohols. *Synlett* **1998**, (3), 247-248.
- 160. Li, F.; Warshakoon, N. C.; Miller, M. J., Synthetic Application of Acylnitroso Diels-Alder Derived Aminocyclopentenols: Total Synthesis of (+)-Streptazolin. J. Org. Chem. 2004, 69, (25), 8836-8841.
- 161. Li, F.; Miller, M. J., Stereoselective Total Synthesis of (+)-Streptazolin by Using a Temporary Silicon-Tethered RCM Strategy. J. Org. Chem. 2006, 71, (14), 5221-5227.
- 162. Nora, G. P.; Miller, M. J.; Moellmann, U., The synthesis and in vitro testing of structurally novel antibiotics derived from acylnitroso Diels-Alder adducts. *Bioorg. Med. Chem. Lett.* **2006**, 16, (15), 3966-3970.
- 163. Ding, P.; Miller, M. J.; Chen, Y.; Helquist, P.; Oliver, A. J.; Wiest, O., Syntheses of Conformationally Constricted Molecules as Potential NAALADase/PSMA Inhibitors. *Org. Lett.* **2004**, 6, (11), 1805-1808.
- 164. Surman, M. D.; Mulvihill, M. J.; Miller, M. J., Novel a-substituted b-amino diesters from acylnitroso-derived hetero-Diels-Alder cycloadducts. *Tetrahedron Lett.* **2002**, 43, (7), 1131-1134.
- 165. Heinz, L. J.; Lunn, W. H. W.; Murff, R. E.; Paschal, J. W.; Spangle, L. A., An Efficient Synthesis of cis- and trans-Methyl 3-hydroxy-2-pyrrolidone-5-carboxylates, Key Intermediates for the Synthesis of g-Substituted Glutamic Acid Analogs. J. Org. Chem. 1996, 61, (14), 4838-4841.
- Ritter, A. R.; Miller, M. J., Asymmetric Syntheses of Novel Amino Acids and Peptides from Acylnitroso-Derived Cycloadducts. *Tetrahedron Lett.* 1994, 35, (50), 9379-9382.
- 167. Shireman, B. T.; Miller, M. J.; Jonas, M.; Wiest, O., Conformational Study and Enantioselective, Regiospecific Syntheses of Novel Aminoxy trans-Proline

Analogues Derived from an Acylnitroso Diels-Alder Cycloaddition. J. Org. Chem. 2001, 66, (18), 6046-6056.

- 168. Jiang, M. X.-W.; Warshakoon, N. C.; Miller, M. J., Chemoenzymatic Asymmetric Total Synthesis of Phosphodiesterase Inhibitors: Preparation of a Polycyclic Pyrazolo[3,4-d]pyrimidine from an Acylnitroso Diels-Alder Cycloadduct-Derived Aminocyclopentenol. J. Org. Chem. 2005, 70, (7), 2824-2827.
- 169. Lee, W.; Miller, M. J., Concise Synthesis of 4-Acylamino Analogues of 2-Aminobicyclo[3.1.0]hexane-2,6-dicarboxylic Acids (LY354740) from an Acylnitroso Diels-Alder Cycloadduct. J. Org. Chem. **2004**, 69, (13), 4516-4519.
- 170. Mineno, T.; Miller, M. J., Stereoselective Total Synthesis of Racemic BCX-1812 (RWJ-270201) for the Development of Neuraminidase Inhibitors as Antiinfluenza Agents. J. Org. Chem. 2003, 68, (17), 6591-6596.
- 171. Surman, M. D.; Miller, M. J., Synthesis of a Conformationally Restricted Substrate Analogue of Siderophore Biosynthetases. *Org. Lett.* **2001**, *3*, (4), 519-521.
- 172. Shireman, B. T.; Miller, M. J., Synthesis of Enantiomerically and Diastereomerically Pure 2(S)-Amino-6(R)-hydroxy-1,7-heptanedioic Acid Dimethyl Ester Hydrochloride from Cycloheptadiene. J. Org. Chem. 2001, 66, (14), 4809-4813.
- 173. Horsewood, P.; Kirby, G. W.; Sharma, R. P.; Sweeny, J. G., Formation and thermal cleavage reactions of the cycloadduct of 9,10-dimethylanthracene and nitrosyl cyanide. *J. Chem. Soc., Perkin Trans. 1* **1981**, (6), 1802-1806.
- 174. Kirsch, G.; Golde, R.; Neef, G., A cycloaddition route to 14-hydroxysteroids. *Tetrahedron Lett.* **1989**, 30, (34), 4497-4500.
- 175. Perez-Medrano, A.; Grieco, P. A., Synthesis of the highly oxygenated ergostane type steroid (+)-withanolide E. J. Am. Chem. Soc. **1991**, 113, (3), 1057-1059.
- Li, F.; Yang, B.; Miller, M. J.; Zajicek, J.; Noll, B. C.; Mollmann, U.; Dahse, H. M.; Miller, P. A., Iminonitroso Diels-Alder Reactions for Efficient Derivatization and Functionalization of Complex Diene-Containing Natural Products. *Org. Lett.* 2007, 9, (15), 2923-2926.
- Gourlay, R. I.; Kirby, G. W., Short synthesis of 14b-acylaminocodeinones from the cycloadducts of thebaine and acylnitroso compounds. *J. Chem. Res., Synop.* 1997, (5), 152-153.
- 178. Sheldrake, G. N.; Soissons, N., Selective Opening of Ring C in the Morphine Skeleton by an Unexpected Cleavage of the C5-C6 Bond in Cycloadducts of Thebaine and Acyl Nitroso Compounds. J. Org. Chem. **2006**, 71, (2), 789-791.

- 179. McAuley, B. J.; Nieuwenhuyzen, M.; Sheldrake, G. N., An Unusual Samarium Diiodide Mediated Reductive Ring Contraction of a Tricyclic Oxazine to a Highly-Functionalized Cyclopentane and Cyclobutane. *Org. Lett.* **2000**, *2*, (10), 1457-1459.
- Keck, G. E., Intramolecular dienophile transfer. Diels-alder adducts of acylnitroso compounds as useful for intramolecular cycloadditions. *Tetrahedron Lett.* 1978, 19, (48), 4767.
- 181. Kibayashi, C.; Aoyagi, S., Nitrogenous natural products synthesis via N-acylnitroso Diels-Alder methodology. *Synlett* **1995**, (9), 873-879.
- 182. Watanabe, Y.; Iida, H.; Kibayashi, C., Total synthesis of (+-.)-dihydropinidine, (+-.)-monomorine I, and (+-.)-indolizidine 223AB (gephyrotoxin 223AB) by intramolecular nitroso Diels-Alder reaction. *J. Org. Chem.* **1989**, 54, (17), 4088-4097.
- 183. Aoyagi, S.; Shishido, Y.; Kibayashi, C., Total synthesis of (-)-nupharamine and (+)-3-epinupharamine via asymmetric nitroso Diels-Alder reaction. *Tetrahedron Lett.* **1991**, 32, (34), 4325-4328.
- 184. Sato, T.; Aoyagi, S.; Kibayashi, C., Enantioselective Total Synthesis of (+)-Azimine and (+)-Carpaine. *Org. Lett.* **2003**, *5*, (21), 3839-3842.
- 185. Ozawa, T.; Aoyagi, S.; Kibayashi, C., Total Synthesis of the Marine Alkaloids (-)-Lepadins A, B, and C Based on Stereocontrolled Intramolecular Acylnitroso-Diels-Alder Reaction. J. Org. Chem. 2001, 66, (10), 3338-3347.
- 186. Ozawa, T.; Aoyagi, S.; Kibayashi, C., Total Synthesis of the Marine Alkaloid (-)-Lepadin B. *Org. Lett.* **2000**, *2*, (19), 2955-2958.
- 187. Naruse, M.; Aoyagi, S.; Kibayashi, C., Total synthesis of (-)-pumiliotoxin C by aqueous intramolecular acylnitroso Diels-Alder approach. *Tetrahedron Lett.* **1994**, 35, (49), 9213-9216.
- 188. Aoyagi, S.; Hirashima, S.; Saito, K.; Kibayashi, C., Convergent Approach to Pumiliotoxin Alkaloids. Asymmetric Total Synthesis of (+)-Pumiliotoxins A, B, and 225F. J. Org. Chem. 2002, 67, (16), 5517-5526.
- 189. Kibayashi, C.; Aoyagi, S., Development of new synthetic methods and strategies for the total syntheses of pumiliotoxin A class alkaloids. *Yuki Gosei Kagaku Kyokaishi* **1999**, 57, (11), 981-992.
- 190. Naruse, M.; Aoyagi, S.; Kibayashi, C., Stereoselective total synthesis of (-)pumiliotoxin C by an aqueous intramolecular acylnitroso Diels-Alder approach. J. *Chem. Soc., Perkin Trans. 1* **1996**, (11), 1113-1124.
- 191. Liddell, J. R., Pyrrolizidine alkaloids. Nat. Prod. Rep. 2002, 19, (6), 773-781.

- 192. Keck, G. E.; Romer, D. R., An acylnitroso cycloaddition approach to the synthesis of highly oxygenated indolizidine alkaloids. *J. Org. Chem.* **1993**, 58, (22), 6083-6089.
- 193. Naruse, M.; Aoyagi, S.; Kibayashi, C., New Chiral Route to (-)-Swainsonine via an Aqueous Acylnitroso Cycloaddition Approach. J. Org. Chem. **1994**, 59, (6), 1358-1364.
- 194. Abe, H.; Aoyagi, S.; Kibayashi, C., Total synthesis of the proposed structure of lepadiformine via intramolecular N-acylnitroso Diels-Alder reaction. *Tetrahedron Lett.* **2000**, 41, (8), 1205-1208.
- 195. Keck, G. E.; Nickell, D. G., Synthetic studies on pyrrolizidine alkaloids. 1. (+-)-Heliotridine and (+-)-retronecine via intramolecular dienophile transfer. *J. Am. Chem. Soc.* **1980**, 102, (10), 3632-4.
- 196. Iida, H.; Watanabe, Y.; Kibayashi, C., Stereospecific total synthesis of (±)-gephyrotoxin 223AB. J. Am. Chem. Soc. **1985**, 107, (19), 5534-5535.
- 197. Yamazaki, N.; Kibayashi, C., Asymmetric synthesis with a,bbis[(methoxymethyl)oxy] ketones. Enantioselective total synthesis of natural (+)indolizidine 195B (bicyclic gephyrotoxin 195B) and (-)-pinidine and their enantiomers from a common chiral synthon. J. Am. Chem. Soc. **1989**, 111, (4), 1396-408.
- 198. Machinaga, N.; Kibayashi, C., Total synthesis of (-)-indolizidine 239CD. J. Chem. Soc., Chem. Commun. 1991, (6), 405-407.
- 199. Shishido, Y.; Kibayashi, C., The total synthesis of (-)-indolizidines 205A and 235B. J. Chem. Soc., Chem. Commun. 1991, (18), 1237-1239.
- 200. Machinaga, N.; Kibayashi, C., General entry to the 3,5-disubstituted indolizidine class of dendrobatid alkaloids. Total syntheses of both enantiomers of indolizidines 195B, 223AB, 239AB, and 239CD from a common chiral synthon. *J. Org. Chem.* **1992**, 57, (19), 5178-5189.
- Shishido, Y.; Kibayashi, C., Enantiogenic total syntheses of (-)-indolizidines (bicyclic gephyrotoxins) 205A, 207A, 209B, and 235B via the intramolecular Diels-Alder reaction of a chiral N-acylnitroso compound. *J. Org. Chem.* 1992, 57, (10), 2876-2883.
- 202. Yamazaki, N.; Ito, T.; Kibayashi, C., Asymmetric Synthesis of (-)-Indolizidines 167B and 209D Based on Stereocontrolled Allylation of a Chiral Tricyclic N-Acyl-N,O-acetal. *Org. Lett.* **2000**, *2*, (4), 465-467.
- 203. Blakemore, P. R.; Kim, S.-K.; Schulze, V. K.; White, J. D.; Yokochi, A. F. T., Asymmetric synthesis of (+)-loline, a pyrrolizidine alkaloid from rye grass and tall fescue. *J. Chem. Soc., Perkin Trans. 1* **2001**, (15), 1831-1847.

- 204. Blakemore, P. R.; Schulze, V. K.; White, J. D., Asymmetric synthesis of (+)loline. *Chem. Commun.* **2000**, (14), 1263-1264.
- 205. Zhu, L.; Lauchli, R.; Loo, M.; Shea, K. J., An Approach to the Synthesis of Stenine. *Org. Lett.* **2007**, *9*, (12), 2269-2271.
- 206. Molina, C. L.; Chow, C. P.; Shea, K. J., Type 2 Intramolecular N-Acylazo Diels-Alder Reaction: Regio- and Stereoselective Synthesis of Bridgehead Bicyclic 1,2-Diazines. J. Org. Chem. 2007, 72, (18), 6816-6823.
- 207. Sparks, S. M.; Chow, C. P.; Zhu, L.; Shea, K. J., Type 2 Intramolecular N-Acylnitroso Diels-Alder Reaction: Scope and Application to the Synthesis of Medium Ring Lactams. J. Org. Chem. **2004**, 69, (9), 3025-3035.
- 208. Chow, C. P.; Shea, K. J.; Sparks, S. M., Type 2 Intramolecular N-Acylnitroso Diels-Alder Reaction: Stereoselective Synthesis of Bridged Bicyclic Oxazinolactams. *Org. Lett.* **2002**, 4, (16), 2637-2640.
- Sparks, S. M.; Vargas, J. D.; Shea, K. J., Type 2 intramolecular nitroso Diels-Alder reaction. Synthesis and structure of bridgehead oxazinolactams. *Org. Lett.* 2000, 2, (10), 1473-1475.
- 210. The, L., Stopping tuberculosis proves hard to do. *Lancet* **2007**, 369, (9566), 965.
- 211. Check, E., After decades of drought, new drug possibilities flood TB pipeline. *Nat. Med.* **2007**, 13, (3), 266.
- 212. WHO Factsheet no. 104. http://www.who.int/mediacentre/factsheets/fs104/en/ (Feb. 24),
- 213. Friedman, L. N.; Williams, M. T.; Singh, T. P.; Frieden, T. R., Tuberculosis, AIDS, and Death among Substance Abusers on Welfare in New York City. *N. Engl. J. Med.* **1996**, 334, (13), 828-833.
- 214. Kaufmann, S. H. E., Tuberculosis and AIDS a devilish liaison. *Drug Discovery Today* **2007**, 12, (21-22), 891-893.
- 215. Guo, H.; Naser, S. A.; Ghobrial, G.; Phanstiel, O. I. V., Synthesis and biological evaluation of new citrate-based siderophores as potential probes for the mechanism of iron uptake in Mycobacteria. *J. Med. Chem.* **2002**, 45, (10), 2056-2063.
- 216. Brennan, P. J.; Nikaido, H., The Envelope of Mycobacteria. *Annu. Rev. Biochem.* **1995,** 64, (1), 29-63.
- 217. Kaufmann, S. H. E.; McMichael, A. J., Annulling a dangerous liaison: vaccination strategies against AIDS and tuberculosis. *Nat. Med.* **2005**, 11, (4s), S33-S44.

- 218. Bastian, I.; Colebunders, R., Treatment and Prevention of Multidrug-Resistant Tuberculosis. *Drugs* **1999**, 58, (4), 633-661.
- 219. Dorman, S. E.; Chaisson, R. E., From magic bullets back to the Magic Mountain: the rise of extensively drug-resistant tuberculosis. *Nat. Med.* **2007**, 13, (3), 295-298.
- 220. Schatz, A.; Bugie, E.; Waksman, S. A., Streptomycin, a substance exhibiting antibiotic activity against gram-positive and gram-negative bacteria. *Proc. Soc. Exp. Biol. Med.* **1944**, 55, 66-69.
- 221. Hinshaw, H. C.; Feldman, W. H.; Pfuetze, K. H., Treatment of tuberculosis with streptomycin. *JAMA* **1946**, 132, 778-782.
- 222. Snider, D. E.; Castro, K. G., The Global Threat of Drug-Resistant Tuberculosis. *N. Engl. J. Med.* **1998**, 338, (23), 1689-1690.
- 223. Ballell, L.; Field, R. A.; Duncan, K.; Young, R. J., New Small-Molecule Synthetic Antimycobacterials. *Antimicrob. Agents Chemother.* 2005, 49, (6), 2153-2163.
- 224. The, L., XDR-TB--a global threat. *Lancet* **2006**, 368, (9540), 964.
- 225. Mandavilli, A., Virtually incurable TB warns of impending disaster. *Nat. Med.* **2007**, 13, (3), 271.
- 226. Raviglione, M. C.; Uplekar, M. W., WHO's new Stop TB Strategy. *Lancet* 2006, 367, (9514), 952.
- 227. Raviglione, M. C., The new Stop TB Strategy and the Global Plan to Stop TB, 2006-2015. *Bulletin of the World Health Organization* **2007**, 85, 327-327.
- Maher, D.; Dye, C.; Floyd, K.; Pantoja, A.; Lonnroth, K.; Reid, A.; Nathanson, E.; Pennas, T.; Fruth, U.; Cunningham, J.; Ignatius, H.; Raviglione, M. C.; Koek, I.; Espinal, M., Planning to improve global health: the next decade of tuberculosis control. *Bulletin of the World Health Organization* 2007, 85, 341-347.
- 229. Harper, C., Tuberculosis, a neglected opportunity? *Nat Med* **2007**, 13, (3), 309-312.
- 230. Quadri, L. E. N.; Sello, J.; Keating, T. A.; Weinreb, P. H.; Walsh, C. T., Identification of a Mycobacterium tuberculosis gene cluster encoding the biosynthetic enzymes for assembly of the virulence-conferring siderophore mycobactin. *Chem. Biol.* **1998**, 5, (11), 631-645.
- 231. Barry, C. E.; Slayden, R. A.; Sampson, A. E.; Lee, R. E., Use of genomics and combinatorial chemistry in the development of new antimycobacterial drugs. *Biochem. Pharmacol.* **1999**, 59, (3), 221-231.

- 232. Young, D. B., Blueprint for the white plague. *Nature* **1998**, 393, (6685), 515-516.
- Cole, S. T.; Brosch, R.; Parkhill, J.; Garnier, T.; Churcher, C.; Harris, D.; Gordon, S. V.; Eiglmeier, K.; Gas, S.; Barry, C. E.; Tekaia, F.; Badcock, K.; Basham, D.; Brown, D.; Chillingworth, T.; Connor, R.; Davies, R.; Devlin, K.; Feltwell, T.; Gentles, S.; Hamlin, N.; Holroyd, S.; Hornsby, T.; Jagels, K.; Krogh, A.; McLean, J.; Moule, S.; Murphy, L.; Oliver, K.; Osborne, J.; Quail, M. A.; Rajandream, M. A.; Rogers, J.; Rutter, S.; Seeger, K.; Skelton, J.; Squares, R.; Squares, S.; Sulston, J. E.; Taylor, K.; Whitehead, S.; Barrell, B. G., Deciphering the biology of Mycobacterium tuberculosis from the complete genome sequence. *Nature* 1998, 393, (6685), 537-544.
- 234. Barry, C. E., New horizons in the treatment of tuberculosis. *Biochem. Pharmacol.* **1997**, 54, (11), 1165-1172.
- 235. Sandra M. Newton, C. L. C. W. W., A review of antimycobacterial natural products. *Phytotherapy Research* **2000**, 14, (5), 303-322.
- 236. Copp, B. R., Antimycobacterial natural products. *Nat. Prod. Rep.* **2003**, 20, (6), 535-557.
- 237. Stover, C. K.; Warrener, P.; VanDevanter, D. R.; Sherman, D. R.; Arain, T. M.; Langhorne, M. H.; Anderson, S. W.; Towell, J. A.; Yuan, Y.; McMurray, D. N.; Kreiswirth, B. N.; Barry, C. E.; Baker, W. R., A small-molecule nitroimidazopyran drug candidate for the treatment of tuberculosis. *Nature* 2000, 405, (6789), 962-966.
- 238. Andries, K.; Verhasselt, P.; Guillemont, J.; Goehlmann, H. W. H.; Neefs, J.-M.; Winkler, H.; Van Gestel, J.; Timmerman, P.; Zhu, M.; Lee, E.; Williams, P.; de Chaffoy, D.; Huitric, E.; Hoffner, S.; Cambau, E.; Truffot-Pernot, C.; Lounis, N.; Jarlier, V., A Diarylquinoline Drug Active on the ATP Synthase of Mycobacterium tuberculosis. *Science* 2005, 307, (5707), 223-227.
- 239. Biava, M.; Porretta, G. C.; Poce, G.; Supino, S.; Deidda, D.; Pompei, R.; Molicotti, P.; Manetti, F.; Botta, M., Antimycobacterial Agents. Novel Diarylpyrrole Derivatives of BM212 Endowed with High Activity toward Mycobacterium tuberculosis and Low Cytotoxicity. *J. Med. Chem.* 2006, 49, (16), 4946-4952.
- 240. Biava, M., BM 212 and its Derivatives as a New Class of Antimycobacterial Active Agents. *Curr. Med. Chem.* **2002**, 9, (21), 1859-1869.
- 241. Metaferia, B. B.; Ray, S.; Smith, J. A.; Bewley, C. A., Design and synthesis of substrate-mimic inhibitors of mycothiol-S-conjugate amidase from Mycobacterium tuberculosis. *Bioorg. Med. Chem. Lett.* **2007**, 17, (2), 444-447.
- 242. Kuo, M. R.; Morbidoni, H. R.; Alland, D.; Sneddon, S. F.; Gourlie, B. B.; Staveski, M. M.; Leonard, M.; Gregory, J. S.; Janjigian, A. D.; Yee, C.; Musser, J.

M.; Kreiswirth, B.; Iwamoto, H.; Perozzo, R.; Jacobs, W. R., Jr.; Sacchettini, J. C.; Fidock, D. A., Targeting Tuberculosis and Malaria through Inhibition of Enoyl Reductase: COMPOUND ACTIVITY AND STRUCTURAL DATA. *J. Biol. Chem.* **2003**, 278, (23), 20851-20859.

- 243. Ma, Y.; Stern, R. J.; Scherman, M. S.; Vissa, V. D.; Yan, W.; Jones, V. C.; Zhang, F.; Franzblau, S. G.; Lewis, W. H.; McNeil, M. R., Drug Targeting Mycobacterium tuberculosis Cell Wall Synthesis: Genetics of dTDP-Rhamnose Synthetic Enzymes and Development of a Microtiter Plate-Based Screen for Inhibitors of Conversion of dTDP-Glucose to dTDP-Rhamnose. *Antimicrob. Agents Chemother.* 2001, 45, (5), 1407-1416.
- 244. Andres, C. J.; Bronson, J. J.; D'Andrea, S. V.; Deshpande, M. S.; Falk, P. J.; Grant-Young, K. A.; Harte, W. E.; Ho, H.-T.; Misco, P. F.; Robertson, J. G.; Stock, D.; Sun, Y.; Walsh, A. W., 4-Thiazolidinones: novel inhibitors of the bacterial enzyme murB. *Bioorg. Med. Chem. Lett.* **2000**, 10, (8), 715-717.
- 245. Miethke, M.; Bisseret, P.; Beckering Carsten, L.; Vignard, D.; Eustache, J.; Marahiel Mohamed, A., Inhibition of aryl acid adenylation domains involved in bacterial siderophore synthesis. *The FEBS journal* **2006**, 273, (2), 409-419.
- 246. Ferreras, J. A.; Ryu, J.-S.; Di Lello, F.; Tan, D. S.; Quadri, L. E. N., Smallmolecule inhibition of siderophore biosynthesis in Mycobacterium tuberculosis and Yersinia pestis. *Nat. Chem. Bio.* **2005**, 1, (1), 29-32.
- Somu, R. V.; Boshoff, H.; Qiao, C.; Bennett, E. M.; Barry, C. E., III; Aldrich, C. C., Rationally Designed Nucleoside Antibiotics That Inhibit Siderophore Biosynthesis of Mycobacterium tuberculosis. J. Med. Chem. 2006, 49, (1), 31-34.
- 248. Ratledge, C., Iron, mycobacteria and tuberculosis. *Tuberculosis* **2004**, 84, (1-2), 110-130.
- 249. De Voss, J. J.; Rutter, K.; Schroeder, B. G.; Barry, C. E., III, Iron Acquisition and Metabolism by Mycobacteria. *J. Bacteriol.* **1999**, 181, (15), 4443-4451.
- 250. Ratledge, C.; Dover, L. G., Iron Metabolism in Pathogenic Bacteria. *Annu. Rev. Microbiol.* **2000**, 54, (1), 881-941.
- 251. Boukhalfa, H.; Crumbliss, A. L., Chemical aspects of siderophore mediated iron transport. *BioMetals* **2002**, 15, (4), 325-339.
- 252. Piddington, D. L.; Kashkouli, A.; Buchmeier, N. A., Growth of Mycobacterium tuberculosis in a Defined Medium Is Very Restricted by Acid pH and Mg2+ Levels. *Infect. Immun.* **2000**, 68, (8), 4518-4522.
- 253. Hentze, M. W.; Muckenthaler, M. U.; Andrews, N. C., Balancing Acts: Molecular Control of Mammalian Iron Metabolism. *Cell* **2004**, 117, (3), 285-297.

- 254. Miethke, M.; Marahiel, M. A., Siderophore-Based Iron Acquisition and Pathogen Control. *Microbiol. Mol. Biol. Rev.* **2007**, 71, (3), 413-451.
- 255. Andrews, S. C.; Robinson, A. K.; Rodriguez-Quinones, F., Bacterial iron homeostasis. *FEMS Microbiol. Rev.* 2003, 27, (2-3), 215-237.
- 256. Roosenberg, J. M.; Yun-Ming, L.; Yong, L.; Miller, M. J., Studies and Syntheses of Siderophores, Microbial Iron Chelators, and Analogs as Potential Drug Delivery Agents. *Curr. Med. Chem.* **2000**, *7*, (2), 159-139.
- 257. Vergne, A. F.; Walz, A. J.; Miller, M. J., Iron chelators from mycobacteria (1954-1999) and potential therapeutic applications. *Nat. Prod. Rep.* **2000**, 17, (1), 99-116.
- 258. Neilands, J. B., Siderophores: Structure and Function of Microbial Iron Transport Compounds. J. Biol. Chem. **1995**, 270, (45), 26723-26726.
- 259. Miller, M. J.; Malouin, F., Microbial iron chelators as drug delivery agents: the rational design and synthesis of siderophore-drug conjugates. *Acc. Chem. Res.* **1993**, 26, (5), 241-249.
- 260. Miller, M. J., Syntheses and therapeutic potential of hydroxamic acid based siderophores and analogs. *Chem. Rev.* **1989**, 89, (7), 1563-1579.
- 261. Schwyn, B. B., Universal chemical assay for the detection and determination of siderophores. *Anal. Biochem.* **1987**, 160, (1), 47.
- 262. Lee, B. H.; Miller, M. J.; Prody, C. A.; Neilands, J. B., Artificial siderophores. 1. Synthesis and microbial iron transport capabilities. *J. Med. Chem.* **1985**, 28, (3), 317-323.
- 263. Bergeron, R. J., Synthesis and solution structure of microbial siderophores. *Chem. Rev.* **1984**, 84, (6), 587-602.
- 264. Raymond, K. N.; Dertz, E. A.; Kim, S. S., Enterobactin: An archetype for microbial iron transport. *Proc. Natl. Acad. Sci. U. S. A.* **2003**, 100, (7), 3584-3588.
- 265. Gardner, R. A.; Kinkade, R.; Wang, C.; Phanstiel, O. I. V., Total Synthesis of Petrobactin and Its Homologues as Potential Growth Stimuli for Marinobacter hydrocarbonoclasticus, an Oil-Degrading Bacteria. *J. Org. Chem.* **2004**, 69, (10), 3530-3537.
- 266. Mulqueen, G. C.; Pattenden, G.; Whiting, D. A., Synthesis of the hydroxamate siderophore nannochelin A. *Tetrahedron* **1993**, 49, (40), 9137-9142.

- 267. Bergeron, R. J.; Phanstiel, O. I. V., The total synthesis of nannochelin: a novel cinnamoyl hydroxamate-containing siderophore. *J. Org. Chem.* **1992**, 57, (26), 7140-7143.
- 268. Konetschny-Rapp, S.; Jung, G.; Meiwes, J.; Zaehner, H., Staphyloferrin A: a structurally new siderophore from staphylococci. *Eur. J. Biochem.* **1990**, 191, (1), 65-74.
- 269. Fadeev, E. A.; Luo, M.; Groves, J. T., Synthesis, Structure, and Molecular Dynamics of Gallium Complexes of Schizokinen and the Amphiphilic Siderophore Acinetoferrin. *J. Am. Chem. Soc.* **2004**, 126, (38), 12065-12075.
- 270. Lee, B. H.; Miller, M. J., Natural ferric ionophores: total synthesis of schizokinen, schizokinen A, and arthrobactin. J. Org. Chem. **1983**, 48, (1), 24-31.
- 271. Luo, M.; Fadeev, E. A.; Groves, J. T., Membrane Dynamics of the Amphiphilic Siderophore, Acinetoferrin. J. Am. Chem. Soc. 2005, 127, (6), 1726-1736.
- 272. Gardner, R. A.; Ghobrial, G.; Naser, S. A.; Phanstiel, O., Synthesis and Biological Evaluation of New Acinetoferrin Homologues for Use as Iron Transport Probes in Mycobacteria. *J. Med. Chem.* **2004**, 47, (20), 4933-4940.
- Wang, Q. X.; Phanstiel, O. I. V., Total Synthesis of Acinetoferrin. J. Org. Chem. 1998, 63, (5), 1491-1495.
- 274. Carrano, C. J.; Drechsel, H.; Kaiser, D.; Jung, G.; Matzanke, B.; Winkelmann, G.; Rochel, N.; Albrecht-Gary, A. M., Coordination Chemistry of the Carboxylate Type Siderophore Rhizoferrin: The Iron(III) Complex and Its Metal Analogs. *Inorg. Chem.* **1996**, 35, (22), 6429-6436.
- 275. Okonya, J. F.; Kolasa, T.; Miller, M. J., Synthesis of the Peptide Fragment of Pseudobactin. J. Org. Chem. 1995, 60, (7), 1932-1935.
- 276. Francis, J.; Madinaveitia, J.; Macturk, H. M.; Snow, G. A., Isolation from Acid-Fast Bacteria of a Growth-Factor for Mycobacterium johnei and of a Precursor of Phthiocol. *Nature* **1949**, 163, 365-366.
- 277. Francis, J.; Macturk, H. M.; Madinaveitia, J.; Snow, G. A., Mycobactin, a growth factor for Mycobacterium johnei. I. Isolation from Mycobacterium phlei. *Biochem. J.* **1953**, 55, (4), 596-607.
- 278. Snow, G. A., Mycobactins: iron-chelating growth factors from mycobacteria. *Bacteriol. Rev.* **1970**, 34, (2), 99-125.
- 279. De Voss, J. J.; Rutter, K.; Schroeder, B. G.; Su, H.; Zhu, Y.; Barry, C. E., The salicylate-derived mycobactin siderophores of Mycobacterium tuberculosis are essential for growth in macrophages. *Proc. Natl. Acad. Sci.* **2000**, 97, (3), 1252-1257.

- Gobin, J.; Moore, C. H.; Reeve, J. R., Jr.; Wong, D. K.; Gibson, B. W.; Horwitz, M. A., Iron Acquisition by Mycobacterium Tuberculosis: Isolation and Characterization of a Family of Iron-Binding Exochelins. *Proc. Natl. Acad. Sci.* 1995, 92, (11), 5189-5193.
- 281. Lane, S. J.; Marshall, P. S.; Upton, R. J.; Ratledge, C.; Ewing, M., Novel extracellular mycobactins, the carboxymycobactins from Mycobacterium avium. *Tetrahedron Lett.* **1995**, 36, (23), 4129-4132.
- 282. Lane, S. J.; Marshall, P. S.; Upton, R. J.; Ratledge, C., Isolation and characterization of carboxymycobactins as the second extracellular siderophores in Mycobacterium smegmatis. *BioMetals* **1998**, 11, (1), 13-20.
- 283. Dong, L.; Miller, M. J.; Moellmann, U., Microbial growth promotion studies of exochelin MN and analogues thereof. *BioMetals* **2004**, 17, (2), 99-104.
- 284. Dhungana, S.; Miller, M. J.; Dong, L.; Ratledge, C.; Crumbliss, A. L., Iron chelation properties of an extracellular siderophore exochelin MN. *J. Am. Chem. Soc.* **2003**, 125, (25), 7654-7663.
- 285. Dong, L.; Miller, M. J., Total Synthesis of Exochelin MN and Analogues. J. Org. Chem. 2002, 67, (14), 4759-4770.
- 286. Sharman, G. J.; Williams, D. H.; Ewing, D. F.; Ratledge, C., Isolation, purification and structure of exochelin MS, the extracellular siderophore from Mycobacterium smegmatis. *Biochem. J.* **1995**, 305, (1), 187-196.
- 287. Sharman, G. J.; Williams, D. H.; Ewing, D. F.; Ratledge, C., Determination of the structure of exochelin MN, the extracellular siderophore from Mycobacterium neoaurum. *Chem. Biol.* **1995**, 2, (8), 553-561.
- 288. Breuer, W.; Epsztejn, S.; Cabantchik, Z. I., Iron acquired from transferrin by K562 cells is delivered into a cytoplasmic pool of chelatable iron(II). *J. Biol. Chem.* **1995**, 270, (41), 24209-24215.
- 289. Olakanmi, O.; Schlesinger, L. S.; Ahmed, A.; Britigan, B. E., Intraphagosomal Mycobacterium tuberculosis Acquires Iron from Both Extracellular Transferrin and Intracellular Iron Pools. *J. Biol. Chem.* **2002**, 277, (51), 49727-49734.
- 290. Qian, Z. M.; Tang, P. L.; Wang, Q., Iron crosses the endosomal membrane by a carrier-mediated process. *Progress in Biophysics and Molecular Biology* **1997**, 67, (1), 1-15.
- 291. Rodriguez, G. M.; Smith, I., Mechanisms of iron regulation in mycobacteria: role in physiology and virulence. *Mol. Microbiol.* **2003**, 47, (6), 1485-1494.

- 292. Ratledge, C.; Patel, P. V.; Mundy, J., Iron transport in *Mycobacterium smegmatis*: the location of mycobactin by electron microscopy. *J. Gen. Microbiol.* **1982**, 128, 1559-1565.
- 293. Ghio, A. J.; Piantadosi, C. A.; Crumbliss, A. L., Hypothesis: iron chelation plays a vital role in neutrophilic inflammation. *BioMetals* **1997**, 10, (2), 135-142.
- 294. Bosne-David, S.; Bricard, L.; Ramiandrasoa, F.; DeRoussent, A.; Kunesch, G.; Andremont, A., Evaluation of growth promotion and inhibition from mycobactins and nonmycobacterial siderophores (Desferrioxamine and FR160) in Mycobacterium aurum. *Antimicrob. Agents Chemother.* **1997**, 41, (8), 1837-1839.
- 295. Carpenter, J. G.; Moore, J. W., Synthesis of an analogue of mycobactin. J. Chem. Soc., Perkin Trans. 1 1969, 12, 1610-1611.
- 296. Maurer, P. J.; Miller, M. J., Total synthesis of a mycobactin: mycobactin S2. J. Am. Chem. Soc. 1983, 105, (2), 240-245.
- 297. Hu, J.; Miller, M. J., Total Synthesis of a Mycobactin S, a Siderophore and Growth Promoter of Mycobacterium Smegmatis, and Determination of its Growth Inhibitory Activity against Mycobacterium tuberculosis. *J. Am. Chem. Soc.* **1997**, 119, (15), 3462-3468.
- 298. Xu, Y.; Miller, M. J., Total Syntheses of Mycobactin Analogs as Potent Antimycobacterial Agents Using a Minimal Protecting Group Strategy. J. Org. Chem. 1998, 63, (13), 4314-4322.
- 299. Walz, A. J.; Mollmann, U.; Miller, M. J., Synthesis and studies of catecholcontaining mycobactin S and T analogs. *Org. Biomol. Chem.* **2007**, *5*, (10), 1621-1628.
- 300. Poreddy, A. R.; Schall, O. F.; Marshall, G. R.; Ratledge, C.; Slomczynska, U., Solid-phase synthesis of methyl carboxymycobactin T 7 and analogs as potential antimycobacterial agents. *Bioorg. Med. Chem. Lett.* **2003**, 13, (15), 2553-2556.
- 301. Pahl, P. M. B.; Yan, X.-D.; Hodges, Y. K.; Rosenthal, E. A.; Horwitz, M. A.; Horwitz, L. D., An Exochelin of Mycobacterium tuberculosis Reversibly Arrests Growth of Human Vascular Smooth Muscle Cells in Vitro. J. Biol. Chem. 2000, 275, (23), 17821-17826.
- 302. Pahl, P.; Horwitz, M.; Horwitz, K.; Horwitz, L., Desferri-exochelin induces death by apoptosis in human breast cancer cells but does not kill normal breast cells. *Breast Cancer Res. Treat.* **2001**, 69, (1), 69-79.
- 303. Hodges, Y. K.; Antholine, W. E.; Horwitz, L. D., Effect on ribonucleotide reductase of novel lipophilic iron chelators: the desferri-exochelins. *Biochem. Biophys. Res. Commun.* **2004**, 315, (3), 595-598.
- 304. Yokokawa, F.; Izumi, K.; Omata, J.; Shioiri, T., Total synthesis of Amamistatin A, an antiproliferative linear peptide from an actinomycete. *Tetrahedron* **2000**, 56, (19), 3027-3034.
- 305. Fennell, K. A.; Miller, M. J., Syntheses of Amamistatin Fragments and Determination of Their HDAC and Antitumor Activity. *Org. Lett.* **2007**, *9*, (9), 1683-1685.
- 306. Fennell, K. A.; Mollmann, U.; Miller, M. J., Syntheses and Biological Activity of Amamistatin B and Analogs. *J. Org. Chem.* **2008**, 73, (3), 1018-1024.
- 307. Kokubo, S.; Suenaga, K.; Shinohara, C.; Tsuji, T.; Uemura, D., Structures of amamistatins A and B, novel growth inhibitors of human tumor cell lines from Nocardia asteroides. *Tetrahedron* **2000**, 56, (35), 6435-6440.
- 308. Suenaga, K.; Kokubo, S.; Shinohara, C.; Tsuji, T.; Uemura, D., Structures of amamistatins A and B, novel growth inhibitors of human tumor cell lines from an actinomycete. *Tetrahedron Lett.* **1999**, 40, (10), 1945-1948.
- 309. Murakami, Y.; Kato, S.; Nakajima, M.; Matsuoka, M.; Kawai, H.; Shin-Ya, K.; Seto, H., Formobactin, a novel free radical scavenging and neuronal cell protecting substance from Nocardia sp. *J. Antibiot.* **1996**, 49, (9), 839-845.
- Ratledge, C.; Snow, G. A., Isolation and structure of nocobactin NA, a lipid-soluble iron-binding compound from Nocardia asteroides. *Biochem. J.* 1974, 139, (2), 407-413.
- 311. Nemoto, A.; Hoshino, Y.; Yazawa, K.; Ando, A.; Mikami, Y., Asterobactin, a New Siderophore Group Antibiotic from Nocardia asteroides. *J. Antibiot.* **2002**, 55, 593-597.
- 312. Tsukamoto, M.; Murooka, K.; Nakajima, S.; Abe, S.; Suzuki, H.; Hirano, K.; Kondo, H.; Kojiri, K.; Suda, H., BE-32030 A, B, C, D and E, New Antitumor Substances Produced by Nocavdia sp. A32030. J. Antibiot. 1997, 50, (10), 815-821.
- 313. Ikeda, Y.; Nonaka, H.; Furumai, T.; Onaka, H.; Igarashi, Y., Nocardimicins A, B, C, D, E, and F, Siderophores with Muscarinic M3 Receptor Inhibiting Activity from Nocardia sp. TP-A0674. *J. Nat. Prod.* **2005**, 68, (7), 1061-1065.
- Ikeda, Y.; Furumai, T.; Igarashi, Y., Nocardimicins G, H and I, Siderophores with Muscarinic M3 Receptor Binding Inhibitory Activity from Nocardia nova JCM 6044. J. Antibiot. 2005, 58, (9), 566-572.
- 315. Mitchell, J. M.; Shaw, J. T., Synthesis and Stereochemical Assignment of Brasilibactin A. *Org. Lett.* **2007**, *9*, (9), 1679-1681.

- 316. Tsuda, M.; Yamakawa, M.; Oka, S.; Tanaka, Y.; Hoshino, Y.; Mikami, Y.; Sato, A.; Fujiwara, H.; Ohizumi, Y.; Kobayashi, J., Brasilibactin A, a Cytotoxic Compound from Actinomycete Nocardia brasiliensis. J. Nat. Prod. 2005, 68, (3), 462-464.
- Sonntag, B.; Gerlitz, M.; Paululat, T.; Rasser, H.-F.; Grün-Wollny, I.; Hansske, F. G., Oxachelin, a Novel Iron Chelator and Antifungal Agent from Streptomyces sp. GW9/1258. J. Antibiot. 2006, 59, (10), 659-663.
- 318. Burgess, E. M.; Penton, H. R.; Taylor, E. A., Thermal reactions of alkyl N-carbomethoxysulfamate esters. *J. Org. Chem.* **1973**, 38, (1), 26-31.
- 319. Wipf, P.; Venkatraman, S., An improved protocol for azole synthesis with PEGsupported Burgess reagent. *Tetrahedron Lett.* **1996**, 37, (27), 4659-4662.
- Phillips, A. J.; Uto, Y.; Wipf, P.; Reno, M. J.; Williams, D. R., Synthesis of functionalized oxazolines and oxazoles with DAST and Deoxo-Fluor. *Org. Lett.* 2000, 2, (8), 1165-1168.
- 321. Miller, M. J.; Xu, Y. Preparation of antimycobacterial agents. 2000-5789736310058, 20000525., 2001.
- 322. Miller, M. J.; Hu, J. Preparation of phenyloxazolinecarboxylates and thiazolinecarboxylates as antituberculosis agents. 2000-5788346403623, 20000525., 2002.
- 323. Strazzolini, P.; Pavsler, A., Highly Efficient, Low-Cost, and Simple Protocol for the Preparation of O-(Phenylmethyl)hydroxylamine. *Ind. Eng. Chem. Res.* **2005**, 44, (6), 1625-1626.
- 324. Welch, J. T.; Seper, K. W., Synthesis, regioselective deprotonation, and stereoselective alkylation of fluoro ketimines. *J. Org. Chem.* **1988**, 53, (13), 2991-2999.
- 325. Levin, J. I.; Turos, E.; Weinreb, S. M., An alternative procedure for the aluminum-mediated conversion of esters to amides. *Synth. Commun.* **1982**, 12, (13), 989-993.
- 326. Pirrung, M. C.; Chau, J. H. L., A Convenient Procedure for the Preparation of Amino Acid Hydroxamates from Esters. J. Org. Chem. **1995**, 60, (24), 8084-8085.
- 327. Guzzo, P. R.; Miller, M. J., Catalytic, Asymmetric Synthesis of the Carbacephem Framework. *J. Org. Chem.* **1994**, 59, (17), 4862-4867.
- 328. Lee, B. H.; Biswas, A.; Miller, M. J., [1,2] Anionic Rearrangements of Substituted N-Hydroxy-2-azetidinones and Applications to the Synthesis of Bicyclic B-Lactams. *J. Org. Chem.* **1986**, 51, 106-109.

- 329. Klijn, J. E.; Engberts, J. B. F. N., Organic chemistry: Fast reactions 'on water'. *Nature* **2005**, 435, (7043), 746-747.
- Narayan, S.; Muldoon, J.; Finn, M. G.; Fokin, V. V.; Kolb, H. C.; Sharpless, K. B., "On water": Unique reactivity of organic compounds in aqueous suspension. *Angew. Chem., Int. Ed.* 2005, 44, (21), 3275-3279.
- 331. Outram, H. S.; Raw, S. A.; Taylor, R. J. K., In situ oxidative diol cleavage-Wittig processes. *Tetrahedron Lett.* **2002**, 43, (35), 6185-6187.
- 332. Varma, R. S.; Saini, R. K.; Meshram, H. M., Selective oxidation of sulfides to sulfoxides and sulfones by microwave thermolysis on wet silica-supported sodium periodate. *Tetrahedron Lett.* **1997**, 38, (37), 6525-6528.
- 333. Mancuso, A. J.; Swern, D., Activated Dimethyl Sulfoxide: Useful Reagents for Synthesis. *Synthesis* **1981**, 165-184.
- 334. Marshall, J. A.; Garofalo, A. W., Oxidative cleavage of mono-, di-, and trisubstituted olefins to methyl esters through ozonolysis in methanolic sodium hydroxide. *J. Org. Chem.* **1993**, 58, (14), 3675-3680.
- Lepore, S. D.; Schacht, A. L.; Wiley, M. R., Preparation of 2hydroxybenzamidines from 3-aminobenzisoxazoles. *Tetrahedron Lett.* 2002, 43, (48), 8777-8779.
- 336. Keck, G. E.; Fleming, S.; Nickell, D.; Weider, P., Mild and efficient methods for the reductive cleavage of nitrogen-oxygen bonds. *Synth. Commun.* **1979**, 9, (4), 281-286.
- 337. Staben, S. T.; Linghu, X.; Toste, F. D., Enantioselective Synthesis of gamma-Hydroxyenones by Chiral Base-Catalyzed Kornblum DeLaMare Rearrangement. *J. Am. Chem. Soc.* **2006**, 128, (39), 12658-12659.
- Travis, B. R.; Narayan, R. S.; Borhan, B., Osmium Tetroxide-Promoted Catalytic Oxidative Cleavage of Olefins: An Organometallic Ozonolysis. *J. Am. Chem. Soc.* 2002, 124, (15), 3824-3825.
- 339. Creary, X., Tosylhydrazone salt pyrolyses. Phenyldiazomethanes. *Org. Synth.* **1986**, 64, 207-216.
- 340. Plietker, B.; Niggemann, M., RuCl3/CeCl3/NaIO4: A New Bimetallic Oxidation System for the Mild and Efficient Dihydroxylation of Unreactive Olefins. *J. Org. Chem.* **2005**, 70, (6), 2402-2405.
- Dess, D. B.; Martin, J. C., Readily accessible 12-I-5 oxidant for the conversion of primary and secondary alcohols to aldehydes and ketones. *J. Org. Chem.* 1983, 48, (22), 4155-4156.

- 342. Jones, N. A.; Nepogodiev, S. A.; MacDonald, C. J.; Hughes, D. L.; Field, R. A., Synthesis of the Branched-Chain Sugar Aceric Acid: A Unique Component of the Pectic Polysaccharide Rhamnogalacturonan-II. *J. Org. Chem.* **2005**, 70, (21), 8556-8559.
- 343. Tavares, F.; Lawson, J. P.; Meyers, A. I., Total Synthesis of Streptogramin Antibiotics. (-)-Madumycin II. J. Am. Chem. Soc. **1996**, 118, (13), 3303-3304.
- 344. Sharma, G. V. M.; Reddy, K. L.; Lakshmi, P. S.; Ravi, R.; Kunwar, A. C., Direct C-Allylation of Aryl-Alkyl/Glycosyl Carbinols: Facile Synthesis of C-linked Carbo- $\beta^2$ - $\gamma^2$ -δ<sup>2</sup>-Amino Acids. *J. Org. Chem.* **2006**, 71, (10), 3967-3969.
- 345. Sarabia, F.; Sanchez-Ruiz, A., Total Synthesis of Bengamide E and Analogues by Modification at C-2 and at Terminal Olefinic Positions. *J. Org. Chem.* **2005**, 70, (23), 9514-9520.
- 346. DeClue, M. S.; Baldridge, K. K.; Kast, P.; Hilvert, D., Experimental and Computational Investigation of the Uncatalyzed Rearrangement and Elimination Reactions of Isochorismate. *J. Am. Chem. Soc.* **2006**, 128, (6), 2043-2051.
- 347. Jeselnik, M.; Leban, I.; Polanc, S.; Kocevar, M., *D*-Gulonolactone as a Synthon for *L*-Noviose: First Preparation of 4-*O*-Demethyl-*L*-noviofuranose and Related Derivatives. *Org. Lett.* **2003**, *5*, (15), 2651-2653.
- 348. Tadano, K.; Kanazawa, S.; Ogawa, S., Synthesis of (3S,4S)-dihydro-4-ethyl-3hydroxy-4-methyl-2(3H)-furanone, a pantolactone homolog isolated from Marshallia tenuifolia. J. Org. Chem. **1988**, 53, (16), 3868-3870.
- 349. De Luca, L.; Giacomelli, G.; Porcheddu, A., A Very Mild and Chemoselective Oxidation of Alcohols to Carbonyl Compounds. *Org. Lett.* **2001**, *3*, (19), 3041-3043.
- 350. Ahn, H. C.; Choi, K., N-(2-Nitrophenyl)proline: An Intramolecular Hydrogen Bond Forming Reagent for the Determination of the Absolute Configuration of Primary Amines. *Org. Lett.* **2007**, 9, (19), 3853-3855.
- Baldwin, J. E.; Killin, S. J.; Adlington, R. M.; Spiegel, U., Synthesis of nbenzyloxycarbonyl-l-[alpha]-aminoadipic acid, [alpha]-benzyl ester. *Tetrahedron* 1988, 44, (9), 2633-2636.
- 352. Inanaga, J.; Hirata, K.; Saeki, H.; Katsuki, T.; Yamaguchi, M., A Rapid Esterification by Means of Mixed Anhydride and Its Application to Large-ring Lactonization. *Bull. Chem. Soc. Jpn.* **1979**, 52, (7), 1989-1993.
- 353. Hassner, A.; Alexanian, V., Direct room temperature esterification of carboxylic acids. *Tetrahedron Lett.* **1978**, 19, (46), 4475-4478.

- 354. Neises, B.; Steglich, W., Simple Method for the Esterification of Carboxylic Acids. *Angew. Chem. Int. Ed. Engl.* **1978**, 17, (7), 522-524.
- 355. Höfle, G.; Steglich, W.; Vorbrüggen, H., 4-Dialkylaminopyridines as Highly Active Acylation Catalysts. [New synthetic method (25)]. *Angew. Chem. Int. Ed. Engl.* **1978**, 17, (8), 569-583.
- 356. Sakaitani, M.; Ohfune, Y., Syntheses and reactions of silyl carbamates. 1. Chemoselective transformation of amino protecting groups via tertbutyldimethylsilyl carbamates. *J. Org. Chem.* **1990**, 55, (3), 870-876.
- 357. Ashton, P. R.; Philip, D.; Spencer, N.; Stoddart, J. F., A new design strategy for the self-assembly of molecular shuttles. *J. Chem. Soc., Chem. Commun.* **1992**, 1124-1128.
- 358. Armspach, D.; Matt, D., The tris(4-tert-butylphenyl)methyl group: a bulky substituant for effective regioselective difunctionalisation of cyclomaltohexaose. *Carbohydr. Res.* **1998**, 310, (1-2), 129-133.
- 359. Zhu, K.; Simpson, J. H.; Delaney, E. J.; Nugent, W. A., Synthesis of Z-5-Carboxymethylene-1,3-dioxolan-4-ones: A Better Way. J. Org. Chem. 2007, 72, (10), 3949-3951.
- 360. Winneroski, L. L.; Xu, Y., Two Complementary Approaches toward 2-Alkoxy Carboxylic Acid Synthesis from 1,3-Dioxolan-4-ones. J. Org. Chem. 2004, 69, (15), 4948-4953.
- Moorlag, H.; Kellogg, R. M.; Kloosterman, M.; Kaptein, B.; Kamphuis, J.; Schoemaker, H. E., Pig-liver-esterase-catalyzed hydrolyses of racemic.alpha.substituted.alpha.-hydroxy esters. J. Org. Chem. 1990, 55, (23), 5878-5881.
- Collins, L.; Franzblau, S. G., Microplate alamar blue assay versus BACTEC 460 system for high- throughput screening of compounds against Mycobacterium tuberculosis and Mycobacterium avium. *Antimicrob. Agents Chemother.* 1997, 41, (5), 1004-1009.
- 363. Falzari, K.; Zhu, Z.; Pan, D.; Liu, H.; Hongmanee, P.; Franzblau, S. G., In Vitro and In Vivo Activities of Macrolide Derivatives against Mycobacterium tuberculosis. *Antimicrob. Agents Chemother.* **2005**, 49, (4), 1447-1454.
- Cho, S. H.; Warit, S.; Wan, B.; Hwang, C. H.; Pauli, G. F.; Franzblau, S. G., Low-oxygen-recovery assay for high-throughput screening of compounds against nonreplicating Mycobacterium tuberculosis. *Antimicrob. Agents Chemother.* 2007, 51, (4), 1380-1385.
- 365. Roy, B. G.; Roy, A.; Achari, B.; Mandal, S. B., A simple one-pot entry to cyclic ethers of varied ring sizes from diols via phosphonium ion induced iodination and

base catalyzed Williamson etherification. *Tetrahedron Lett.* **2006**, 47, (44), 7783-7787.

- 366. Lang, S.; Murphy, J. A., Azide rearrangements in electron-deficient systems. *Chem. Soc. Rev.* **2006**, 35, (2), 146-156.
- 367. Braese, S.; Gil, C.; Knepper, K.; Zimmermann, V., Organic azides. An exploding diversity of a unique class of compounds. *Angew. Chem., Int. Ed.* **2005,** 44, (33), 5188-5240.
- 368. Scriven, E. F. V.; Turnbull, K., Azides: their preparation and synthetic uses. *Chem. Rev.* **1988**, 88, (2), 297-368.
- 369. Scriven, E. F. V., *Azides and Nitrenes: Reactivity and Utility*. Academic Press, Inc.: Orlando, Florida, 1984; p 542.
- 370. Dori, Z.; Ziolo, R. F., Chemistry of coordinated azides. *Chem. Rev.* **1973**, 73, (3), 247-54.
- L'Abbe, G., Decomposition and addition reactions of organic azides. *Chem. Rev.* 1969, 69, (3), 345-63.
- 372. Lwowski, W., Nitrenes and the decomposition of carbonyl azides. *Angew. Chem., Int. Ed. Engl.* **1967,** 6, (11), 897-906.
- 373. Lwowski, W.; Mattingly, T. W., Jr., The decomposition of ethyl azidoformate in cyclohexene and in cyclohexane. *J. Am. Chem. Soc.* **1965**, 87, (9), 1947-1958.
- 374. Lwowski, W., Nitrenes. John Wiley & Sons, Inc.: New York, 1970; p 457.
- 375. Abramovitch, R. A., *Reactive Intermediates*. Plenum Press: New York, 1982; Vol. 2, p 599.
- 376. Kwart, H.; Kahn, A. A., Copper-Catalyzed Decomposition of Benzenesulfonyl Azide in Hydroxylic Media. J. Am. Chem. Soc. **1967**, 89, (8), 1950-1951.
- 377. Kwart, H.; Khan, A. A., Copper-Catalyzed Decomposition of Benzenesulfonyl Azide in Cyclohexene Solution. *J. Am. Chem. Soc.* **1967**, 89, (8), 1951-1953.
- 378. Yamada, Y.; Yamamoto, T.; Okawara, M., Synthesis and Reaction of new Type I–N Ylide, N-Tosyliminoiodinane. *Chem. Lett.* **1975**, 4, (4), 361-362.
- 379. Hillhouse, G. L.; Haymore, B. L., Interaction of aryl azides with tungsten complexes. Three new types of reactions yielding coordinated RN3PR3, RN3H, and RN. J. Organomet. Chem. **1978**, 162, (2), C23-C26.
- 380. Paul Mueller, C. B. I. N., Rhodium(II)-catalyzed nitrene transfer with phenyliodonium ylides. J. Phys. Org. Chem. **1998**, 11, (8-9), 597-601.

- Brandt, P.; Sodergren, M. J.; Andersson, P. G.; Norrby, P. O., Mechanistic Studies of Copper-Catalyzed Alkene Aziridination. J. Am. Chem. Soc. 2000, 122, (33), 8013-8020.
- 382. Levites-Agababa, E.; Menhaji, E.; Perlson, L. N.; Rojas, C. M., Amidoglycosylation via Metal-Catalyzed Internal Nitrogen Atom Delivery. *Org. Lett.* **2002**, 4, (5), 863-865.
- 383. Mueller, P.; Fruit, C., Enantioselective Catalytic Aziridinations and Asymmetric Nitrene Insertions into CH Bonds. *Chem. Rev.* **2003**, 103, (8), 2905-2919.
- 384. Liang, J.-L.; Yuan, S.-X.; Chan, P. W. H.; Che, C.-M., Chiral rhodium(II,II) dimers catalyzed enantioselective intramolecular aziridination of sulfonamides and carbamates. *Tetrahedron Lett.* **2003**, 44, (31), 5917-5920.
- 385. Wehn, P. M.; Lee, J.; Du Bois, J., Stereochemical Models for Rh-Catalyzed Amination Reactions of Chiral Sulfamates. *Org. Lett.* **2003**, **5**, (25), 4823-4826.
- 386. Halfen, J. A., Recent Advances in Metal-Mediated Carbon-Nitrogen Bond Formation Reactions: Aziridination and Amidation. *Curr. Org. Chem.* **2005**, 9, (7), 657.
- 387. Kawabata, H.; Omura, K.; Katsuki, T., Asymmetric aziridination: a new entry to optically active non-N-protected aziridines. *Tetrahedron Lett.* **2006**, 47, (10), 1571-1574.
- 388. Ruppel, J. V.; Kamble, R. M.; Zhang, X. P., Cobalt-Catalyzed Intramolecular C-H Amination with Arylsulfonyl Azides. *Org. Lett.* **2007**.
- 389. Rolf, H., 1,3-Dipolar Cycloadditions. Past and Future. Angew. Chem. Int. Ed. Engl. 1963, 2, (10), 565-598.
- 390. Rostovtsev, V. V.; Green, L. G.; Fokin, V. V.; Sharpless, K. B., A stepwise Huisgen cycloaddition process: copper(I)-catalyzed regioselective "ligation" of azides and terminal alkynes. *Angew. Chem., Int. Ed.* **2002**, 41, (14), 2596-2599.
- 391. Himo, F.; Lovell, T.; Hilgraf, R.; Rostovtsev, V. V.; Noodleman, L.; Sharpless, K. B.; Fokin, V. V., Copper(I)-Catalyzed Synthesis of Azoles. DFT Study Predicts Unprecedented Reactivity and Intermediates. J. Am. Chem. Soc. 2005, 127, (1), 210-216.
- 392. Bock, V. D.; Hiemstra, H.; van Maarseveen, J. H., Cu(I)-catalyzed alkyne-azide click cycloadditions from a mechanistic and synthetic perspective. *Eur. J. Org. Chem.* **2005**, (1), 51-68.
- 393. Zhang, L.; Chen, X.; Xue, P.; Sun, H. H. Y.; Williams, I. D.; Sharpless, K. B.; Fokin, V. V.; Jia, G., Ruthenium-Catalyzed Cycloaddition of Alkynes and Organic Azides. J. Am. Chem. Soc. 2005, 127, (46), 15998-15999.

- 394. Kolb, H. C.; Finn, M. G.; Sharpless, K. B., Click chemistry: diverse chemical function from a few good reactions. *Angew. Chem., Int. Ed.* **2001,** 40, (11), 2004-2021.
- 395. Kolb, H. C.; Sharpless, K. B., The growing impact of click chemistry on drug discovery. *Drug Discovery Today* **2003**, **8**, (24), 1128-1137.
- 396. Lewis, W. G.; Green, L. G.; Grynszpan, F.; Radic, Z.; Carlier, P. R.; Taylor, P.; Finn, M. G.; Sharpless, K. B., Click chemistry in situ: Acetylcholinesterase as a reaction vessel for the selective assembly of a femtomolar inhibitor from an array of building blocks. *Angew. Chem., Int. Ed.* **2002**, 41, (6), 1053-1057.
- 397. Tornoe, C. W.; Christensen, C.; Meldal, M., Peptidotriazoles on Solid Phase: [1,2,3]-Triazoles by Regiospecific Copper(I)-Catalyzed 1,3-Dipolar Cycloadditions of Terminal Alkynes to Azides. *J. Org. Chem.* **2002**, 67, (9), 3057-3064.
- 398. Seo, T. S.; Li, Z.; Ruparel, H.; Ju, J., Click Chemistry to Construct Fluorescent Oligonucleotides for DNA Sequencing. *J. Org. Chem.* **2003**, 68, (2), 609-612.
- 399. Wang, Q.; Chan, T. R.; Hilgraf, R.; Fokin, V. V.; Sharpless, K. B.; Finn, M. G., Bioconjugation by copper(I)-catalyzed azide-alkyne [3 + 2] cycloaddition. *J. Am. Chem. Soc.* **2003**, 125, (11), 3192-3193.
- 400. Speers, A. E.; Adam, G. C.; Cravatt, B. F., Activity-based protein profiling in vivo using a copper(I)-catalyzed azide-alkyne [3 + 2] cycloaddition. *J. Am. Chem. Soc.* **2003**, 125, (16), 4686-4687.
- Lee, L. V.; Mitchell, M. L.; Huang, S.-J.; Fokin, V. V.; Sharpless, K. B.; Wong, C.-H., A Potent and Highly Selective Inhibitor of Human a-1,3-Fucosyltransferase via Click Chemistry. J. Am. Chem. Soc. 2003, 125, (32), 9588-9589.
- 402. Lummerstorfer, T.; Hoffmann, H., Click Chemistry on Surfaces: 1,3-Dipolar Cycloaddition Reactions of Azide-Terminated Monolayers on Silica. *J. Phys. Chem. B* **2004**, 108, (13), 3963-3966.
- 403. Collman, J. P.; Devaraj, N. K.; Chidsey, C. E. D., "Clicking" Functionality onto Electrode Surfaces. *Langmuir* **2004**, 20, (4), 1051-1053.
- 404. Bourne, Y.; Kolb, H. C.; Radic, Z.; Sharpless, K. B.; Taylor, P.; Marchot, P., Freeze-frame inhibitor captures acetylcholinesterase in a unique conformation. *Proc. Natl. Acad. Sci.* **2004**, 101, (6), 1449-1454.
- 405. Bodine, K. D.; Gin, D. Y.; Gin, M. S., Synthesis of Readily Modifiable Cyclodextrin Analogues via Cyclodimerization of an Alkynyl-Azido Trisaccharide. J. Am. Chem. Soc. **2004**, 126, (6), 1638-1639.

- 406. Suarez, P. L.; Gandara, Z.; Gomez, G.; Fall, Y., Vitamin D and click chemistry. Part 1: a stereoselective route to vitamin D analogues with triazole rings in their side chains. *Tetrahedron Lett.* **2004**, 45, (24), 4619-4621.
- 407. Suh, B.-C.; Jeon, H.; Posner, G. H.; Silverman, S. M., Vitamin D side chain triazole analogs via cycloaddition 'click' chemistry. *Tetrahedron Lett.* **2004**, 45, (24), 4623-4625.
- 408. Manetsch, R.; Krasinski, A.; Radic, Z.; Raushel, J.; Taylor, P.; Sharpless, K. B.; Kolb, H. C., In Situ Click Chemistry: Enzyme Inhibitors Made to Their Own Specifications. J. Am. Chem. Soc. **2004**, 126, (40), 12809-12818.
- 409. Ramachary, D. B.; Barbas, C. F., III, Towards organo-click chemistry: Development of organocatalytic multicomponent reactions through combinations of Aldol, Wittig, Knoevenagel, Michael, Diels-Alder and Huisgen cycloaddition reactions. *Chem.--Eur. J.* **2004**, 10, (21), 5323-5331.
- 410. Punna, S.; Kuzelka, J.; Wang, Q.; Finn, M. G., Head-to-tail peptide cyclodimerization by copper-catalyzed azide-alkyne cycloaddition. *Angew. Chem., Int. Ed.* **2005,** 44, (15), 2215-2220.
- 411. Parent, M.; Mongin, O.; Kamada, K.; Katan, C.; Blanchard-Desce, M., New chromophores from click chemistry for two-photon absorption and tuneable photoluminescence. *Chem. Commun.* **2005**, (15), 2029-2031.
- 412. Krasinski, A.; Radic, Z.; Manetsch, R.; Raushel, J.; Taylor, P.; Sharpless, K. B.; Kolb, H. C., In Situ Selection of Lead Compounds by Click Chemistry: Target-Guided Optimization of Acetylcholinesterase Inhibitors. *J. Am. Chem. Soc.* 2005, 127, (18), 6686-6692.
- 413. Ryu, E.-H.; Zhao, Y., Efficient Synthesis of Water-Soluble Calixarenes Using Click Chemistry. *Org. Lett.* **2005**, 7, (6), 1035-1037.
- 414. Pagliai, F.; Pirali, T.; Del Grosso, E.; Di Brisco, R.; Tron Gian, C.; Sorba, G.; Genazzani Armando, A., Rapid Synthesis of Triazole-Modified Resveratrol Analogues via Click Chemistry. J. Med. Chem. 2006, 49, (2), 467-70.
- 415. Luo, S.; Xu, H.; Mi, X.; Li, J.; Zheng, X.; Cheng, J. P., Evolution of Pyrrolidine-Type Asymmetric Organocatalysts by "Click" Chemistry. *J. Org. Chem.* **2006**, 71, (24), 9244-9247.
- 416. Lipshutz, B. H.; Taft, B. R., Heterogeneous Copper-in-Charcoal-Catalyzed Click Chemistry. *Angew. Chem. Int. Ed.* **2006**, 45, (48), 8235-8238.
- 417. Aucagne, V.; Berna, J.; Crowley, J. D.; Goldup, S. M.; Hanni, K. D.; Leigh, D. A.; Lusby, P. J.; Ronaldson, V. E.; Slawin, A. M. Z.; Viterisi, A.; Walker, D. B., Catalytic "Active-Metal" Template Synthesis of [2]Rotaxanes, [3]Rotaxanes, and

Molecular Shuttles, and Some Observations on the Mechanism of the Cu(I)-Catalyzed Azide-Alkyne 1,3-Cycloaddition. J. Am. Chem. Soc. 2007.

- 418. Salisbury, C. M.; Cravatt, B. F., Optimization of Activity-Based Probes for Proteomic Profiling of Histone Deacetylase Complexes. *J. Am. Chem. Soc.* **2008**.
- 419. Bodnar, B. S.; Miller, M. J., Reactions of Nitroso Hetero-Diels-Alder Cycloadducts with Azides: Stereoselective Formation of Triazolines and Aziridines. J. Org. Chem. 2007, 72, (10), 3929-3932.
- 420. Miller, M. J., Isourea-mediated preparation of dehydro amino acids. J. Org. Chem. 1980, 45, (15), 3131-3132.
- 421. Srinivasan, A.; Richards, K. D.; Olsen, R. K., Comments on assignment of stereochemistry to 2-acylaminocrotonates. *Tetrahedron Lett.* **1976**, (12), 891-894.
- 422. Panella, L.; Aleixandre, A. M.; Kruidhof, G. J.; Robertus, J.; Feringa, B. L.; deVries.J.G; Minnaard, A. J., Enantioselective Rh-Catalyzed Hydrogenation of N-Formyl Dehydroamino Esters with Monodentate Phosphoramidite Ligands. *J. Org. Chem.* **2006**, 71, (5), 2026-2036.
- 423. Gangadasu, B.; Narender, P.; Bharath Kumar, S.; Ravinder, M.; Ananda Rao, B.; Ramesh, C.; China Raju, B.; Jayathirtha Rao, V., Facile and selective synthesis of chloronicotinaldehydes by the Vilsmeier reaction. *Tetrahedron* **2006**, 62, (35), 8398-8403.
- 424. Alvarez, S. G.; Alvarez, M. T., A practical procedure for the synthesis of alkyl azides at ambient temperature in dimethyl sulfoxide in high purity and yield. *Synthesis* **1997**, (4), 413-414.
- 425. Pollex, A.; Hiersemann, M., Catalytic Asymmetric Claisen Rearrangement in Natural Product Synthesis: Synthetic Studies toward (-)-Xeniolide F. *Org. Lett.* **2005**, 7, (25), 5705-5708.
- 426. Lindsay, R. O.; Allen, C. F. H., Phenyl azide. Org. Synth. 1955, 3, 710.
- 427. Shaw, E., Observations on the cyclization of a substituted a-formamidoamidine to aminoimidazolecarboxamide derivatives. *J. Org. Chem.* **1965**, 30, (10), 3371-3373.
- 428. Shaw, E., 5-Amino-4-imidazolecarboxamide riboside from inosine. Ring-opening reactions of purine nucleosides. *J. Am. Chem. Soc.* **1958**, 80, 3899-902.
- 429. Shaw, E.; Woolley, D. W., A new and convenient synthesis of 4-amino-5imidazolecarboxamide. J. Biol. Chem. **1949**, 181, 89-93.
- 430. Spaltenstein, A.; Holler, T. P.; Hopkins, P. B., New synthesis of 4- and 5imidazolethiols. J. Org. Chem. 1987, 52, (14), 2977-2979.

- 431. Reiter, L. A., Synthesis of 4(5)-acyl-, 1-substituted 5-acyl- and 1-substituted 4-acyl-1H-imidazoles from 4-aminoisoxazoles. *J. Org. Chem.* **1987**, 52, (13), 2714-26.
- 432. Reiter, L. A., A general synthesis of 4(5)-acylimidazoles from 4-acylaminoisoxazoles. *Tetrahedron Lett.* **1985**, 26, (29), 3423-6.
- 433. Toplak, R.; Svete, J.; Grdadolnik, S. G.; Stanovnik, B., Methyl (Z)-2-[(benzyloxycarbonyl)amino]-3-dimethyl-aminopropenoate in the synthesis of heterocyclic systems. Synthesis of (benzyloxycarbonyl)amino substituted fused pyrimidinones. *Collect. Czech. Chem. Commun.* **1999**, 64, (2), 177-189.
- 434. Celestina, T.; Golobic, A.; Svete, J.; Stanovnik, B., Intramolecular thermal cyclizations of methyl (E)-3-arylamino-2-benzoylaminobut-2-enoates. *ARKIVOC* **2004**, (7), 169-176.
- 435. Stanovnik, B.; Svete, J., Synthesis of Heterocycles from Alkyl 3-(Dimethylamino)propenoates and Related Enaminones. *Chem. Rev.* 2004, 104, (5), 2433-2480.
- 436. Bratusek, U.; Recnik, S.; Svete, J.; Meden, A.; Stanovnik, B., Transformations of methyl 2-benzoylamino-2-oxobutanoate. The synthesis of oxazolo[4,5-c]quinoline and 1-substituted 1H-imidazole-4-carboxylate derivatives. *Heterocycles* **2003**, 60, (5), 1161-1172.
- 437. Stanovnik, B.; Svete, J., Alkyl 2-substituted 3-(dimethylamino)propenoates and related compounds versatile reagents in heterocyclic chemistry. *Synlett* **2000**, (8), 1077-1091.
- 438. Gurjar, M. K.; Karmakar, S.; Mohapatra, D. K.; Phalgune, U. D., A novel application of a [3+2] cycloaddition reaction for the synthesis of the piperazinone rings of pseudotheonamides A1 and A2. *Tetrahedron Lett.* **2002**, 43, (10), 1897-1900.
- 439. Konda, Y.; Sato, T.; Tsushima, K.; Dodo, M.; Kusunoki, A.; Sakayanagi, M.; Sato, N.; Takeda, K.; Harigaya, Y., An efficient synthesis of (pyrrolidin-2-ylidene)glycinate using intramolecular 1, 3-dipolar cycloaddition of azide and olefin. *Tetrahedron* **1999**, 55, (44), 12723-12740.
- 440. Prager, R. H.; Razzino, P., Heterocyclic Synthesis With Azides. III. Reactions of Triazolines Made From Arylmethylidenemalonates. *Aust. J. Chem.* **1994**, 47, (7), 1375-1385.
- 441. Sha, C. K.; Ouyang, S. L.; Hsieh, D. Y.; Chang, R. C.; Chang, S. C., Intramolecular 1,3-dipolar cycloaddition of alkyl azide enones and rearrangements of the triazoline intermediates. Formal total synthesis of (+-.)desamylperhydrohistrionicotoxin. J. Org. Chem. **1986**, 51, (9), 1490-1494.

- 442. Benati, L.; Montevecchi, P. C.; Spagnolo, P., Thermal reactions of aryl azides with trans-1,2-dibenzoyl- and trans-1,2-diacetylethylene. Reactivity of 4,5-dibenzoyl- and 4,5-diacetyl-1-aryltriazolines. *J. Chem. Soc., Perkin Trans. 1* **1989**, (12), 2235-2243.
- 443. Reddy, D. S.; Judd, W. R.; Aube, J., Lewis Acid-Mediated Reactions of Alkyl Azides with a,b-Unsaturated Ketones. *Org. Lett.* **2003**, *5*, (21), 3899-3902.
- 444. Dubreuil, J. F.; Bazureau, J. P., Rate accelerations of 1,3-dipolar cycloaddition reactions in ionic liquids. *Tetrahedron Lett.* **2000**, 41, (38), 7351-7355.
- 445. Creary, X.; Willis, E. D.; Moura-Letts, G.; Curran, D. P., Preparation of 1-butyl-3-methylimidazolium tetrafluoroborate. *Org. Synth.* **2005**, 82, 166-169.
- 446. Shireman, B. T.; Miller, M. J., Synthesis of Enantiomerically and Diastereomerically Pure 2(S)-Amino-6(R)-hydroxy-1,7-heptanedioic Acid Dimethyl Ester Hydrochloride from Cycloheptadiene. *J. Org. Chem.* 2001, 66, 4809-4813.
- 447. Shea, K. J.; Kim, J. S., Influence of strain on chemical reactivity. Relative reactivity of torsionally strained double bonds in 1,3-dipolar cycloadditions. *J. Am. Chem. Soc.* **1992**, 114, (12), 4846-4855.
- 448. Becker, K. B.; Hohermuth, M. K., 1,3-Dipolar cycloadditions to strained olefins. *Helv. Chim. Acta* **1979**, 62, (6), 2025-36.
- 449. Taniguchi, H.; Ikeda, T.; Imoto, E., 1,3-Dipolar cycloadditions to bicyclic olefins.
  V. The stereo- and regiochemistry of 1,3-dipolar cycloadditions to several unsymmetrical bicyclic olefins. *Bull. Chem. Soc. Jpn.* 1978, 51, (6), 1859-65.
- 450. Scheiner, P., Triazoline photodecomposition. Preparation of aziridines. *Tetrahedron* **1968**, 24, (6), 2757-66.
- 451. Scheiner, P., Triazoline photodecomposition. Stereochemistry and mechanism. J. *Am. Chem. Soc.* **1968**, 90, (4), 988-92.
- 452. Rolf, H., 1,3-Dipolar Cycloadditions. Past and Future. *Angew. Chem., Int. Ed. Engl.* **1963**, 2, (10), 565-598.
- 453. Tarabara, I. N.; Kas'yan, A. O.; Yarovoi, M. Y.; Shishkina, S. V.; Shishkin, O. V.; Kas'yan, L. I., Reactions of bicyclo[2.2.1]hept-5-ene-2,3-dicarboximides with aromatic azides. *Russ. J. Org. Chem.* **2004**, 40, (7), 992-998.
- 454. Ishikura, M.; Katagiri, N., Synthetic application of 2-azabicyclo[2.2.1]hept-5-en-3-one (ABH) as a versatile synthon. *Recent Res. Devel. Organic Chem.* **2004**, 8, (Pt. 1), 197-205.

- 455. Ishikura, M.; Murakami, A.; Katagiri, N., First synthesis of 2',3'-epiminocarbocyclic nucleosides. *Org. Biomol. Chem.* **2003**, 1, (3), 452-453.
- 456. Ishikura, M.; Kudo, S.; Hino, A.; Ohnuki, N.; Katagiri, N., A concise access to 6azabicyclo[3.1.0]hexanes via high-pressure promoted cycloaddition reaction of azides to ABH. *Heterocycles* **2000**, 53, (7), 1499-1504.
- 457. Stout, D. M.; Takaya, T.; Meyers, A. I., Unequivocal synthesis of N-substituted 1,4-dihydropyridines. J. Org. Chem. 1975, 40, (5), 563-9.
- 458. Huenig, S.; Kraft, P., Azobridges from azines. XIII. Azobridged polycycles: skeletal rearrangements by denitrogenation of triazolines and solvolyses of aziridines. *Heterocycles* **1995**, 40, (2), 639-52.
- 459. Tarabara, I. N.; Kas'yan, A. O.; Yarovoi, M. Y.; Shishkina, S. V.; Shishkin, O. V.; Kas'yan, L. I., Reactions of Bicyclo[2.2.1]hept-5-ene-2,3-dicarboximides with Aromatic Azides. *Russ. J. Org. Chem.* **2004**, 40, (7), 992-998.
- 460. Malpass, J. R.; Belkacemi, D.; Griffith, G. A.; Robertson, M. D., Cycloaddition of phenyl azide to unsymmetrical azaicyclic alkenes. *ARKIVOC* **2002**, (6), 164-174.
- 461. Reymond, J.-L.; Vogel, P., Stereoselective amino-hydroxylation of the double bond in 7-oxabicyclo[2.2.1]hept-5-en-2-yl derivatives. Remote substituent participation in acid-catalyzed decompositions of aziridines and triazolines. *Tetrahedron Lett.* **1988**, 29, (30), 3695-3698.
- 462. Stout, D. M.; Takaya, T.; Meyers, A. I., Unequivocal synthesis of N-substituted 1,4-dihydropyridines. J. Org. Chem. 1975, 40, (5), 563-569.
- 463. Peterson, W. R., Jr.; Arkles, B.; Washburne, S. S., Reaction of trimethylsilyl azide with bridged bicyclic olefins. *J. Organomet. Chem.* **1976**, 121, (3), 285-91.
- 464. Hong, K. B.; Donahue, M. G.; Johnston, J. N., On the Nature of Rate Acceleration in the Synthesis and Fragmentation of Triazolines by Bronsted Acid: Secondary Catalysis by Water (Hydronium Triflate). *J. Am. Chem. Soc.* **2008**, 130, (7), 2323-2328.
- Krasinski, A.; Fokin, V. V.; Sharpless, K. B., Direct Synthesis of 1,5-Disubstituted-4-magnesio-1,2,3-triazoles, Revisited. Org. Lett. 2004, 6, (8), 1237-1240.
- 466. Rodionov, V. O.; Fokin, V. V.; Finn, M. G., Mechanism of the ligand-free Culcatalyzed azide-alkyne cycloaddition reaction. *Angew. Chem., Int. Ed.* **2005,** 44, (15), 2210-2215.
- 467. Jones, G. O.; Houk, K. N., Predictions of Substituent Effects in Thermal Azide 1,3-Dipolar Cycloadditions: Implications for Dynamic Combinatorial

(Reversible) and Click (Irreversible) Chemistry. J. Org. Chem. 2008, 73, (4), 1333-1342.

- 468. Baskin, J. M.; Prescher, J. A.; Laughlin, S. T.; Agard, N. J.; Chang, P. V.; Miller, I. A.; Lo, A.; Codelli, J. A.; Bertozzi, C. R., From the Cover: Copper-free click chemistry for dynamic in vivo imaging. *Proc. Natl. Acad. Sci.* 2007, 104, (43), 16793-16797.
- 469. Agard, N. J.; Prescher, J. A.; Bertozzi, C. R., A Strain-Promoted [3 + 2] Azide-Alkyne Cycloaddition for Covalent Modification of Biomolecules in Living Systems. J. Am. Chem. Soc. 2004, 126, (46), 15046-15047.
- 470. Gorpinchenko, V.; Yatsynich, E.; Petrov, D.; Karachurina, L.; Hisamutdinova, R.; Baschenko, N.; Dokichev, V.; Tomilov, Y.; Yunusov, M.; Nefedov, O., Synthesis and Antiarrhythmic Activity of 5-Amino-exo-3-azatricyclo[5.2.1.02,6]decan-4one. *Pharm. Chem. J.* 2005, 39, (6), 289-291.
- 471. Yatsynich, E. A.; Petrov, D. V.; Dokichev, V. A.; Tomilov, Y. V., Synthesis of Amines from Norbornane Series. *Russ. J. Org. Chem.* **2005**, 41, (8), 1187-1191.
- 472. Christl, M.; Brunn, E.; Roth, W. R.; Lennartz, H. W., 7-Methyl- and 7phenylcyclohepta-1,3,5-trienes from benzvalene via 3,3a,4,5,6,6a-hexahydro-4,5,6-methenocyclopentapyrazoles and tetracyclo[4.1.0.02,4,03,5]heptanes. *Tetrahedron* **1989**, 45, (10), 2905-2915.
- 473. Lal, K.; Salomon, R. G., 6-Substituted bicyclo[2.2.1]hept-5-en-2-one ketals. J. Org. Chem. **1989**, 54, (11), 2628-2632.
- 474. Adam, W.; Carballeira, N.; Craemer, E.; Lucchini, V.; Peters, E. M.; Peters, K.; Von Schnering, H. G., 1,3-Dipolar cycloadditions with the keto- and spirocyclopropane-substituted norbornenes bicyclo[2.2.1]hept-5-ene-2-one and spiro[bicyclo[2.2.1]hept-5-ene-2,1'-cyclopropane]. *Chem. Ber.* **1987**, 120, (5), 695-703.
- 475. Wilt, J. W.; Tufano, M. D., 8-Methylene-exo-3,3diphenyltricyclo[3.2.1.02,4]octane, a probe for addition reaction mechanism. J. Org. Chem. **1985**, 50, (14), 2600-2601.
- 476. Rieber, N.; Boehm, H., 1,3-Dipolar additions to 3,4diazatricyclo[4.2.1.02.5]nona-3,7-diene and 3,4-diazatricyclo[4.2.1.02.5]non-7ene derivatives. *J. Heterocycl. Chem.* **1981**, 18, 1-7.
- 477. Ishikura, M.; Murakami, A.; Katagiri, N., Preparation of 2',3'-methanocarbocyclic nucleosides through the addition of diazo-methane to 2azabicyclo[2.2.1]hept-5-en-3-one. *Heterocycles* **2002**, 58, 317-324.

- 478. Katagiri, N.; Yamatoya, Y.; Ishikura, M., Synthesis of nucleosides and their related compounds. ILV. The first synthesis of a 2',3'-methano carbocyclic nucleoside. *Tetrahedron Lett.* **1999**, 40, (51), 9069-9072.
- 479. Allred, E. L.; Hinshaw, J. C., Tricyclo[3.1.0.02,4]hexane. J. Am. Chem. Soc. **1968**, 90, (24), 6885-7.
- 480. Gorpinchenko, V. A.; Petrov, D. V.; Spirikhin, L. V.; Dokichev, V. A.; Tomilov, Y. V., Synthesis of N3-substituted methyl exo-3,4-diazatricyclo[5.2.1.02,6]dec-4-ene-5-carboxylates. *Russ. J. Org. Chem.* **2006**, 42, (11), 1706-1710.
- 481. Wulfman, D. S.; McDaniel, R. S., Jr., Decomposition of a 1-pyrazoline by cupric tetrafluoroborate. *Tetrahedron Lett.* **1975**, 16, (50), 4523-4524.
- 482. Kumar, S.; Murray, R. W., Carbonyl oxide chemistry. The NIH shift. J. Am. Chem. Soc. 1984, 106, (4), 1040-1045.
- 483. Fischer, W.; Anselme, J. P., Diazoalkanes from hydrazone anions. Novel oxidation reaction. J. Am. Chem. Soc. **1967**, 89, (20), 5312-13.
- 484. Smith, L. I.; Howard, K. L., Diphenyldiazomethane. Org. Synth. 1955, 3, 351.
- 485. Miller, J. B., Preparation of crystalline diphenyldiazomethane. J. Org. Chem. 1959, 24, 560-1.
- Blomquist, A. T.; Meinwald, Y. C., Synthesis of Some Conjugated Cyclobutane Polyolefins and their 1,2-Cycloaddition to Tetracyanoethylene. *J. Am. Chem. Soc.* 1959, 81, (3), 667-672.
- 487. Hall, H. K., Synthesis of Two Atom-Bridged Tetracyclic Ketones. J. Org. Chem. **1960**, 25, (1), 42-44.
- 488. Sasaki, T.; Eguchi, S.; Sugimoto, M.; Hibi, F., Reactions of isoprenoids. XVII. Cycloaddition reactions of norbornadiene with some unsymmetrically substituted dienophiles. Competitive ionic additions with homo-Diels-Alder reactions. *J. Org. Chem.* **1972**, 37, (14), 2317-20.
- 489. Lautens, M.; Edwards, L. G., Stereoselectivity in the homo Diels-Alder reaction. *Tetrahedron Lett.* **1989**, 30, (49), 6813-6816.
- 490. Lautens, M.; Blackwell, J., The utility of the homo Diels-Alder reaction in the synthesis of natural product skeletons. A cycloaddition-fragmentation-cyclization sequence to diquinanes and triquinanes. *Synthesis* **1998**, (Spec.), 537-546.
- 491. Lautens, M.; Edwards, L. G., Regiochemical control in the homo-Diels-Alder reaction: substituent effects. J. Org. Chem. 1991, 56, (12), 3761-3763.

- 492. Lautens, M.; Edwards, L. G.; Tam, W.; Lough, A. J., Nickel-catalyzed  $[2\pi + 2\pi + 2\pi]$  (homo-Diels-Alder) and  $[2\pi + 2\pi]$  cycloadditions of bicyclo[2.2.1]hepta-2,5-dienes. *J. Am. Chem. Soc.* **1995**, 117, (41), 10276-10291.
- 493. Lautens, M.; Tam, W.; Blackwell, J., Synthesis of Highly Functionalized Diquinanes by the Regio- and Stereoselective Cleavage of Homo-Diels-Alder Cycloadducts. J. Am. Chem. Soc. **1997**, 119, (3), 623-624.
- 494. Lautens, M.; Tam, W.; Edwards, L. G., Cobalt-catalyzed intramolecular homo Diels-Alder reactions. J. Org. Chem. **1992**, 57, (1), 8-9.
- 495. Lautens, M.; Tam, W.; Lautens, J. C.; Edwards, L. G.; Crudden, C. M.; Smith, A. C., Cobalt-Catalyzed  $[2\pi + 2\pi + 2\pi]$  (Homo-Diels-Alder) and  $[2\pi + 2\pi + 4\pi]$  Cycloadditions of Bicyclo[2.2.1]hepta-2,5-dienes. *J. Am. Chem. Soc.* **1995**, 117, (26), 6863-6879.
- 496. Gugelchuk, M. M.; Doherty-Kirby, A. L., Kinetic and Modeling Studies on the Nickel-Catalyzed Homo-Diels-Alder Addition of 7-Substituted Norbornadienes. *J. Org. Chem.* **1996**, 61, (10), 3490-3495.
- 497. Gugelchuk, M. M.; Doherty-Kirby, A. L., Kinetic and Modeling Studies on the Nickel-Catalyzed Homo-Diels-Alder Addition of 7-Substituted Norbornadienes. *J. Org. Chem.* **1996**, 61, (10), 3490-3495.
- 498. Chen, Y.; Snyder, J. K., Opening norbornadiene homo Diels-Alder adducts to bicyclic systems. *Tetrahedron Lett.* **1997**, 38, (9), 1477-1480.
- 499. Chen, Y.; Snyder, J. K., Opening the [4 + 2 + 2] Cycloadducts of Bicyclo[2.2.1]hepta-2,5-dienes (Norbornadienes) to Cis-Fused Bicyclo[5.3.0]decanes. J. Org. Chem. 2001, 66, (21), 6943-6957.
- 500. Bachurin, S. O.; Shevtsova, E. P.; Kireeva, E. G.; Oxenkrug, G. F.; Sablin, S. O., Mitochondria as a Target for Neurotoxins and Neuroprotective Agents. *Ann. N. Y. Acad. Sci.* **2003**, 993, (1), 334-344.
- 501. Grigor'ev, V. V.; Dranyi, O. A.; Bachurin, S. O., Comparative Study of Action Mechanisms of Dimebon and Memantine on AMPA- and NMDA-Subtypes Glutamate Receptors in Rat Cerebral Neurons. *Bull. Exp. Biol. Med.* 2003, 136, (5), 474-477.
- 502. Bachurin, S.; Bukatina, E.; Lermontova, N.; Tkachenko, S.; Afanasiev, A.; Grigoriev, V.; Grigorieva, I.; Ivanov, Y. U.; Sablin, S.; Zefirov, N., Antihistamine Agent Dimebon As a Novel Neuroprotector and a Cognition Enhancer. *Ann. N. Y. Acad. Sci.* **2001**, 939, (1), 425-435.
- 503. Lermontova, N. N.; Redkozubov, A. E.; Shevtsova, E. F.; Serkova, T. P.; Kireeva,
   E. G.; Bachurin, S. O., Dimebon and Tacrine Inhibit Neurotoxic Action of β-

Amyloid in Culture and Block L-type Ca2+ Channels. *Bull. Exp. Biol. Med.* **2001**, 132, (5), 1079-1083.

- 504. Lermontova, N. N.; Lukoyanov, N. V.; Serkova, T. P.; Lukoyanova, E. A.; Bachurin, S. O., Dimebon improves learning in animals with experimental Alzheimer's disease. *Bull. Exp. Biol. Med.* **2000**, 129, (6), 544-546.
- 505. Bachurin, S. O.; Grigoriev, V. V. Methods and compositions for slowing aging. 2004-US410812005055951, 20041208., 2005.
- 506. Bachurin, S. O.; Grigoriev, V. V.; Morozova, M. A.; Beniashvili, A. G. Methods and compositions using hydrogenated pyrido[4,3-b]indoles for treating schizophrenia. 2007-US21172007087425, 20070125., 2007.
- 507. Hung, D. Methods and compositions for treating Huntington's disease using hydrogenated pyrido[4,3-b]indoles. 2006-US390772007041697, 20061004., 2007.
- 508. Lomberget, T.; Chataigner, I.; Bouyssi, D.; Maddaluno, J.; Balme, G., Diels-Alder cycloadditions of functionalized (Z)-1-benzylidene-2-methylene cyclohexanes: the beneficial effect of high pressure. *Tetrahedron Lett.* **2004**, 45, (17), 3437-3441.
- 509. Haber, M.; Pindur, U., N-benzoylindole-2,3-quinodimethane: Diels-Alder reactivity and synthetic applications for annellated indoles. *Tetrahedron* **1991**, 47, (10-11), 1925.
- 510. Saroja, B.; Srinivasan, P. C., A simple route to indole-2,3-quinodimethane a facile synthesisof carbazoles. *Tetrahedron Lett.* **1984**, 25, (47), 5429.
- 511. Marinelli, E. R., Generation and cycloaddition reactions of indole-2,3quinodimethanes. *Tetrahedron Lett.* **1982**, 23, (27), 2745.
- 512. Vice, S. F.; Nandin de Carvalho, H.; Taylor, N. G.; Dmitrienko, G. I., Indole-2,3quinodimethanes: Generation, reactivity and regioselective control of intermolecular Diels-Alder reactions. *Tetrahedron Lett.* **1989**, 30, (52), 7289.
- 513. Vice, S. F.; Copeland, C. R.; Forsey, S. P.; Dmitrienko, G. I., Selective side-chain bromination of n-acyl-2,3-dialkylindoles: synthesis and chemical modification of pyrrolo[1,2-a]indoles. *Tetrahedron Lett.* **1985**, 26, (43), 5253-5256.