

Identifiability and numerical algebraic geometry

Jonathan Hauenstein

Publication Date

08-12-2023

License

This work is made available under a Exclusive rights in copyrighted work license and should only be used in accordance with that license.

Citation for this work (American Psychological Association 7th edition)

Hauenstein, J. (2018). *Identifiability and numerical algebraic geometry* (Version 1). University of Notre Dame. <https://doi.org/10.7274/R03T9F91>

This work was downloaded from CurateND, the University of Notre Dame's institutional repository.

For more information about this work, to report or an issue, or to preserve and share your original work, please contact the CurateND team for assistance at curate@nd.edu.

Identifiability and numerical algebraic geometry

Daniel J. Bates^{*} Jonathan D. Hauenstein[†] Nicolette Meshkat[‡]

March 11, 2018

Abstract

A common problem when analyzing models, such as a mathematical modeling of a biological process, is to determine if the unknown parameters of the model can be determined from given input-output data. Identifiable models are models such that the unknown parameters can be determined to have a finite number of values given input-output data. The total number of such values over the complex numbers is called the identifiability degree of the model. Unidentifiable models are models such that the unknown parameters can have an infinite number of values given input-output data. For unidentifiable models, a set of identifiable functions of the parameters are sought so that the model can be reparametrized in terms of these functions yielding an identifiable model. In this work, we use numerical algebraic geometry to determine if a model is identifiable or unidentifiable. For identifiable models, we present a novel approach to compute the identifiability degree. For unidentifiable models, we present a novel numerical differential algebra technique aimed at computing a set of algebraically independent identifiable functions. Several examples are used to demonstrate the new techniques.

Keywords: Identifiability, Numerical Algebraic Geometry, Numerical Differential Algebra

1 Introduction

Parameter identifiability analysis for dynamical system models consisting of ordinary differential equations (ODEs) addresses the question of which unknown parameters can be determined from given input-output data. In this paper, we address *structural* identifiability, which concerns whether the parameters of a model can be determined from *perfect* input-output data, i.e., noise-free and of any time duration required. This is a necessary condition for the *practical* or *numerical* identifiability problem, which involves parameter estimation with real, and often noisy, data. For this reason, structural identifiability is often referred to as *a priori* identifiability [7]. Even if a model fails to be structurally identifiable, some useful information about the parameters can still be determined, which is the main motivation for this paper.

^{*}Department of Mathematics, Colorado State University, Fort Collins, CO, 80523–1874, USA (bates@math.colostate.edu, www.math.colostate.edu/~bates). Partially supported by NSF grants ACI-1440467 and DMS-1719658.

[†]Department of Applied and Computational Mathematics and Statistics, University of Notre Dame, Notre Dame, IN 46556, USA (hauenstein@nd.edu, www.nd.edu/~jhauenst). Partially supported by Army YIP W911NF-15-1-0219, Sloan Research Fellowship BR2014-110 TR14, and NSF grant ACI-1460032.

[‡]Department of Mathematics and Computer Science, Santa Clara University, Santa Clara, CA, 95053, USA (nmeshkat@scu.edu, math.scu.edu/~nmeshkat/). Partially supported by the Luce Foundation.

There are two possible outcomes of the structural identifiability check of a mathematical model. If the parameters of the model have a unique or finite number of values given input-output data, then the model and its parameters are said to be *identifiable*. However, if some subset of the parameters can take on an infinite number of values and yet yield the same input-output data, then the model and this subset of parameters are called *unidentifiable*. In the latter case, we attempt to find a set of *identifiable functions* of the parameters. These can then be used to reparameterize the model and also to give additional insight into which parameters should be experimentally measured [24].

Several methods have been proposed to find identifiable functions. In linear models, this can be done using the transfer function method [6]. However, in nonlinear models, the problem has been more challenging with only *ad hoc* methods proposed, e.g., [12, 15, 24]. For example, the approach in [24] requires the calculation of many Gröbner bases and can thus be computationally expensive. It should be noted, however, that even in the linear case, the identifiable functions of parameters found using the transfer function method are not necessarily (and are usually not) the *simplest* identifiable functions of parameters. Since our goal is to reparametrize a model over identifiable functions of the parameters, simpler functions are preferred.

In this paper, we use techniques from numerical algebraic geometry (e.g., see [4, 40] for a general overview) to investigate both identifiable and unidentifiable models. For an identifiable model, we compute the finite number of values of the parameters given input-output data. The total number of such values over the complex numbers is called the *identifiability degree* which is computed in two ways. The first method relies on differential algebra tools to first generate the *input-output equations* while the second does not utilize these equations.

For unidentifiable models, we also introduce two novel approaches for finding identifiable functions of the parameters. The first method relies on knowing the *input-output equations* and uses them to find *globally identifiable* functions of parameters, as in [24]. In the case where these input-output equations cannot be calculated using conventional differential algebra techniques, we also introduce a method to compute *locally identifiable* functions of parameters. This combination of numerical algebraic geometry and differential algebra could be thought of as *numerical differential algebra*. We demonstrate our methods on various models.

The layout of our paper is as follows. Section 2 provides details related specifically to identifiability. For an identifiable model, Section 3 describes the computation of the identifiability degree. For unidentifiable models, Section 4 focuses on the use of sampling and exactness recovery methods to construct identifiable functions. Several examples are considered in Section 5.¹

2 Identifiability

We consider ODE models of the form:

$$\begin{aligned}\dot{\mathbf{x}}(t) &= \mathbf{f}(\mathbf{x}(t), \mathbf{p}, \mathbf{u}(t), t) \\ \mathbf{y}(t) &= \mathbf{g}(\mathbf{x}(t), \mathbf{p}, t)\end{aligned}\tag{1}$$

where \mathbf{f} and \mathbf{g} are vectors of rational functions, $\mathbf{x}(t)$ is the *state variable* vector, \mathbf{p} is the *parameter* vector which is assumed to be constant, $\mathbf{u}(t)$ is the *input* vector, and $\mathbf{y}(t)$ is the *output* vector. In the following, only the input $\mathbf{u}(t)$ and output $\mathbf{y}(t)$ vectors are assumed to be known, i.e., the state variables $\mathbf{x}(t)$ and the parameters \mathbf{p} are unknown.

¹Data for computations available at <http://dx.doi.org/10.7274/R03T9F91>.

2.1 Input-output equations

One approach to determine identifiability properties of the model (1) using known input-output data is via the *input-output equations*, i.e., equations that relate the input $\mathbf{u}(t)$, output $\mathbf{y}(t)$, and parameters \mathbf{p} . Thus, the input-output equations result from eliminating the state variables $\mathbf{x}(t)$. Several methods have been proposed to compute the input-output equations for nonlinear ODE models, e.g., [1, 5, 10, 15, 16, 21, 25, 31, 32, 36], including the so-called *differential algebra approach* to identifiability [21, 31, 36]. Using differential algebra, the state variables $\mathbf{x}(t)$ are eliminated using differential elimination techniques. If the number of outputs $\mathbf{y}(t)$ is m , this procedure produces m differential polynomial equations that are solely in input and output variables with rational coefficients in the parameters so that the j^{th} one can be written as

$$\sum_i c_{ji}(\mathbf{p}) \psi_i(\mathbf{u}, \mathbf{y}) = 0$$

where each $\psi_i(\mathbf{u}, \mathbf{y})$ is a differential monomial. Each $c_{ji}(\mathbf{p})$ is a rational function in the parameters \mathbf{p} , forming a collection $\mathbf{c}(\mathbf{p})$ called the *coefficients of the input-output equations*. The coefficients of each input-output equation can be determined uniquely by taking each input-output equation to be monic.

2.2 Deciding identifiability

Let m_1 denote the number of independent parameters \mathbf{p} and m_2 denote the total number of non-monic coefficients taken from all m input-output equations. Thus, we can treat the coefficients of the input-output equations as a rational map $\mathbf{c} : \mathbb{C}^{m_1} \rightarrow \mathbb{C}^{m_2}$. Identifiability refers to whether it is possible to recover the parameters of the model only by observing the relations among the input and output variables. In other words, assuming known input-output data for a sufficient number of time instances so that \mathbf{c} can theoretically be computed, identifiability asks whether it is possible to recover the parameters \mathbf{p} .

Definition 2.1. Let \mathbf{c} be the coefficients of the input-output equations for a model (1). For general $\mathbf{p} \in \mathbb{C}^{m_1}$, let

$$X_{\mathbf{p}} = \mathbf{c}^{-1}(\mathbf{c}(\mathbf{p})) = \{\mathbf{q} \in \mathbb{C}^{m_1} \mid \mathbf{c}(\mathbf{q}) = \mathbf{c}(\mathbf{p})\} \subset \mathbb{C}^{m_1}, \quad (2)$$

$\ell = \dim X_{\mathbf{p}} \geq 0$, and $k = \#X_{\mathbf{p}} \in \mathbb{N} \cup \{\infty\}$. That is, ℓ is the dimension of a general fiber of \mathbf{c} and \mathbf{c} is generically a k -to-one map when $\ell = 0$. The model (1) is *identifiable* from \mathbf{c} if $\ell = 0$, i.e., $k \in \mathbb{N}$, and *unidentifiable* if $\ell > 0$, i.e., $k = \infty$.

When identifiable, the number $k \in \mathbb{N}$ is called the *identifiability degree*. If $k = 1$, the model (1) is called *globally identifiable* and called *locally identifiable* if $1 < k < \infty$.

When unidentifiable, the number $\ell \geq 1$ is called the *dimension of unidentifiability*.

To distinguish between identifiable and unidentifiable models, one simply needs to compute the dimension ℓ of a general fiber of \mathbf{c} . As defined in [40, §13.4], the *rank* of \mathbf{c} , denoted $\text{rank } \mathbf{c}$, is the rank of the Jacobian matrix of \mathbf{c} evaluated at a general, i.e., random, $\mathbf{p} \in \mathbb{C}^{m_1}$. The *corank* of \mathbf{c} is $\text{corank } \mathbf{c} = m_1 - \text{rank } \mathbf{c}$. The following, which is [40, Lemma 13.4.1] (see also [27]), relates ℓ and $\text{corank } \mathbf{c}$.

Proposition 2.2. For a general $\mathbf{p} \in \mathbb{C}^{m_1}$, $\ell = \dim X_{\mathbf{p}}$ as defined in (2) is equal to $\text{corank } \mathbf{c}$ where \mathbf{c} is the set of coefficients of the input-output equations. In particular, the model (1) is identifiable if and only if \mathbf{c} has full rank and the dimension of unidentifiability is equal to $\text{corank } \mathbf{c}$.

In particular, Prop. 2.2 indicates a method to distinguish between identifiable and unidentifiable models provided that the coefficients \mathbf{c} of the input-output equations can be computed, which is summarized in the following pseudocode.

Method 1: Computing dimension of unidentifiability from input-output equations

Input: m_2 input-output equation coefficients $\mathbf{c}(\mathbf{q})$, depending on parameters $\mathbf{q} \in \mathbb{C}^{m_1}$.

Output: Dimension of unidentifiability $\ell = \text{corank } \mathbf{c} = \dim \mathbf{c}^{-1}(\mathbf{c}(\mathbf{p}))$ for general $\mathbf{p} \in \mathbb{C}^{m_1}$.

Choose random, complex values \mathbf{p} of parameters \mathbf{q} .

Return $\ell = \text{corank } J\mathbf{c}(\mathbf{p})$ where $J\mathbf{c}(\mathbf{p})$ is the Jacobian matrix of \mathbf{c} evaluated at \mathbf{p} .

Example 2.3. Linear compartment models are frequently used models arising in pharmacokinetics, toxicology, cell biology, physiology, and ecology [8, 9, 13, 30, 41]. The following from [27] is an example of a linear three-compartment model with input $u(t)$, output $y(t)$, state variables $\mathbf{x}(t) = (x_1(t), x_2(t), x_3(t))$, and unknown parameters $\mathbf{p} = (k_{01}, k_{02}, k_{03}, k_{12}, k_{13}, k_{21}, k_{32})$:

$$\begin{aligned} \dot{x}_1 &= -(k_{01} + k_{21})x_1 + k_{12}x_2 + k_{13}x_3 + u \\ \dot{x}_2 &= k_{21}x_1 - (k_{02} + k_{12} + k_{32})x_2 \\ \dot{x}_3 &= k_{32}x_2 - (k_{03} + k_{13})x_3 \\ y &= x_1. \end{aligned} \tag{3}$$

Figure 1 presents a pictorial representation of this model.




Figure 1: A 3-compartment model with input (represented by the arrowhead) and output (represented by the line segment with a circle at the end) in the first compartment and “leaks” from every compartment (represented by arrows leaving the compartments). Here, the input could represent a drug injection and the first compartment could represent blood, with the other two compartments representing organs, e.g., tissue and stomach. The unknown parameters represent rates of transfer from one compartment to another (drawn as arrows in the figure), or in the case of leaks, from one compartment to outside the system. The state variables represent drug concentration in the blood and organs, with output from the first compartment representing measured drug concentration in the blood.

The approach described in [26, 27] yields the input-output equation:

$$\ddot{y} - c_1(\mathbf{p})\ddot{y} + c_2(\mathbf{p})\dot{y} - c_3(\mathbf{p})y - \ddot{u} + c_4(\mathbf{p})\dot{u} - c_5(\mathbf{p})u = 0$$

where

$$\begin{aligned}
c_1(\mathbf{p}) &= E_1(-(k_{01} + k_{21}), -(k_{02} + k_{12} + k_{32}), -(k_{03} + k_{13})) \\
c_2(\mathbf{p}) &= E_2(-(k_{01} + k_{21}), -(k_{02} + k_{12} + k_{32}), -(k_{03} + k_{13})) - k_{12}k_{21} \\
c_3(\mathbf{p}) &= E_3(-(k_{01} + k_{21}), -(k_{02} + k_{12} + k_{32}), -(k_{03} + k_{13})) + k_{12}k_{21}(k_{03} + k_{13}) + k_{21}k_{13}k_{32} \\
c_4(\mathbf{p}) &= E_1(-(k_{02} + k_{12} + k_{32}), -(k_{03} + k_{13})) \\
c_5(\mathbf{p}) &= E_2(-(k_{02} + k_{12} + k_{32}), -(k_{03} + k_{13}))
\end{aligned}$$

such that $E_k(z_1, \dots, z_m)$ is the k^{th} elementary symmetric polynomial in z_1, \dots, z_m . Thus, for $\mathbf{c} = (c_1, \dots, c_5)$, it is easy to verify that $\text{rank } \mathbf{c} = 5$ and $\text{corank } \mathbf{c} = 2$ so that this model is unidentifiable with 2 dimensions of unidentifiability.

For an identifiable model, one approach to distinguish between global and local identifiability is to solve the system of equations $\mathbf{c}(\mathbf{q}) = \mathbf{c}(\mathbf{p})$ given a general point $\mathbf{p} \in \mathbb{C}^{m_1}$. If there is a unique solution, namely $\mathbf{q} = \mathbf{p}$, the model is globally identifiable. If there are a finite number of solutions, the model is locally identifiable. Such an approach, for example, is implemented in the software package DAISY [7, 35] which randomly selects a point \mathbf{p} and uses Gröbner basis methods to count the number of solutions to $\mathbf{c}(\mathbf{q}) = \mathbf{c}(\mathbf{p})$ yielding the identifiability degree. Since such an approach can only be applied when \mathbf{c} has first been computed, we will consider the following problem using numerical algebraic geometric methods.

Problem 2.4. Given a model (1), decide if it is identifiable or unidentifiable. If identifiable, determine its identifiability degree to decide if it is globally identifiable or locally identifiable.

One technique for determining whether a model is identifiable without computing \mathbf{c} is via the Exact Arithmetic Rank (EAR) approach [20]. In particular, rather than eliminating to compute the corank of \mathbf{c} , one considers projections of a system that still involves the state variables derived by replacing functions with Taylor series expansions and taking a finite-size system via the Cartan-Kuranishi Theorem that underlies differential elimination, e.g., see [33]. The following, from [19, Lemma 3], is essential for computing $\text{corank } \mathbf{c}$ without first computing \mathbf{c} .

Proposition 2.5. Let $F : \mathbb{C}^N \rightarrow \mathbb{C}^n$ be a polynomial system, $V \subset \{x \in \mathbb{C}^N \mid F(x) = 0\} \subset \mathbb{C}^N$ be irreducible of multiplicity 1 with respect to F , and $\pi(x_1, \dots, x_N) = (x_1, \dots, x_a)$ for some $a \leq N$. For general $z \in V$,

$$\dim \overline{\pi(V)} = \text{corank}_0 JF(z) - \text{corank}_a JF(z)$$

where $JF(z)$ is the Jacobian matrix of F evaluated at z and $\text{corank}_j M$ is the corank of the last $N - j$ columns of M .

Example 2.6. With the setup from Ex. 2.3, write the function $\mathbf{x}(t)$, $u(t)$, and $y(t)$ using a Taylor series expansion centered at $t = 0$, namely

$$\mathbf{x}(t) = \sum_{j=0}^{\infty} \mathbf{x}_j \cdot t^j / j!, \quad u(t) = \sum_{j=0}^{\infty} u_j \cdot t^j / j!, \quad \text{and} \quad y(t) = \sum_{j=0}^{\infty} y_j \cdot t^j / j!. \quad (4)$$

Since (3) holds for all t , one obtains equations by substituting these Taylor series expansions into (3) and taking coefficients with respect to t . For $r \geq 0$, let F_r be the system obtained by

taking coefficients of $1, t, t^2, \dots, t^r$. For this linear compartment model, the coefficients of t^j are

$$G_j = \begin{bmatrix} -(k_{01} + k_{21})x_{j1} + k_{12}x_{j2} + k_{13}x_{j3} + u_j - x_{j+1,1} \\ k_{21}x_{j1} - (k_{02} + k_{12} + k_{32})x_{j2} - x_{j+1,2} \\ k_{32}x_{j2} - (k_{03} + k_{13})x_{j3} - x_{j+1,3} \\ y_j - x_{j1} \end{bmatrix} \quad \text{so that} \quad F_r = \begin{bmatrix} G_0 \\ G_1 \\ \vdots \\ G_r \end{bmatrix}.$$

Based on the structure of each G_j , it is clear that the Jacobian matrix of F_r has full rank, namely $4(r+1)$, at every point. In fact, $F_r = 0$ defines an irreducible and smooth solution set of codimension $4(r+1)$ ($\dim = 11 + r$). We can compute a random point on this solution set by randomly selecting the following $11 + r$ values: \mathbf{p} , \mathbf{x}_0 , and u_0, \dots, u_r , and trivially computing the unique \mathbf{x}_{j+1} and y_j sequentially for $j = 0, \dots, r$ via $G_j = 0$.

Next, one treats the coefficients of the input $u(t)$ and output $y(t)$ as constants in F_r . Thus, we have that F_r depends upon $N_r = 13 + 3r$ variables and apply Prop. 2.5 to compute

$$d_r = \text{corank}_0 JF_r(\mathbf{p}, \mathbf{x}_0, \dots, \mathbf{x}_{r+1}) - \text{corank}_7 JF_r(\mathbf{p}, \mathbf{x}_0, \dots, \mathbf{x}_{r+1})$$

since $\pi_r(\mathbf{p}, \mathbf{x}_0, \dots, \mathbf{x}_{r+1}) = \mathbf{p} \in \mathbb{C}^7$. We trivially know $d_r \geq d_{r+1}$ since F_r is a subset of F_{r+1} . Hence, $\{d_r\}_{r=0}^\infty$ is a sequence of nonincreasing nonnegative integers that stabilizes with

$$\lim_{r \rightarrow \infty} d_r = \text{corank } \mathbf{c}.$$

This limit is obtained at a finite value of r in accordance with the Cartan-Kuranishi Theorem and can be observed by checking for stabilization between the values obtained from r to $r+1$ as demonstrated in the following table:

r	N_r	$\text{corank}_0 JF_r$	$\text{corank}_7 JF_r$	d_r
0	13	9	2	7
1	16	8	1	7
2	19	7	0	7
3	22	6	0	6
4	25	5	0	5
5	28	4	0	4
6	31	3	0	3
7	34	2	0	2
8	37	2	0	2
9	40	2	0	2
10	43	2	0	2

We see that $d_7 = d_8 = 2 = \text{corank } \mathbf{c}$ and provide the extra rows to show how the entries stabilize. In particular, this confirms that (3) is unidentifiable with 2 dimensions of unidentifiability.

We summarize this computation of the dimension of unidentifiability without first explicitly computing the input-output equations \mathbf{c} in the following pseudocode.

Method 2: Computing dimension of unidentifiability without input-output equations

Input: For each $r \geq 0$, system $F_r(\mathbf{q}, \mathbf{x}, \mathbf{u}, \mathbf{y})$ consisting of the coefficients of $1, t, t^2, \dots, t^r$ and general point \mathbf{z}_r such that $F_r(\mathbf{z}_r) = 0$ where \mathbf{q} consists of m_1 parameters.

Output: Dimension of unidentifiability $\ell = \text{corank } \mathbf{c} = \dim \mathbf{c}^{-1}(\mathbf{c}(\mathbf{p}))$ for general $\mathbf{p} \in \mathbb{C}^{m_1}$.
For $r = 0, 1, 2, \dots$

Compute $d_r = \text{corank}_0 JF_r(\mathbf{z}_r) - \text{corank } J_{m_1} F_r(\mathbf{z}_r)$

If either $d_r = 0$ or $r > 0$ and $d_r = d_{r-1}$, return d_r .

Such an approach naturally extends to problems when the parameters and initial conditions are restricted to an irreducible component by simply appending to F_r the requested constraints and taking the test points to be general on the corresponding irreducible component. The following demonstrates this on [34, Ex. 1].

Example 2.7. Consider the following three-compartment model [42]:

$$\begin{aligned} \dot{x}_1 &= p_{13}x_3 + p_{12}x_2 - p_{21}x_1 + u \\ \dot{x}_2 &= p_{21}x_1 - p_{12}x_2 \\ \dot{x}_3 &= -p_{13}x_3 \\ y &= x_2 \end{aligned} \tag{5}$$

with state variables $\mathbf{x}(t) = (x_1(t), x_2(t), x_3(t))$, input $u(t)$, output $y(t)$, and unknown parameters $\mathbf{p} = (p_{12}, p_{13}, p_{21})$. Using a similar setup from Ex. 2.6 summarized in Method 2, the following table shows that the model (5) is identifiable:

r	N_r	$\text{corank}_0 JF_r$	$\text{corank}_3 JF_r$	d_r
0	9	5	2	3
1	12	4	1	3
2	15	3	0	3
3	18	2	0	2
4	21	1	0	1
5	24	0	0	0

Let F'_r be the system obtained by adding the constraint $x_3(0) = 0$ to F_r . The following table shows that the model (5) is now unidentifiable with one dimension of unidentifiability:

r	N_r	$\text{corank}_0 JF'_r$	$\text{corank}_3 JF'_r$	d_r
0	9	4	1	3
1	12	3	0	3
2	15	2	0	2
3	18	1	0	1
4	21	1	0	1

2.3 Identifiable functions

When a model (1) is unidentifiable, one can ask for functions of the parameters \mathbf{p} which are actually functions of the coefficients $\mathbf{c}(\mathbf{p})$ of the input-output equations. Such functions are called *identifiable functions*. For example, every element of \mathbf{c} is itself an identifiable function. This is algebraically formalized in the following.

Definition 2.8. Let \mathbf{c} be as above. A real-valued function $f(\mathbf{p})$ is *identifiable* if the field extension $\mathbb{R}(f, \mathbf{c})/\mathbb{R}(\mathbf{c})$ is an algebraic field extension.

One can also consider the global and local identifiability of functions.

Definition 2.9. Let \mathbf{c} be as above and f be an identifiable function. The function f is called *globally identifiable* from \mathbf{c} if there exists a function ϕ such that $\phi \circ \mathbf{c} = f$. The function f is called *locally identifiable* from \mathbf{c} if there exists a multi-valued function ξ such that $\xi \circ \mathbf{c} = f$.

Example 2.10. With the setup from Ex. 2.3, the function $f(\mathbf{p}) = k_{01} + k_{21}$ is globally identifiable with $f = \phi \circ \mathbf{c}$ where $\phi(x_1, \dots, x_5) = x_4 - x_1$, i.e., $f = c_4 - c_1$. The function $g(\mathbf{p}) = k_{02} + k_{12} + k_{32}$ is locally identifiable with $g^2 + c_4g + c_5 = 0$, i.e., $\xi \circ \mathbf{c} = g$ where

$$\xi(x_1, \dots, x_5) = \frac{-x_4 \pm \sqrt{x_4^2 - 4x_5}}{2}.$$

When a model is unidentifiable with $\ell = \text{corank } \mathbf{c}$ dimensions of unidentifiability, the goal is to compute $d = \text{rank } \mathbf{c}$ algebraically independent identifiable functions. The problem of finding d “nice” algebraically independent identifiable functions can be described in the following way, where “nice” could be taken to mean low degree, sparse, or are easy to interpret in terms of the model, depending on the context.

Problem 2.11. For rational functions \mathbf{c} with $d = \text{rank } \mathbf{c}$, compute a “nice” transcendence basis of the field extension $\mathbb{R}(\mathbf{c})/\mathbb{R}$ given by the functions f_1, \dots, f_d .

One way to locate identifiable functions is by computing Gröbner bases with respect to various elimination orderings of the ideal $\langle \mathbf{c}(\mathbf{q}) - \mathbf{c}(\mathbf{p}) \rangle$. This approach is described in [24, 28] and has been implemented in the web application COMBOS [28]. In addition to requiring \mathbf{c} , e.g., computed using differential elimination techniques, the biggest disadvantage of this method is that Gröbner basis computations can be computationally expensive. Thus, COMBOS can fail even for relatively simple models. Alternatively, the program DAISY [7, 35] can sometimes be used to find identifiable functions. Specifically, the DAISY program gives the solution to the system of equations $\mathbf{c}(\mathbf{q}) = \mathbf{c}(\mathbf{p})$ for a randomly chosen numerical point \mathbf{p} . Sometimes one can algebraically manipulate the solution to obtain functions of the form $f(\mathbf{q}) = f(\mathbf{p})$, but there are many cases where this cannot be done [24, 28]. Nonetheless, if one is able to obtain such f , the following shows that they are indeed identifiable functions.

Proposition 2.12. *If $f(\mathbf{q}) - f(\mathbf{p})$ is an element of the ideal $I = \langle \mathbf{c}(\mathbf{q}) - \mathbf{c}(\mathbf{p}) \rangle \subset \mathbb{R}[\mathbf{p}, \mathbf{q}]$, then f is an identifiable function. That is, if f is constant on irreducible components of generic fibers of \mathbf{c} , then f is an identifiable function.*

Proof. If $f(\mathbf{q}) - f(\mathbf{p})$ is contained in I , then the dimension of the image of the combined map (\mathbf{c}, f) is equal to the dimension of the image of the map \mathbf{c} . In other words, the field extension $\mathbb{R}(f, \mathbf{c})/\mathbb{R}(\mathbf{c})$ is an algebraic field extension showing that f is identifiable. \square

2.4 Reparametrization and other uses of identifiable functions

If one can solve Problem 2.11, one then tries to use the new basis to reparametrize the model. In [26], a method to find identifiable scaling reparametrizations is given for a certain class of linear compartment models where the identifiable functions are monomials. Currently, there is no

general approach to find identifiable reparametrizations and, for most models, the reparametrizations are found using *ad hoc* approaches which work on a case-by-case basis.

Even if a reparametrization cannot be found, identifiable functions have other important uses. From the identifiable functions, one can determine which parameters need to be known in order to render the entire model identifiable. While this information can also be determined from the solution to the system of equations $\mathbf{c}(\mathbf{q}) = \mathbf{c}(\mathbf{p})$, identifiable functions give us additional information if only a subset of those parameters can be determined. In other words, we can obtain a simpler set of identifiable functions of parameters if a subset of the parameters are known and, perhaps, for this new set of identifiable functions, computing an identifiable reparametrization is possible. This is the case for Ex. 5.1 below where knowledge of either the pair (a_{34}, a_{43}) or the pair (a_{33}, a_{44}) renders all the identifiable functions to be monomials, in which case the method in [26] can be used to find an identifiable scaling reparametrization.

3 Computing identifiability degree

For a model (1) that is identifiable, Problem 2.4 can be solved by computing the identifiability degree $k \in \mathbb{N}$ in order to distinguish between globally identifiable ($k = 1$) and locally identifiable ($k > 1$) models. k is simply the number of solutions of $\mathbf{c}(\mathbf{q}) = \mathbf{c}(\mathbf{p})$ for general \mathbf{p} , where \mathbf{c} is the collection of coefficients of the input-output equations. As mentioned in §2.2, the software package DAISY [7, 35] uses such an approach with Gröbner basis methods to count the number of solutions. One could also use numerical homotopy methods, e.g., as summarized in [4, 40], to compute k , as illustrated in the following example.

Example 3.1. As shown in Ex. 2.3, the model (3) has 2 dimensions of unidentifiability. With the aim of constructing an identifiable model, we modify (3) by adding the extra constraints $k_{01} = k_{03} = 0$ yielding a new model with only one leak parameter k_{02} . The coefficients \mathbf{c} of the input-output equation for this simplified model are

$$\mathbf{c}(k_{02}, k_{12}, k_{13}, k_{21}, k_{32}) = \begin{bmatrix} k_{02} + k_{12} + k_{13} + k_{21} + k_{32} \\ k_{02}k_{13} + k_{02}k_{21} + k_{12}k_{13} + k_{13}k_{21} + k_{13}k_{32} + k_{21}k_{32} \\ k_{02}k_{13}k_{21} \\ k_{02} + k_{12} + k_{13} + k_{32} \\ k_{02}k_{13} + k_{12}k_{13} + k_{13}k_{32} \end{bmatrix}, \quad (6)$$

which is easily seen to have rank 5, i.e., the model is identifiable. For general $\alpha_i \in \mathbb{C}$, the system

$$\mathbf{c}(k_{02}, k_{12}, k_{13}, k_{21}, k_{32}) - \mathbf{c}(\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5) = 0 \quad (7)$$

consists of 5 equations (1 cubic, 2 quadratic, and 2 linear) in 5 variables. Using a total degree homotopy (see [4, 40] for more details), one tracks $3 \cdot 2^2 \cdot 1^2 = 12$, i.e., the total degree of (7), solution paths. Tracking these paths with homotopy continuation, e.g., via Bertini [3], yields 2 solutions to (7). One can also use a Gröbner basis computation to see that (7) has 2 solutions. These computations show that the model (3) with $k_{01} = k_{03} = 0$ is locally identifiable with identifiability degree of 2.

We summarize this most basic approach for computing the identifiability degree when the input-output equations are known in the following pseudocode.

Method 3: Computing identifiability degree from input-output equations (direct solving)

Input: m_2 input-output equation coefficients $\mathbf{c}(\mathbf{q})$, depending on parameters $\mathbf{q} \in \mathbb{C}^{m_1}$ for which $\text{corank } \mathbf{c} = 0$, i.e., corresponding model is identifiable.

Output: Identifiability degree $k \in \mathbb{N}$.

Choose random, complex values \mathbf{p} of parameters \mathbf{q} .

Use homotopy continuation to compute $Z = \{\mathbf{q} \in \mathbb{C}^{m_1} \mid \mathbf{c}(\mathbf{q}) = \mathbf{c}(\mathbf{p})\}$.

Return $k = \#Z$.

Rather than using a direct global solving method which is based on knowing the coefficients \mathbf{c} , we next consider an alternative approach based on monodromy computations in numerical algebraic geometry that also can be used without computing \mathbf{c} . We first describe the approach when \mathbf{c} is known and then extend to the case when \mathbf{c} is not explicitly computed.

3.1 Identifiability degree from input-output equations

Suppose that (1) is identifiable with identifiability degree $k \in \mathbb{N}$ and \mathbf{c} is the set of coefficients of the input-output equations. Following §2.2, let m_1 be the number of independent parameters \mathbf{p} and m_2 be the number of entries in \mathbf{c} so that $\mathbf{c} : \mathbb{C}^{m_1} \rightarrow \mathbb{C}^{m_2}$. Assume that the model is identifiable so that $\text{corank } \mathbf{c} = 0$ and $\text{rank } \mathbf{c} = \dim X = m_1$ where $X = \overline{\mathbf{c}(\mathbb{C}^{m_1})} \subset \mathbb{C}^{m_2}$. The continuity of \mathbf{c} yields that X is irreducible. The graph of \mathbf{c} , namely

$$\text{Graph}(\mathbf{c}) = \{(\mathbf{p}, \mathbf{c}(\mathbf{p})) \mid \mathbf{p} \in \mathbb{C}^{m_1}\} \subset \mathbb{C}^{m_1} \times \mathbb{C}^{m_2}$$

is also irreducible of dimension m_1 . In fact, for the projection map $\pi : \mathbb{C}^{m_1} \times \mathbb{C}^{m_2} \rightarrow \mathbb{C}^{m_2}$, we know that $X = \overline{\pi(\text{Graph}(\mathbf{c}))}$ and π restricted to $\text{Graph}(\mathbf{c})$ is generically a k -to-1 map.

One can compute k via a pseudowitness point set [19] for X . To that end, let $\mathcal{L}_2 \subset \mathbb{C}^{m_2}$ be a general linear space of codimension m_1 . The finite set $W = \text{Graph}(\mathbf{c}) \cap (\mathbb{C}^{m_1} \times \mathcal{L}_2)$ is a *pseudowitness point set* for X with respect to $\text{Graph}(\mathbf{c})$ and π where $\#W = k \cdot \deg X$ and $\#\pi(W) = \deg X$, i.e., $k = \#W / \#\pi(W)$. In order to compute W , we follow the approach in [17] using monodromy loops [39], as follows.

We first note that it is trivial to construct one point $w \in W$ as follows. One first selects a general point $(\mathbf{p}, \mathbf{c}(\mathbf{p})) \in \text{Graph}(\mathbf{c})$ and then constructs a general linear space $\mathcal{L}_2 \subset \mathbb{C}^{m_2}$ of codimension m_1 that passes through $\mathbf{c}(\mathbf{p})$. Hence, $w = (\mathbf{p}, \mathbf{c}(\mathbf{p})) \in W$.

Next, the irreducibility of $\text{Graph}(\mathbf{c})$ ensures that pairs of points in W are connected via smooth paths on $\text{Graph}(\mathbf{c})$. We aim to discover such connecting paths using random monodromy loops. For $t \in [0, 1]$, let $\mathcal{L}(t)$ be a smooth path consisting of general linear spaces of codimension m_1 in \mathbb{C}^{m_2} such that $\mathcal{L}(0) = \mathcal{L}(1) = \mathcal{L}_2$. Hence, this defines paths $w(t)$ defined by $\text{Graph}(\mathbf{c}) \cap (\mathbb{C}^{m_1} \times \mathcal{L}(t))$ where $w(1) \in W$ is known. Homotopy continuation computes the endpoint $w(0)$, which must also be a point in W . If $w(0) \neq w(1)$, the resulting loop has produced a nontrivial monodromy action and potentially yielded a previously unknown point in W .

Example 3.2. For $\mathbf{c} : \mathbb{C}^5 \rightarrow \mathbb{C}^5$ in (6), we know that $X = \overline{\mathbf{c}(\mathbb{C}^5)} = \mathbb{C}^5$, i.e., $\deg X = 1$. Hence, we have that the identifiability degree $k = \#W$ where W is a pseudowitness point set for X .

For illustrative purposes, consider $\mathbf{p} = (-1, -2, 5, -1, -3)$ with $\mathbf{c}(\mathbf{p}) = (-2, -31, 5, -1, -30)$ so $\mathcal{L}_2 = \{(-2, -31, 5, -1, -30)\}$ has codimension 5 with $\mathbf{c}(\mathbf{p}) \in \mathcal{L}_2$. Consider the loop

$$\mathcal{L}(t) = \{(-2, -31 - 15s(t), 5 + 5s(t), -1, -30 + 35s(t))\}$$

where $s(t) = 1 - e^{2\pi i(1-t)}$ and $i = \sqrt{-1}$. Hence, $\mathcal{L}(t)$ is a loop with $\mathcal{L}(0) = \mathcal{L}(1) = \mathcal{L}_2$. For the path $w(t) \in \text{Graph}(\mathbf{c}) \cap (\mathbb{C}^5 \times \mathcal{L}(t))$ with $w(1) = (\mathbf{p}, \mathbf{c}(\mathbf{p}))$, we have $w(0) = (\mathbf{q}, \mathbf{c}(\mathbf{q}))$ where $\mathbf{q} = (5/6, -2, -6, -1, 37/6)$ and $\mathbf{c}(\mathbf{q}) = \mathbf{c}(\mathbf{p})$ showing $\{w(0), w(1)\} \subset W$ and $k = \#W \geq 2$.

Running finitely many random monodromy loops necessarily yields a set $\widehat{W} \subset W$ that may fail to achieve the goal of equality. However, information about the model can be obtained even if $\widehat{W} \subsetneq W$. For example, if $(\mathbf{p}_1, \mathbf{c}(\mathbf{p}_1))$ and $(\mathbf{p}_2, \mathbf{c}(\mathbf{p}_2))$ are known points in W with $\mathbf{c}(\mathbf{p}_1) = \mathbf{c}(\mathbf{p}_2)$ and $\mathbf{p}_1 \neq \mathbf{p}_2$, then one knows the identifiability degree is larger than 1, i.e., the model is locally identifiable. A heuristic stopping criterion for when $\widehat{W} = W$ provided in [17] is simply to have many different random monodromy loops yielding no new points. We use trace tests [18, 38] to provide a stopping criterion to recognize when $\widehat{W} = W$.

Trace tests can be summarized as follows. Suppose that $Y \subset \mathbb{C}^n$ is irreducible of dimension r . Let $\mathcal{L}(t)$ be a general family of codimension r parallel linear spaces. For $Z \subset Y \cap \mathcal{L}(0)$, the trace test of [38] shows that $Z = Y \cap \mathcal{L}(0)$, i.e., $\#Z = \deg Y$, if and only if $\sum_{z \in Z} p_z(t)$ is linear in t where $p_z(t) \in Y \cap \mathcal{L}(t)$ with $p_z(0) = z$.

Returning to the setup from above, since X is irreducible, we can apply this so-called “classical” trace test to verify that $\pi(\widehat{W}) = \pi(W)$, i.e., $\#\pi(\widehat{W}) = \#\pi(W) = \deg X$.

To verify that $\widehat{W} = W$, we need to compute additional points to use the multihomogeneous trace test of [18]. Let \mathcal{M}_2 and \mathcal{H}_2 be general linear spaces in \mathbb{C}^{m_2} of codimension $m_1 - 1$ and 1, respectively, with $\mathcal{L}_2 = \mathcal{M}_2 \cap \mathcal{H}_2$. By [37, Thm. 3.42], $\mathcal{C} = \text{Graph}(\mathbf{c}) \cap (\mathbb{C}^{m_1} \times \mathcal{M}_2)$ is an irreducible curve in $\mathbb{C}^{m_1} \times \mathbb{C}^{m_2}$, say, of multidegree (d_1, d_2) . In particular, $d_2 = \#(\mathcal{C} \cap (\mathbb{C}^{m_1} \times \mathcal{H}_2)) = \#W$ meaning that a trace test for validating the completeness of a multihomogeneous witness point set for \mathcal{C} will show the completeness of W . To that end, let $\mathcal{H}_1 \subset \mathbb{C}^{m_1}$ be a general hyperplane and $\mathcal{H}(t)$ be a general family of parallel $(1, 1)$ hypersurfaces in $\mathbb{C}^{m_1} \times \mathbb{C}^{m_2}$ with

$$\mathcal{H}(0) = (\mathcal{H}_1 \times \mathbb{C}^{m_2}) \cup (\mathbb{C}^{m_1} \times \mathcal{H}_2). \quad (8)$$

For $Z \subset \mathcal{C} \cap \mathcal{H}(0)$, the trace test of [18] shows that $Z = \mathcal{C} \cap \mathcal{H}(0)$ if and only if $\sum_{z \in Z} \sigma(p_z(t))$ is linear in t where $p_z(t) \in \mathcal{C} \cap \mathcal{H}(t)$ and $\sigma : \mathbb{C}^{m_1} \times \mathbb{C}^{m_2} \rightarrow \mathbb{C}^{m_1 m_2 + m_1 + m_2}$ is the Segre embedding.

To make these monodromy and trace test computations more efficient, see [11, 14].

Example 3.3. To show that Ex. 3.2 computed both points in W , i.e., the degree of identifiability $k = 2$, for illustrative purposes, we consider the following three linear spaces in \mathbb{C}^5 :

$$\begin{aligned} \mathcal{H}_1 &= \{4x_1 + 5x_2 - 2x_3 + 4x_4 - 2x_5 - 1 = 0\}, & \mathcal{H}_2 &= \{2y_1 + 4y_2 - y_3 - 6y_4 - 4y_5 + 7 = 0\}, \\ &\text{and } \mathcal{M}_2 &= \{(-2 - 5r, -31 - 3r, 5 - 3r, -1 + 2r, -30 + 4r) \mid r \in \mathbb{C}\}, \end{aligned}$$

with $\mathcal{L}_2 = \mathcal{M}_2 \cap \mathcal{H}_2$. We take

$$\mathcal{H}(t) = \{(4x_1 + 5x_2 - 2x_3 + 4x_4 - 2x_5 - 1)(2y_1 + 4y_2 - y_3 - 6y_4 - 4y_5 + 7) - 2t = 0\} \subset \mathbb{C}^5 \times \mathbb{C}^5$$

which satisfies (8). The irreducible curve $\mathcal{C} = \text{Graph}(\mathbf{c}) \cap (\mathbb{C}^5 \times \mathcal{M}_2)$ has multidegree $(5, 2)$, which is verified using the multihomogeneous trace test applied to $\mathcal{C} \cap \mathcal{H}(t)$. This yields $k = 2$.

We summarize this computation in the following pseudocode.

Method 4: Computing identifiability degree from input-output equations (monodromy)

Input: m_2 input-output equation coefficients $\mathbf{c}(\mathbf{q})$, depending on parameters $\mathbf{q} \in \mathbb{C}^{m_1}$ for which $\text{corank } \mathbf{c} = 0$, i.e., model is identifiable, and an integer *maxUselessLoops*.

Output: Identifiability degree $k \in \mathbb{N}$ or *error* along with a lower bound on k if the number of loops in a row that do not yield any new points is more than *maxUselessLoops*.

Choose random, complex values \mathbf{p} of parameters \mathbf{q} and compute $\mathbf{c}(\mathbf{p})$.

Form $\mathbf{w} = (\mathbf{p}, \mathbf{c}(\mathbf{p}))$ and $\mathbf{W} = \{\mathbf{w}\}$.

Construct general linear space $\mathcal{L}_2 \subset \mathbb{C}^{m_2}$ of codimension m_1 that passes through $\mathbf{c}(\mathbf{p})$.

Set *numUselessLoops* = 0.

While *numUselessLoops* < *maxUselessLoops*

 Increment *numUselessLoops* = *numUselessLoops* + 1.

 Construct a general loop of linear spaces $\mathcal{L}(t)$ such that $\mathcal{L}(0) = \mathcal{L}(1) = \mathcal{L}_2$.

 For each $\mathbf{w} \in \mathbf{W}$

 Use homotopy continuation applied to the homotopy $\text{Graph}(\mathbf{c}) \cap (\mathbb{C}^{m_1} \times \mathcal{L}(t))$ to track from \mathbf{w} at $t = 1$ to $t = 0$ yielding \mathbf{w}' .

 If $\mathbf{w}' \notin \mathbf{W}$

 Update $\mathbf{W} = \{\mathbf{W}, \mathbf{w}'\}$ and *numUselessLoops* = 0.

 If trace test passes, return $k = \#\mathbf{W}$.

Return *error* with $k = \#\mathbf{W}$.

The advantage to using such a monodromy approach is that the structure of \mathbf{c} may be such that k is small but this structure is not known *a priori* meaning that a homotopy for solving $\mathbf{c}(\mathbf{q}) = \mathbf{c}(\mathbf{p})$ requires tracking many homotopy paths. The disadvantage is that many monodromy loops may be needed to find all points necessary for the trace test to validate completeness when k is large.

3.2 Identifiability degree without input-output equations

In §3.1, we computed the degree of a general fiber of a generically finite-to-one coefficient map. This is based on the fact that one has the same input-output equation if and only if the coefficients agree. However, when we are using a truncated system as described in §2.2, namely F_r which depends upon the parameters \mathbf{p} , input $\mathbf{U} = \{\mathbf{u}_0, \dots, \mathbf{u}_r\}$, output $\mathbf{Y} = \{\mathbf{y}_0, \dots, \mathbf{y}_r\}$, and state variables $\mathbf{X} = \{\mathbf{x}_0, \dots, \mathbf{x}_{r+1}\}$, it provides necessary conditions to have the same input-output as shown in the following example.

Example 3.4. The following model is a modification of an HIV model from [29]:

$$\begin{aligned} \dot{x}_1 &= p_1 - p_2 x_1 - p_3 x_1 x_3 \\ \dot{x}_2 &= p_3 x_1 x_3 - p_4 x_2 \\ \dot{x}_3 &= p_1 p_4 x_2 - p_5 x_3 \\ y &= x_3 \end{aligned} \tag{9}$$

As in §3.1, the following table shows that the system F_7 provides the model (9) is identifiable:

r	$\text{corank}_0 JF_r$	$\text{corank}_5 JF_r$	d_r
0	7	2	5
1	6	1	5
2	5	0	5
3	4	0	4
4	3	0	3
5	2	0	2
6	1	0	1
7	0	0	0

For example, consider the sufficiently general truncated output

$$\mathbf{Y} = (y_0, \dots, y_7) = (0.5, -0.03, -0.15, -0.2, -0.2, -0.17, -0.16, -0.15).$$

We know that there are finitely many values of the parameters \mathbf{p} which yield this output. Monodromy yields the following 12 values of the parameters (listed to four decimal places):

p_1	p_2	p_3	p_4	p_5	y_8
± 0.1253	-2.4825	4.4249	-0.9210	-0.2137	0.1706
± 0.2602	-2.4825	4.4249	-0.2137	-0.9210	0.1706
$0.3023 \pm 0.0779i$	$-3.5234 \pm 0.5105i$	$4.2201 \pm 1.9168i$	$-1.3367 \mp 0.0298i$	$-0.1080 \mp 0.2292i$	$0.1107 \mp 0.4040i$
$-0.3023 \pm 0.0779i$	$-3.5234 \mp 0.5105i$	$4.2201 \mp 1.9168i$	$-1.3367 \pm 0.0298i$	$-0.1080 \pm 0.2292i$	$0.1107 \pm 0.4040i$
$0.6847 \pm 0.2133i$	$-3.5234 \mp 0.5105i$	$4.2201 \mp 1.9168i$	$-0.1080 \pm 0.2292i$	$-1.3367 \pm 0.0298i$	$0.1107 \pm 0.4040i$
$-0.6847 \pm 0.2133i$	$-3.5234 \pm 0.5105i$	$4.2201 \pm 1.9168i$	$-0.1080 \mp 0.2292i$	$-1.3367 \mp 0.0298i$	$0.1107 \mp 0.4040i$

Table 1: 12 possible values of parameters of system F_7 from model (9).

This table shows that there are 3 distinct values of y_8 , each of which is obtained by 4 values of the parameters yielding the identifiability degree is 4.

This example shows that even though F_r is enough to show identifiability, we may only need to consider a subset of the corresponding parameter values which have the same input-output.

The structure of (1) clearly shows that the solution set of $F_r = 0$ is irreducible, smooth, and parameterized by \mathbf{p} , \mathbf{U} , and \mathbf{x}_0 . Thus, it is trivial to construct a generic point $(\mathbf{p}^*, \mathbf{X}^*, \mathbf{U}^*, \mathbf{Y}^*)$ in the solution set of $F_r = 0$. From this point, we can use Prop. 2.5 to compute the dimension $d \geq 0$ of the solution set of $F_r(\mathbf{p}^*, \mathbf{X}, \mathbf{U}^*, \mathbf{Y}^*) = 0$, i.e., the dimension of the state variables. If $d > 0$, we can add d general linear slices in \mathbf{X} to F_r to reduce to the case when $d = 0$.

With this reduction, we repeatedly apply random monodromy loops to compute all values of \mathbf{p} such that there exists \mathbf{X} with

$$F_r(\mathbf{p}, \mathbf{X}, \mathbf{U}^*, \mathbf{Y}^*) = 0.$$

By testing the finitely many values of the parameters \mathbf{p} , the identifiability degree k is the number of which correspond with the same input-output. To verify the completeness, we simply apply the multihomogeneous trace test via the parameter space and the input-output space.

To save space, we exclude pseudocode for this method as it is so similar to Method 4. The primary change is that the set of coefficients \mathbf{c} is replaced by the truncated system F_r for some value of r along with an extra computation to test for the same input-output values.

Example 3.5. Reconsidering the model (9) in Ex. 3.4 which has no input, we first restrict the output space to, for illustration purposes, the sufficiently general line

$$\mathbf{Y}(s) = (s + 0.5, 4s - 0.03, 3s - 0.15, -2s - 0.2, -s - 0.2, -3s - 0.17, 3s - 0.16, 4s - 0.15).$$

Thus, we apply the multihomogeneous trace test by solving $F_7 = 0$ on this line intersected with the sufficiently general family of bilinear hyperplanes in the parameter and output space:

$$\mathcal{H}(t) = \{(2p_1 - 3p_2 - 4p_3 - p_4 - 4p_5 - 5)(3y_0 + 4y_1 + 5y_2 + y_3 + y_4 - 4y_5 + 4y_6 - y_7 - 0.42) - t = 0\}.$$

Monodromy followed by the trace test confirms that the bidegree is $(222, 12)$. Hence, the number of elements in Table 1 is complete.

We can simplify this computation, for example, by instead taking the following family

$$\mathcal{H}(t) = \{(3p_5 - 4)(3y_0 + 4y_1 + 5y_2 + y_3 + y_4 - 4y_5 + 4y_6 - y_7 - 0.42) - t = 0\}.$$

The bidegree in p_5 and the output space is $(60, 12)$ which again shows that Table 1 is complete.

4 Computing identifiable functions

A model (1) is identifiable if and only if every function of the parameters is an identifiable function. In particular, each irreducible component of a generic fiber of the coefficients \mathbf{c} of the input-output equations is a singleton for an identifiable model. Since every function of the parameter is trivially constant on each singleton, Prop. 2.2 yields that every function is identifiable. To be a globally identifiable function, it must take the same constant value on all of the irreducible components of a general fiber.

Example 4.1. With the setup from Ex. 3.4, the model (9) is identifiable with identifiability degree 4. Hence, for example, we know that $f_1 = p_4$ and $f_2 = p_5$ are both identifiable functions. From the first two rows of Table 1, we see that both f_1 and f_2 are not globally identifiable since each of them take two different values. The functions $g_1 = p_2$, $g_2 = p_3$, and $g_3 = p_4 + p_5$ are all globally identifiable since each of them take the same value at all four points.

To compute identifiable functions, we will first use numerical algebraic geometry to sample points from fibers. Then, given a finite collection of terms, we will use exactness recovery techniques, e.g., [2], or interpolation to construct identifiable functions from the sample data. Computing globally identifiable functions simply requires computing points on all irreducible components and adding additional constraints.

4.1 Sampling

In the case that input-output equations have been computed, let \mathbf{c} be the collection of coefficients of the input-output equations and suppose that $d \geq 0$ is the dimension of unidentifiability. Thus, for a given generic point \mathbf{p} , the point $\mathbf{q} = \mathbf{p}$ is a smooth point on an irreducible component $V_{\mathbf{p}}$ of dimension d of the solution set defined by $\mathbf{c}(\mathbf{q}) - \mathbf{c}(\mathbf{p}) = 0$. Hence, when $d > 0$, we can sample other points in this irreducible component as follows. Let $\mathcal{L}_{\mathbf{p}}$ be a general linear space of codimension d passing through \mathbf{p} and \mathcal{L} be some other general linear space of codimension d . By using homotopy continuation, we can track the solution path $\mathbf{q}(t)$ defined by $\mathbf{q}(1) = \mathbf{p}$ and

$$\begin{aligned} \mathbf{c}(\mathbf{q}(t)) - \mathbf{c}(\mathbf{p}) &= 0 \\ \mathbf{q}(t) &\in t \cdot \mathcal{L}_{\mathbf{p}} + (1 - t) \cdot \mathcal{L}. \end{aligned} \tag{10}$$

This yields the point $\mathbf{q}(0)$ which is also a generic point in $V_{\mathbf{p}}$.

One can easily compute other points in this same fiber $V_{\mathbf{p}}$ by repeating with a different linear space \mathcal{L} and sample other fibers by repeating the process with a different generic point \mathbf{p} .

With the aim of computing globally identifiable functions, sample points in every irreducible component of $\mathbf{c}(\mathbf{q}) - \mathbf{c}(\mathbf{p}) = 0$ are needed. In this case, one simply constructs an identifiable system by restricting the parameters to a general linear space of codimension d and applying the techniques from §3.1 to the resulting system. That is, if $\mathbf{p} \in \mathbb{C}^{m_1}$, we take a general affine linear mapping $\mathbf{b} : \mathbb{C}^{m_1-d} \rightarrow \mathbb{C}^{m_1}$ so that $\mathbf{c}(\mathbf{b}(\hat{\mathbf{q}})) - \mathbf{c}(\mathbf{b}(\hat{\mathbf{p}})) = 0$ has finitely many solutions for generic $\hat{\mathbf{p}}$, say $\mathbf{q}_1 = \mathbf{b}(\hat{\mathbf{q}}_1), \dots, \mathbf{q}_k = \mathbf{b}(\hat{\mathbf{q}}_k)$, i.e., the model with parameters $\mathbf{p} = \mathbf{b}(\hat{\mathbf{p}})$ is identifiable over \mathbb{C}^{m_1-d} with identifiability degree k . Applying the slice moving described above, one can sample points in all components of the fiber over \mathbf{p} using the points $\mathbf{q}_1, \dots, \mathbf{q}_k$.

Example 4.2. Reconsider (3) in Ex. 2.3 for which \mathbf{c} shows the model has $d = 2$ dimensions of unidentifiability. For illustration, with $\mathbf{p} = (1, 2, 3, 4, 5, 6, 7)$, we can take

$$\mathcal{L}_{\mathbf{p}} = \{k_{01} - k_{02} + k_{03} - k_{12} + k_{13} - k_{21} + k_{32} = 4, 2k_{02} - k_{01} + 2k_{03} + k_{12} - k_{13} - 2k_{21} + 2k_{32} = 10\} \quad (11)$$

and

$$\mathcal{L} = \left\{ \begin{array}{l} k_{01} + k_{02}(3 - i) + k_{03}(-3 + 2i) + k_{12}(1 + i) - k_{13}(2 + 2i) + k_{21}(2 - i) - 2k_{32} = 1 \\ k_{01}(1 - 3i) - 3ik_{02} + k_{03}(-2 + 2i) - 2k_{12} - ik_{13} + k_{21}(3 + 2i) - ik_{32} = 1 \end{array} \right\} \quad (12)$$

where $i = \sqrt{-1}$. Tracking the path defined by (10) yields the endpoint (to four decimal places):

$$(0.6709 - 2.1940i, 3.6921 + 2.5919i, 2.8774 + 0.5068i, 3.3852 - 1.1735i, 5.1226 - 0.5068i, 6.3291 + 2.1940i, 5.9227 - 1.4185i). \quad (13)$$

Hence, we immediately see that each parameter itself is an unidentifiable function.

If, for illustration, we take the affine linear mapping $\mathbf{b} : \mathbb{C}^5 \rightarrow \mathbb{C}^7$ defined by

$$\mathbf{b}(\hat{\mathbf{p}}) = \begin{bmatrix} \hat{\mathbf{p}}_1 \\ \hat{\mathbf{p}}_2 \\ \hat{\mathbf{p}}_3 \\ \hat{\mathbf{p}}_4 \\ \hat{\mathbf{p}}_5 \\ \hat{\mathbf{p}}_1 + 3\hat{\mathbf{p}}_2 - \hat{\mathbf{p}}_3 - 3\hat{\mathbf{p}}_4 + 2\hat{\mathbf{p}}_5 + 4 \\ 2\hat{\mathbf{p}}_1 + 3\hat{\mathbf{p}}_2 + 5\hat{\mathbf{p}}_3 + \hat{\mathbf{p}}_4 - 3\hat{\mathbf{p}}_5 - 5 \end{bmatrix}$$

the resulting model is identifiable with identifiability degree 8 and the following 7 other points corresponding with $\mathbf{b}(1, 2, 3, 4, 5) = (1, 2, 3, 4, 5, 6, 7)$:

$$\begin{aligned} &(9.2814, -10.3208, 10.7201, -10.52, -2.7201, -2.2814, 33.8409), \\ &(108.0762, -66.9431, 13.0118, -0.23744, -0.0118, -101.0762, 75.1805), \\ &(2.4938, 0.3612, 4.3645, 5.326, 8.6355, 4.5062, 2.3128), \\ &(52.0709, -31.4763, 8.1035, -0.5325, -0.1035, -45.0709, 45.0087), \\ &(8.6814, -13.22, 14.5081, -14.2737, -1.5081, -1.6814, 35.4937), \\ &(-9.615, 5.8203, 5.6325, 1.4445, 7.3675, 16.615, 0.7352), \\ &(-13.625, 9.5193, 3.6057, 1.1636, 4.3943, 20.625, 2.317). \end{aligned} \quad (14)$$

Thus, we have computed at least one point in each irreducible component of the fiber over \mathbf{p} .

Without input-output equations, one simply uses a truncated system F_r as described in §2.2 to perform the same computations. The only potential issues were addressed in §3.2, namely reduction to the case that the state variables are generically zero-dimensional over the parameter-input-output space and restricting to the irreducible components which have the same input-output. The latter is accomplished by simply ignoring the components which have different input-output values.

Example 4.3. To illustrate moving on an irreducible component, we describe the setup to yield the same corresponding endpoint in (13). To that end, following Ex. 2.6, we utilize F_7 . Starting with parameter values $\mathbf{p} = (1, 2, 3, 4, 5, 6, 7)$, the structure of F_7 makes it trivial to generate general input, output, and state variables satisfying $F_7 = 0$, i.e., randomly selecting input \mathbf{U} and initial conditions \mathbf{x}_0 for the state variables trivially yields the values of the other state variables $\mathbf{x}_1, \dots, \mathbf{x}_8$ and output \mathbf{Y} . Then, by holding the input \mathbf{U} and output \mathbf{Y} fixed, we track along the solution path where the variables consist of the model parameters and the state variables defined by $F_7 = 0$ that deforms $\mathcal{L}_{\mathbf{p}}$ in (11) to \mathcal{L} in (12). The resulting endpoint corresponds with the endpoint in (13).

4.2 Functions from samples

From the ability to sample points described in §4.1, we can reconstruct identifiable functions in a given finite-dimensional vector space of functions, say $\mathcal{F} = \text{span}\{f_1, \dots, f_j\}$. Following Prop. 2.2, an identifiable function $f \in \mathcal{F}$ is constant on irreducible components of generic fibers of \mathbf{c} , which corresponds with computing null spaces of linear equations described as follows.

We can express every $f \in \mathcal{F}$ as $f = \sum_{i=1}^j a_i f_i$ where $\mathbf{a} = (a_1, \dots, a_j) \in \mathbb{C}^j$. If \mathbf{p} is a generic value of the parameters, using §4.1, we can compute a generic $\mathbf{q}_{\mathbf{p}}$ in the same irreducible component $V_{\mathbf{p}}$. Hence, the condition $f(\mathbf{q}_{\mathbf{p}}) = f(\mathbf{p})$ imposes a linear constraint on \mathbf{a} , namely

$$\begin{bmatrix} f_1(\mathbf{q}_{\mathbf{p}}) - f_1(\mathbf{p}) & \cdots & f_j(\mathbf{q}_{\mathbf{p}}) - f_j(\mathbf{p}) \end{bmatrix} \cdot \mathbf{a} = 0.$$

One option is to keep imposing more such conditions by selecting other general values of \mathbf{p} with corresponding $\mathbf{q}_{\mathbf{p}}$. The dimension of the null space reduces by one monotonically with each new condition until it reaches the dimension of the linear span of the identifiable functions in \mathcal{F} .

Alternatively, for computing identifiable functions with integer coefficients, i.e., $\mathbf{a} \in \mathbb{Z}^j$, one general point is enough via exactness recovery methods [2].

Example 4.4. Let $\mathcal{F} = \text{span}\{k_{01}, k_{02}, k_{03}, k_{12}, k_{13}, k_{21}, k_{32}\}$, $\mathbf{p} = (1, 2, 3, 4, 5, 6, 7)$, and $\mathbf{q}_{\mathbf{p}}$ as in (13). Then, integer solutions to $(\mathbf{q}_{\mathbf{p}} - \mathbf{p}) \cdot \mathbf{a} = 0$ computed using [2] correspond to:

$$k_{01} + k_{21}, \quad k_{03} + k_{13}, \quad k_{02} + k_{12} + k_{32}.$$

Alternatively, one can sample $V_{\mathbf{p}}$ for five general values of \mathbf{p} and observe that the first four impose a new linear constraint on the coefficients \mathbf{a} while the fifth one is redundant. This shows that there is a three-dimensional linear space of identifiable functions in \mathcal{F} spanned by the three linear functions above.

We bring all methods of §4 together in the following brief high-level pseudocode.

Method 5: Computing identifiable functions via sampling

Input: Input-output equation coefficients $\mathbf{c}(\mathbf{q})$, depending on parameters $\mathbf{q} \in \mathbb{C}^{m_1}$ (if available), else the truncated system F_r for some r and a basis f_1, \dots, f_j for a linear space of polynomials \mathcal{F} of interest.

Output: Identifiable functions in \mathcal{F} .

Choose random, complex values \mathbf{p} of parameters \mathbf{q} .

Compute a point on each irreducible component of $\mathbf{c}^{-1}(\mathbf{c}(\mathbf{p}))$ using either \mathbf{c} or F_r .

Use homotopy sampling to additional points on each irreducible component.

Use the sample points together with exactness recovery methods to find all identifiable functions in \mathcal{F} .

Return all discovered identifiable functions.

Globally identifiable functions are computed by simply adding the condition that the function takes the same constant value on *all* irreducible components of general fibers which are sampled using the methods of §4.1. Since globally identifiable functions are a subset of the identifiable functions, one need only search inside of the space of identifiable functions in \mathcal{F} .

Example 4.5. From the seven points in (14) corresponding with $\mathbf{p} = (1, 2, 3, 4, 5, 6, 7)$, we see that $k_{01} + k_{21}$ is globally identifiable (always taking the value 7 on these eight points) whereas $k_{03} + k_{13}$ and $k_{02} + k_{12} + k_{32}$ are not globally identifiable. However, from the sample points, it is easy to see that their sum, namely $k_{02} + k_{03} + k_{12} + k_{13} + k_{32}$, is globally identifiable.

The selection of the test space \mathcal{F} is a user-defined input and is based on the structure of the identifiable functions of interest, e.g., linear functions, polynomials of low degree, or linear span of rational monomials where the numerator and denominator have low degree.

5 Examples

We now demonstrate our methods on two larger examples. Throughout the paper, for illustrative purposes, the examples presented typically selected small integer values for random numbers. In practice, including the following examples, we select random complex numbers.

Example 5.1. The following is a 4-compartment model from [26, Ex. 6.3]:

$$\begin{aligned} \dot{x}_1 &= a_{11}x_1 + a_{12}x_2 + u \\ \dot{x}_2 &= a_{21}x_1 + a_{22}x_2 + a_{23}x_3 \\ \dot{x}_3 &= a_{33}x_3 + a_{34}x_4 \\ \dot{x}_4 &= a_{42}x_2 + a_{43}x_3 + a_{44}x_4 \\ y &= x_1. \end{aligned}$$

This model, which has parameters $\mathbf{p} = (a_{11}, a_{12}, a_{21}, a_{22}, a_{23}, a_{33}, a_{34}, a_{42}, a_{43}, a_{44})$, input $u(t)$, state variables $x_1(t), x_2(t), x_3(t), x_4(t)$, and output $y(t)$, does not fit the criteria presented in [26] for computing identifiable functions. Nonetheless, the method provided in [26, 27] is able to

compute the input-output equations where the set $\mathbf{c} : \mathbb{C}^{10} \rightarrow \mathbb{C}^7$ of coefficients is

$$\begin{aligned} & a_{11}a_{23}a_{34}a_{42} + a_{12}a_{21}a_{34}a_{43} - a_{11}a_{22}a_{34}a_{43} - a_{12}a_{21}a_{33}a_{44} + a_{11}a_{22}a_{33}a_{44} \\ & a_{12}a_{21}a_{33} - a_{11}a_{22}a_{33} - a_{23}a_{34}a_{42} + a_{11}a_{34}a_{43} + a_{22}a_{34}a_{43} + a_{12}a_{21}a_{44} - a_{11}a_{22}a_{44} - a_{11}a_{33}a_{44} - a_{22}a_{33}a_{44} \\ & - a_{12}a_{21} + a_{11}a_{22} + a_{11}a_{33} + a_{22}a_{33} - a_{34}a_{43} + a_{11}a_{44} + a_{22}a_{44} + a_{33}a_{44} \\ & - a_{11} - a_{22} - a_{33} - a_{44} \\ & a_{23}a_{34}a_{42} - a_{22}a_{34}a_{43} + a_{22}a_{33}a_{44} \\ & - a_{22}a_{33} + a_{34}a_{43} - a_{22}a_{44} - a_{33}a_{44} \\ & a_{22} + a_{33} + a_{44}. \end{aligned}$$

Using Prop 2.2, the model is unidentifiable with 4 dimensions of unidentifiability. Therefore, to solve Problem 2.11, we need to compute 6 algebraically independent identifiable functions.

We utilize the method of §4.1 to sample and that of §4.2 to construct the identifiable functions. For example, sampling yields the following two values of the parameters (rounded to four decimal places) so that every identifiable function must take the same value on both:

a_{11}	$-0.6690 - 0.1758i$	$-0.6690 - 0.1758i$
a_{12}	$-0.1669 + 0.3165i$	$1.3705 - 0.4117i$
a_{21}	$2.3433 + 0.6225i$	$-0.5219 + 0.3086i$
a_{22}	$-0.6286 - 0.1868i$	$-0.6286 - 0.1868i$
a_{23}	$0.4005 - 0.5144i$	$2.5585 + 0.5746i$
a_{33}	$2.1248 - 0.6011i$	$0.2095 - 0.4521i$
a_{34}	$1.1295 - 0.8604i$	$0.8611 + 0.5272i$
a_{42}	$-0.4210 + 0.6785i$	$0.2734 - 0.0567i$
a_{43}	$-1.1126 - 0.0416i$	$-0.1132 - 0.7724i$
a_{44}	$-0.6880 + 0.3317i$	$1.2273 + 0.1827i$

In particular, we immediately see that both $f_1 = a_{11}$ and $f_2 = a_{22}$ are identifiable. Applying §4.2 to the space of linear forms also yields the identifiable linear function $f_3 = a_{33} + a_{44}$.

Considering the space of polynomials of degree at most 2 which are algebraically independent of f_1, f_2, f_3 yields $f_4 = a_{12}a_{21}$ and $f_5 = a_{33}a_{44} - a_{34}a_{43}$.

Finally, the space of polynomials of degree at most 3 which are algebraically independent of f_1, \dots, f_5 yields $f_6 = a_{23}a_{34}a_{42}$.

To show that f_1, \dots, f_6 are actually globally identifiable, we use the approach in §4.1 to sample points from every irreducible component. The result of this process is that a generic fiber only has one irreducible component thereby showing global identifiability. We could also have used Defn. 2.9 to show global identifiability. This is demonstrated by the following:

$$\begin{aligned} f_1 &= -(c_4 + c_7) \\ f_2 &= \frac{c_6^2 + c_6c_7^2 + c_4c_6c_7 + c_3c_6 + c_5c_7 + c_1 + c_4c_5}{c_2 + c_5 + c_4c_6 + c_6c_7} \\ f_3 &= -\frac{c_2 + c_5 + c_4c_6 + c_6c_7}{c_7^2 + c_4c_7 + c_3 + c_6} \\ f_4 &= -(c_7^2 + c_4c_7 + c_3 + c_6) \\ f_5 &= \frac{c_1 + c_4c_5 + c_5c_7}{c_7^2 + c_4c_7 + c_3 + c_6} \\ f_6 &= -\frac{c_1^2 + 2c_1c_4c_5 + c_1c_4c_6c_7 + 2c_1c_5c_7 + c_1c_6^2 + c_1c_6c_7^2 + c_3c_1c_6 + c_4^2c_5^2 + c_4c_5^2c_7}{c_2c_3 + c_2c_6 + c_3c_5 + c_5c_6 + c_2c_7^2 + c_4c_6^2 + c_5c_7^2 + c_6^2c_7 + c_6c_7^3 + c_2c_4c_7 + c_3c_4c_6 + c_3c_6c_7 + c_4c_5c_7 + 2c_4c_6c_7^2 + c_4^2c_6c_7} \end{aligned}$$

Example 5.2. The following is a model from biochemical reaction network theory for the mitogen-activated protein kinase (MAPK) pathway [22] which is part of a molecular signaling network that governs the growth, proliferation, differentiation, and survival of many cell types:

$$\begin{aligned}
K\dot{S}_{00} &= -a_{00}KS_{00} + b_{00}KS_{00} + \gamma_{0100}FS_{01} + \gamma_{1000}FS_{10} + \gamma_{1100}FS_{11} \\
K\dot{S}_{01} &= -a_{01}KS_{01} + b_{01}KS_{01} + c_{0001}KS_{00} - \alpha_{01}FS_{01} + \beta_{01}FS_{01} + \gamma_{1101}FS_{11} \\
K\dot{S}_{10} &= -a_{10}KS_{10} + b_{10}KS_{10} + c_{0010}KS_{00} - \alpha_{10}FS_{10} + \beta_{10}FS_{10} + \gamma_{1110}FS_{11} \\
F\dot{S}_{01} &= -\alpha_{11}FS_{11} + \beta_{11}FS_{11} + c_{0111}KS_{01} + c_{1011}KS_{10} + c_{0011}KS_{00} \\
F\dot{S}_{10} &= a_{00}KS_{00} - (b_{00} + c_{0001} + c_{0010} + c_{0011})KS_{00} \\
F\dot{S}_{11} &= a_{01}KS_{01} - (b_{01} + c_{0111})KS_{01} \\
\dot{K} &= a_{10}KS_{10} - (b_{10} + c_{1011})KS_{10} \\
\dot{F} &= \alpha_{01}FS_{01} - (\beta_{01} + \gamma_{0100})FS_{01} \\
\dot{S}_{00} &= \alpha_{10}FS_{10} - (\beta_{10} + \gamma_{1000})FS_{10} \\
\dot{S}_{01} &= \alpha_{11}FS_{11} - (\beta_{11} + \gamma_{1101} + \gamma_{1110} + \gamma_{1100})FS_{11} \\
\dot{S}_{10} &= -a_{00}KS_{00} + (b_{00} + c_{0001} + c_{0010} + c_{0011})KS_{00} - a_{01}KS_{01} \\
&\quad + (b_{01} + c_{0111})KS_{01} - a_{10}KS_{10} + (b_{10} + c_{1011})KS_{10} \\
\dot{S}_{11} &= -\alpha_{01}FS_{01} + (\beta_{01} + \gamma_{0100})FS_{01} - \alpha_{10}FS_{10} + (\beta_{10} + \gamma_{1000})FS_{10} - \alpha_{11}FS_{11} \\
&\quad + (\beta_{11} + \gamma_{1101} + \gamma_{1110} + \gamma_{1100})FS_{11}.
\end{aligned}$$

This model has 12 state variables

$$KS_{00}(t), KS_{01}(t), KS_{10}(t), FS_{01}(t), FS_{10}(t), FS_{11}(t), K(t), F(t), S_{00}(t), S_{01}(t), S_{10}(t), S_{11}(t)$$

and 22 parameters

$$\begin{aligned}
&a_{00}, \quad a_{01}, \quad a_{10}, \quad b_{00}, \quad b_{01}, \quad b_{10}, \quad c_{0001}, \quad c_{0010}, \quad c_{0011}, \quad c_{0111}, \quad c_{1011}, \\
&\alpha_{01}, \quad \alpha_{10}, \quad \alpha_{11}, \quad \beta_{01}, \quad \beta_{10}, \quad \beta_{11}, \quad \gamma_{0100}, \quad \gamma_{1000}, \quad \gamma_{1100}, \quad \gamma_{1101}, \quad \gamma_{1110}.
\end{aligned}$$

We will consider several different cases of what is measured as output. In all of our examples, we attempted to first compute input-output equations using differential elimination via the command *RosenfeldGroebner* in *Maple* [23]. In all of our attempts, the differential elimination failed to terminate meaning that we will just utilize the model equations in the following.

First, for taking the standard 6 measurable outputs:

$$y_1 = K, \quad y_2 = F, \quad y_3 = S_{00}, \quad y_4 = S_{01}, \quad y_5 = S_{10}, \quad y_6 = S_{11},$$

the following table, computed in about a minute on a single processor, shows that the resulting model is identifiable:

r	$\text{corank}_0 JF_r$	$\text{corank}_{22} JF_r$	d_r
0	28	6	22
1	23	1	22
2	18	0	18
3	13	0	13
4	8	0	8
5	3	0	3
6	0	0	0

For comparison of methods, neither DAISY [7, 35] nor COMBOS [28] finished the identifiability computations for this model after running for 24 hours. To the best of our knowledge, this is the first successful implementation of a structural identifiability test for this model.

Second, if we adjust the model so that we only take the following 2 measurable outputs:

$$y_1 = K, \quad y_2 = F,$$

the following table shows that the resulting model is still identifiable:

r	$\text{corank}_0 JF_r$	$\text{corank}_{22} JF_r$	d_r
0	32	10	22
1	30	8	22
2	28	6	22
3	26	4	22
4	24	2	22
5	22	0	22
6	20	0	20
7	18	0	18
8	16	0	16
9	14	0	14
10	12	0	12
11	10	0	10
12	8	0	8
13	6	0	6
14	4	0	4
15	2	0	2
16	0	0	0

Third, if we take the following 4 measurable outputs:

$$y_1 = S_{00}, \quad y_2 = S_{01}, \quad y_3 = S_{10}, \quad y_4 = S_{11},$$

the following table shows that the resulting model is still identifiable:

r	$\text{corank}_0 JF_r$	$\text{corank}_{22} JF_r$	d_r
0	30	8	22
1	26	4	22
2	22	0	22
3	18	0	18
4	14	0	14
5	10	0	10
6	6	0	6
7	2	0	2
8	0	0	0

Finally, we consider 10 new mixing parameters, namely

$$ms_{00}, \quad mks_{00}, \quad ms_{01}, \quad mks_{01}, \quad mfs_{01}, \quad ms_{10}, \quad mks_{10}, \quad mfs_{10}, \quad ms_{11}, \quad mfs_{11},$$

with the following 4 measurable outputs:

$$\begin{aligned} y_1 &= ms_{00}S_{00} + mks_{00}KS_{00} \\ y_2 &= ms_{01}S_{01} + mks_{01}KS_{01} + mfs_{01}FS_{01} \\ y_3 &= ms_{10}S_{10} + mks_{10}KS_{10} + mfs_{10}FS_{10} \\ y_4 &= ms_{11}S_{11} + mfs_{11}FS_{11}. \end{aligned}$$

The following table shows that the resulting model, which has a total of 32 parameters, is unidentifiable with one dimension of unidentifiability:

r	$\text{corank}_0 JF_r$	$\text{corank}_{32} JF_r$	d_r
0	40	8	32
1	36	4	32
2	32	0	32
3	28	0	28
4	24	0	24
5	20	0	20
6	16	0	16
7	12	0	12
8	8	0	8
9	4	0	4
10	1	0	1
11	1	0	1

Using the results from §4, we can observe from sampling that each irreducible component of a general fiber is simply a line and the following 16 parameters are all identifiable:

$$b_{00}, b_{01}, b_{10}, c_{0001}, c_{0010}, c_{0011}, c_{0111}, c_{1011}, \beta_{01}, \beta_{10}, \beta_{11}, \gamma_{0100}, \gamma_{1000}, \gamma_{1100}, \gamma_{1101}, \gamma_{1110}$$

meaning $a_{00}, a_{01}, a_{10}, \alpha_{01}, \alpha_{10}, \alpha_{11}$ and the 10 mixing parameters are unidentifiable. In fact, no nonconstant linear function in these 16 unidentifiable parameters is identifiable.

6 Conclusion

In this article, we considered the problems of determining the identifiability of an ODE model, computing the identifiability degree in the case that the model is identifiable and identifiable functions in the case that the model is unidentifiable. To summarize, the results of this article include numerical methods for the following:

1. compute the dimension of unidentifiability with or without input-output equations;
2. for identifiable models, compute the identifiability degree with or without input-output equations using basic homotopy continuation or monodromy loops;
3. for unidentifiable models, compute identifiable and globally identifiable functions inside of a linear family of functions with or without input-output equations.

These methods were illustrated on several examples, including the first known structural identifiability result for MAPK in Example 5.2.

In the future, we hope to apply similar numerical algebraic geometry methods to other areas in biological modelling, such as controllability, observability, and indistinguishability.

References

- [1] S. Audoly, G. Bellu, L. D’Angiò, M. P. Saccomani, and C. Cobelli, Global identifiability of nonlinear models of biological systems, *IEEE Transactions on Biomedical Engineering* **48** (2001) 55-65.
- [2] D.J. Bates, J.D. Hauenstein, T. McCoy, C. Peterson, and A.J. Sommese, Recovering exact results from inexact numerical data in algebraic geometry, *Experimental Math.* **22**(1):38–50, 2013.
- [3] D.J. Bates, J.D. Hauenstein, A.J. Sommese, and C.W. Wampler, Bertini: Software for numerical algebraic geometry, Available for download at bertini.nd.edu/.
- [4] D.J. Bates, J.D. Hauenstein, A.J. Sommese, and C.W. Wampler, *Numerically solving polynomial systems with Bertini*, SIAM, 2013.
- [5] D. J. Bearup, N. D. Evans, and M. J. Chappell, The input-output relationship approach to structural identifiability analysis, *Computer Methods and Programs in Biomedicine* **109** (2013) 171-181.
- [6] R. Bellman and K. J. Astrom, On structural identifiability, *Math. Biosci.* **7** (1970) 329-339.
- [7] G. Bellu, M. P. Saccomani, S. Audoly, and L. D’Angiò, DAISY: A new software tool to test global identifiability of biological and physical systems, *Computers in Biology and Medicine* **88** (2007) 52-61.
- [8] M. Berman and R. Schoenfeld, Invariants in Experimental Data on Linear Kinetics and the Formulation of Models, *Journal of Applied Physics* **27** (1956) 1361-1370.
- [9] M. Berman, E. Shahn, and M. F. Weiss, Some Formal Approaches to the Analysis of Kinetic Data in Terms of Linear Compartmental Systems, *Biophysical Journal* **2** (1962) 289-316.
- [10] F. Boulier, Differential Elimination and Biological Modelling, *Radon Series Comp. Appl. Math* **2** (2007) 111-139.
- [11] D.A. Brake, J.D. Hauenstein, and A.C. Liddell, Decomposing solution sets of polynomial systems using derivatives, *Mathematical Software – ICMS 2016, LNCS* **9725** (2016) 127-135.
- [12] M. J. Chappell and R. N. Gunn, A procedure for generating locally identifiable reparameterisations of unidentifiable non-linear systems by the similarity transformation approach, *Math. Biosci.* **148** (1998) 21-41.
- [13] J. J. DiStefano III, *Dynamic Systems Biology Modeling and Simulation*. Elsevier, London, 2014.
- [14] T. Duff, C. Hill, A. Jensen, K. Lee, A. Leykin, and J. Sommars, Solving polynomial systems via homotopy continuation and monodromy, [arXiv 1609.08722](https://arxiv.org/abs/1609.08722).
- [15] N. D. Evans and M. J. Chappell, Extensions to a procedure for generating locally identifiable reparameterisations of unidentifiable systems, *Math. Biosci.* **168** (2000) 137-159.

- [16] N. D. Evans, H. Moyse, D. Lowe, D. Briggs, R. Higgins, D. Mitchell, D. Zehnder, and M. J. Chappell, Structural identifiability of surface binding reactions involving heterogeneous analyte: application to surface plasmon resonance experiments, *Automatica* **49** (2012) 48-57.
- [17] J.D. Hauenstein, L. Oeding, G. Ottaviani, and A.J. Sommese. Homotopy techniques for tensor decomposition and perfect identifiability. To appear in *J. Reine Angew. Math.*
- [18] J.D. Hauenstein and J. Rodriguez, Multiprojective witness sets and a trace test. [arXiv 1507.07069](https://arxiv.org/abs/1507.07069).
- [19] J.D. Hauenstein and A.J. Sommese, Witness sets of projections, *Appl. Math. and Comput.* **217(7)** (2010) 3349-3354.
- [20] J. Karlsson, M. Anguelova, and M. Jirstrand, An Efficient Method for Structural Identifiability Analysis of Large Dynamic Systems, *IFAC Proceedings Volumes* **45 (16)** (2012) 941-946.
- [21] L. Ljung and T. Glad, On global identifiability for arbitrary model parameterization, *Automatica* **30(2)** (1994) 265-276.
- [22] A. K. Manrai and J. Gunawardena, The geometry of multisite phosphorylation, *Biophys. J.* **95** (2008) 5533-5543.
- [23] Maple documentation. <http://www.maplesoft.com/support/help/maple/view.aspx?path=DifferentialAlgebra>
- [24] N. Meshkat, M. Eisenberg, and J. J. DiStefano III, An algorithm for finding globally identifiable parameter combinations of nonlinear ODE models using Gröbner Bases, *Math. Biosci.* **222** (2009) 61-72.
- [25] N. Meshkat, C. Anderson, and J. J. DiStefano III, Alternative to Ritt's Pseudodivision for finding the input-output equations of multi-output models, *Math. Biosci.* **239** (2012) 117-123.
- [26] N. Meshkat and S. Sullivant, Identifiable reparameterizations of linear compartment models, *J. Symb. Comp.* **63** (2014) 46-67.
- [27] N. Meshkat, S. Sullivant, and M. Eisenberg, Identifiability results for several classes of linear compartment models. *Bull. of Math. Bio.* **77(8)** (2015) 1620-1651.
- [28] N. Meshkat, C. E. Kuo, J. J. DiStefano III, On Finding and Using Identifiable Parameter Combinations in Nonlinear Dynamic Systems Biology Models and COMBOS: A Novel Web Implementation, *PLoS ONE* **9(10)** (2014) e110261.
- [29] H. Miao, X. Xia, A. Perelson, H. Wu, On identifiability of nonlinear ODE models and applications in viral dynamics, *SIAM Review* **53(1)** (2011) 3-39.
- [30] R.J. Mulholland and M.S. Keener, Analysis of linear compartment models for ecosystems, *Journal of Theoretical Biology* **44** (1974) 105-116.

- [31] F. Ollivier, Le probleme de l'identifiabilite structurelle globale: etude theoretique, methodes effectives and bornes de complexite, PhD thesis, Ecole Polytechnique, 1990.
- [32] H. Pohjanpalo, System identifiability based on the power series expansion of the solution, *Math. Biosci.* **41** (1978) 21-33.
- [33] G.J. Reid, P. Lin, and A.D. Wittkopf, Differential elimination-completion algorithms for DAE and PDAE, *Stud. Appl. Math.*, **106**(1) (2001) 1-45.
- [34] M.P. Saccomani, S. Audoly, and L. D'Angiò, Parameter identifiability of nonlinear systems: the role of initial conditions, *Automatica* **39** (2003) 619-632.
- [35] M.P. Saccomani, S. Audoly, G. Bellu, and L. D'Angiò, Examples of testing global identifiability of biological and biomedical models with the DAISY software, *Computers in Biology and Medicine* **40** (2010) 402-407.
- [36] M.P. Saccomani, S. Audoly, G. Bellu, and L. D'Angiò, A new differential algebra algorithm to test identifiability of nonlinear systems with given initial conditions, *Proceedings of the 40th IEEE Conference on Decision and Control* (2001) 3108-3113.
- [37] B. Shiffman and A.J. Sommese, *Vanishing theorems on complex manifolds*, Progress in Mathematics, vol. 56, Birkhäuser Boston, Inc., Boston, MA, 1985.
- [38] A.J. Sommese, J. Verschelde, and C.W. Wampler, Symmetric functions applied to decomposing solution sets of polynomial systems, *SIAM J. Numer. Anal.* **40**(6) (2002) 2026-2046.
- [39] A.J. Sommese, J. Verschelde, and C.W. Wampler. Using monodromy to decompose solution sets of polynomial systems into irreducible components. *NATO Science Series* **36** (2001) 297-315.
- [40] A.J. Sommese and C.W. Wampler, *The numerical solution of polynomial systems arising in engineering and science*, World Scientific, 2005.
- [41] J.G. Wagner, History of pharmacokinetics, *Pharmacology & Therapeutics* **12** (1981) 537-562.
- [42] E. Walter, *Identifiability of state space models*, *Lecture Notes in Biomathematics* **46**, Springer, Berlin, 1982.